



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:24 AM UTC

PDB ID : 3Q32 / pdb_00003q32
Title : Structure of Janus kinase 2 with a pyrrolotriazine inhibitor
Authors : Sack, J.S.
Deposited on : 2010-12-21
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	6% 72% 22% ...
1	B	301	5% 75% 19% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5039 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase JAK2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	P	S	0	0	0
			2425	1536	421	452	2	14			
1	B	296	Total	C	N	O	P	S	0	0	0
			2454	1553	429	455	2	15			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	838	MET	-	initiating methionine	UNP O60674
A	1133	HIS	-	expression tag	UNP O60674
A	1134	HIS	-	expression tag	UNP O60674
A	1135	HIS	-	expression tag	UNP O60674
A	1136	HIS	-	expression tag	UNP O60674
A	1137	HIS	-	expression tag	UNP O60674
A	1138	HIS	-	expression tag	UNP O60674
B	838	MET	-	initiating methionine	UNP O60674
B	1133	HIS	-	expression tag	UNP O60674
B	1134	HIS	-	expression tag	UNP O60674
B	1135	HIS	-	expression tag	UNP O60674
B	1136	HIS	-	expression tag	UNP O60674
B	1137	HIS	-	expression tag	UNP O60674
B	1138	HIS	-	expression tag	UNP O60674

- Molecule 2 is 2-(2,6-difluoro-4-methoxyphenyl)-1-(4-{4-[(3-methyl-1H-pyrazol-5-yl)amino]pyrrolo[2,1-f][1,2,4]triazin-2-yl}piperazin-1-yl)ethanone (CCD ID: J2I) (formula: C₂₃H₂₄F₂N₈O₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 35	C 23	F 2	N 8	O 2	0	0
2	B	1	Total 35	C 23	F 2	N 8	O 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	42	Total O 42 42	0	0
3	B	48	Total O 48 48	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein kinase JAK2



- Molecule 1: Tyrosine-protein kinase JAK2



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	110.59Å 110.59Å 69.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.46 – 2.50 49.46 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.46-2.50) 100.0 (49.46-2.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.36 (at 2.51Å)	Xtriage
Refinement program	BUSTER 2.9.7, autoBUSTER	Depositor
R, R_{free}	0.178 , 0.210 (Not available) , 0.222	Depositor DCC
R_{free} test set	649 reflections (2.21%)	wwPDB-VP
Wilson B-factor (Å ²)	43.7	Xtriage
Anisotropy	0.554	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.046 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5039	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: J2I, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	1/2443 (0.0%)	1.42	25/3287 (0.8%)
1	B	0.85	2/2473 (0.1%)	1.43	20/3326 (0.6%)
All	All	0.86	3/4916 (0.1%)	1.43	45/6613 (0.7%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1102	MET	SD-CE	-8.51	1.58	1.79
1	B	964	MET	SD-CE	-5.50	1.65	1.79
1	A	1102	MET	SD-CE	-5.07	1.66	1.79

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	923	ARG	CA-C-N	7.61	136.08	121.54
1	B	923	ARG	C-N-CA	7.61	136.08	121.54
1	B	894	ASP	CA-CB-CG	6.45	119.05	112.60
1	B	919	SER	CA-C-N	6.44	133.84	121.54
1	B	919	SER	C-N-CA	6.44	133.84	121.54
1	A	860	PHE	CA-CB-CG	6.29	120.09	113.80
1	A	948	ILE	N-CA-C	5.96	116.27	107.15
1	A	1120	ALA	CA-C-N	5.83	128.03	120.44
1	A	1120	ALA	C-N-CA	5.83	128.03	120.44
1	B	1027	THR	N-CA-C	5.78	117.66	111.36
1	A	1018	ILE	N-CA-C	5.75	117.17	110.62
1	B	841	PRO	CA-C-N	5.73	128.53	120.28
1	B	841	PRO	C-N-CA	5.73	128.53	120.28
1	B	908	ASP	CA-CB-CG	5.69	118.29	112.60
1	A	933	PRO	N-CA-C	5.67	120.94	113.86
1	A	861	GLY	N-CA-C	5.55	118.62	110.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1096	ASP	CA-C-N	5.51	127.93	120.38
1	A	1096	ASP	C-N-CA	5.51	127.93	120.38
1	B	1035	SER	N-CA-C	-5.44	104.99	111.69
1	A	948	ILE	CA-C-N	5.43	130.62	120.95
1	A	948	ILE	C-N-CA	5.43	130.62	120.95
1	B	945	LYS	N-CA-C	5.40	117.86	111.33
1	B	968	GLY	CA-C-N	5.40	128.05	120.28
1	B	968	GLY	C-N-CA	5.40	128.05	120.28
1	A	888	THR	N-CA-C	5.39	117.89	110.35
1	A	1017	PRO	CA-C-N	5.38	128.11	120.53
1	A	1017	PRO	C-N-CA	5.38	128.11	120.53
1	A	922	ARG	N-CA-C	-5.32	106.95	113.50
1	A	841	PRO	CA-C-N	5.30	129.08	120.60
1	A	841	PRO	C-N-CA	5.30	129.08	120.60
1	B	941	LEU	CA-C-N	5.27	127.29	120.44
1	B	941	LEU	C-N-CA	5.27	127.29	120.44
1	A	995	PHE	CA-C-N	5.24	125.89	120.03
1	A	995	PHE	C-N-CA	5.24	125.89	120.03
1	A	967	LEU	CA-C-N	5.21	126.66	120.14
1	A	967	LEU	C-N-CA	5.21	126.66	120.14
1	A	987	GLU	CB-CG-CD	5.18	121.41	112.60
1	A	960	ILE	CA-C-N	5.15	127.50	120.54
1	A	960	ILE	C-N-CA	5.15	127.50	120.54
1	B	893	ARG	CA-C-N	5.09	127.41	120.54
1	B	893	ARG	C-N-CA	5.09	127.41	120.54
1	A	1105	CYS	N-CA-C	-5.08	105.93	111.82
1	B	924	ASN	CA-C-N	5.06	128.28	121.05
1	B	924	ASN	C-N-CA	5.06	128.28	121.05
1	B	885	GLN	N-CA-C	-5.02	106.84	113.12

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2425	0	2396	21	0
1	B	2454	0	2426	17	0
2	A	35	0	24	4	0
2	B	35	0	24	1	0
3	A	42	0	0	0	0
3	B	48	0	0	0	0
All	All	5039	0	4870	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:932:LEU:H	2:B:2:J2I:HN24	1.15	0.92
1:A:932:LEU:H	2:A:1:J2I:HN24	1.28	0.78
1:A:917:CYS:HB3	1:A:926:LYS:HB2	1.77	0.66
1:A:1011:LYS:HD3	1:A:1011:LYS:H	1.62	0.64
1:A:1008:PTR:HE1	1:A:1010:VAL:HB	1.86	0.57
1:B:917:CYS:HB3	1:B:926:LYS:HB2	1.88	0.55
1:A:888:THR:HG23	1:A:891:HIS:HB2	1.89	0.55
1:B:1064:MET:HE2	1:B:1081:LEU:HD11	1.90	0.54
1:B:888:THR:HG22	1:B:890:GLU:H	1.74	0.52
1:A:935:GLY:HA2	2:A:1:J2I:C11	2.40	0.51
1:B:1071:GLY:HA2	1:B:1074:ILE:HD12	1.92	0.51
1:A:1002:PRO:HG2	1:A:1005:LYS:HB2	1.91	0.51
1:A:920:ALA:H	1:A:923:ARG:HB3	1.76	0.50
1:B:966:TYR:O	1:B:969:THR:HB	2.12	0.49
1:A:845:GLU:HB3	1:A:848:HIS:HD2	1.78	0.48
1:A:860:PHE:H	1:A:860:PHE:HD1	1.61	0.48
1:A:863:VAL:HG23	2:A:1:J2I:O33	2.14	0.47
1:B:1045:TYR:HB2	1:B:1102:MET:HE3	1.96	0.47
1:B:840:ASP:HB3	1:B:843:GLN:HG2	1.97	0.47
1:B:999:LYS:HD2	1:B:1008:PTR:HE1	1.97	0.46
1:A:886:HIS:HB3	1:A:888:THR:HG22	2.00	0.44
1:A:1098:ILE:HD11	1:A:1126:ILE:HG21	2.00	0.43
1:A:1050:TYR:CD2	1:B:1112:GLN:HG3	2.53	0.43
1:A:976:ASP:HB2	1:A:997:LEU:HD12	2.00	0.43
1:A:954:LEU:HB3	1:A:1127:ARG:CG	2.49	0.43
1:A:1064:MET:HE2	1:A:1081:LEU:HD11	2.01	0.42
1:B:867:ARG:HG2	1:B:869:ASP:HB2	2.02	0.42
2:A:1:J2I:C7	2:A:1:J2I:H20	2.49	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1062:MET:HE1	1:B:1074:ILE:HD13	2.01	0.42
1:A:1090:ARG:HG3	1:A:1099:TYR:HB2	2.02	0.41
1:B:923:ARG:HG2	1:B:924:ASN:N	2.36	0.41
1:B:1114:PRO:HG2	1:B:1119:LEU:HD21	2.03	0.41
1:A:1008:PTR:C	1:A:1008:PTR:HD1	2.51	0.41
1:B:948:ILE:CG2	1:B:953:LEU:HD13	2.51	0.41
1:A:848:HIS:CG	1:A:870:PRO:HA	2.55	0.41
1:B:923:ARG:O	1:B:924:ASN:HB3	2.21	0.40
1:B:967:LEU:HG	1:B:972:TYR:HB2	2.04	0.40
1:A:973:ILE:HG12	1:A:1001:LEU:HD13	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/301 (96%)	270 (93%)	13 (4%)	6 (2%)	5	9
1	B	292/301 (97%)	275 (94%)	9 (3%)	8 (3%)	4	6
All	All	581/602 (96%)	545 (94%)	22 (4%)	14 (2%)	4	8

All (14) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1011	LYS
1	A	1070	GLN
1	B	920	ALA
1	B	924	ASN
1	B	1070	GLN
1	A	925	LEU
1	A	1012	GLU
1	A	1071	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	1013	PRO
1	B	1015	GLU
1	B	1071	GLY
1	A	923	ARG
1	B	1016	SER
1	B	1014	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/273 (97%)	235 (89%)	30 (11%)	5	12
1	B	268/273 (98%)	240 (90%)	28 (10%)	7	14
All	All	533/546 (98%)	475 (89%)	58 (11%)	6	13

All (58) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	860	PHE
1	A	884	LEU
1	A	885	GLN
1	A	886	HIS
1	A	890	GLU
1	A	892	LEU
1	A	893	ARG
1	A	949	ASP
1	A	950	HIS
1	A	951	ILE
1	A	953	LEU
1	A	954	LEU
1	A	977	LEU
1	A	1001	LEU
1	A	1011	LYS
1	A	1012	GLU
1	A	1026	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1047	LEU
1	A	1051	ILE
1	A	1052	GLU
1	A	1065	ILE
1	A	1069	LYS
1	A	1072	GLN
1	A	1073	MET
1	A	1082	LEU
1	A	1096	ASP
1	A	1117	ARG
1	A	1119	LEU
1	A	1121	LEU
1	A	1125	GLN
1	B	839	ARG
1	B	842	THR
1	B	846	GLU
1	B	854	GLN
1	B	872	GLN
1	B	884	LEU
1	B	890	GLU
1	B	892	LEU
1	B	893	ARG
1	B	924	ASN
1	B	932	LEU
1	B	953	LEU
1	B	967	LEU
1	B	969	THR
1	B	989	ARG
1	B	1001	LEU
1	B	1003	GLN
1	B	1012	GLU
1	B	1026	LEU
1	B	1030	LYS
1	B	1033	VAL
1	B	1047	LEU
1	B	1065	ILE
1	B	1072	GLN
1	B	1073	MET
1	B	1082	LEU
1	B	1088	LEU
1	B	1121	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	854	GLN
1	A	872	GLN
1	A	874	ASN
1	A	1067	ASN
1	A	1084	ASN
1	A	1125	GLN
1	A	1129	ASN
1	B	843	GLN
1	B	885	GLN
1	B	1067	ASN
1	B	1112	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PTR	A	1008	1	15,16,17	1.50	4 (26%)	17,22,24	1.58	3 (17%)
1	PTR	A	1007	1	15,16,17	1.50	4 (26%)	17,22,24	1.23	1 (5%)
1	PTR	B	1007	1	15,16,17	1.27	0	17,22,24	1.24	2 (11%)
1	PTR	B	1008	1	15,16,17	1.35	3 (20%)	17,22,24	1.30	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	1008	1	-	1/10/11/13	0/1/1/1
1	PTR	A	1007	1	-	0/10/11/13	0/1/1/1
1	PTR	B	1007	1	-	0/10/11/13	0/1/1/1
1	PTR	B	1008	1	-	0/10/11/13	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1007	PTR	CE2-CZ	3.17	1.44	1.38
1	A	1008	PTR	CE2-CZ	2.93	1.44	1.38
1	B	1008	PTR	CE2-CZ	2.90	1.44	1.38
1	A	1008	PTR	CE2-CD2	2.67	1.43	1.38
1	A	1008	PTR	P-OH	-2.35	1.55	1.59
1	B	1008	PTR	CE2-CD2	2.30	1.42	1.38
1	A	1007	PTR	CE1-CZ	2.25	1.43	1.38
1	A	1008	PTR	CD2-CG	2.11	1.43	1.38
1	A	1007	PTR	CD1-CG	2.07	1.43	1.38
1	B	1008	PTR	P-O1P	2.06	1.56	1.50
1	A	1007	PTR	CB-CA	2.00	1.57	1.53

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1007	PTR	O3P-P-OH	3.88	116.80	105.32
1	A	1007	PTR	O3P-P-OH	3.38	115.31	105.32
1	A	1008	PTR	OH-CZ-CE2	3.27	129.01	119.22
1	A	1008	PTR	O3P-P-OH	3.17	114.68	105.32
1	A	1008	PTR	OH-CZ-CE1	-2.41	112.00	119.22
1	B	1007	PTR	P-OH-CZ	-2.29	115.73	123.88
1	B	1008	PTR	P-OH-CZ	-2.26	115.85	123.88
1	B	1008	PTR	CE2-CZ-CE1	-2.26	116.87	120.16
1	B	1008	PTR	CD1-CE1-CZ	2.14	122.18	119.73
1	B	1008	PTR	OH-P-O1P	2.14	116.62	109.48

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	1008	PTR	CZ-OH-P-O3P

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1008	PTR	2	0
1	B	1008	PTR	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	J2I	B	2	-	38,39,39	1.09	2 (5%)	45,56,56	1.95	10 (22%)
2	J2I	A	1	-	38,39,39	1.13	4 (10%)	45,56,56	2.31	15 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	J2I	B	2	-	-	3/18/28/28	0/5/5/5
2	J2I	A	1	-	-	2/18/28/28	0/5/5/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	J2I	C15-N25	3.20	1.41	1.34
2	A	1	J2I	C5-C13	2.71	1.42	1.38
2	A	1	J2I	C16-N30	2.53	1.40	1.35
2	A	1	J2I	C15-N25	2.51	1.39	1.34
2	B	2	J2I	C5-C13	2.45	1.42	1.38
2	A	1	J2I	C3-C9	2.15	1.41	1.37

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	J2I	C4-C10-C7	-5.87	120.08	124.45
2	A	1	J2I	C3-C9-C7	-5.74	120.18	124.45
2	A	1	J2I	C13-N31-C14	5.33	133.84	123.72
2	B	2	J2I	C4-C10-C7	-5.24	120.55	124.45
2	B	2	J2I	C13-N31-C14	5.03	133.28	123.72
2	B	2	J2I	C3-C9-C7	-4.65	120.99	124.45
2	A	1	J2I	C21-C12-C5	3.97	134.56	128.62
2	A	1	J2I	C10-C7-C9	3.87	119.76	114.54
2	A	1	J2I	C2-C11-N28	3.65	108.65	106.22
2	B	2	J2I	C12-N27-N24	3.63	114.11	105.86
2	A	1	J2I	F35-C10-C7	3.46	121.95	117.64
2	B	2	J2I	C21-C12-C5	3.45	133.79	128.62
2	A	1	J2I	C12-N27-N24	3.29	113.35	105.86
2	A	1	J2I	C15-N25-C14	3.17	122.65	116.21
2	B	2	J2I	C2-C11-N28	3.09	108.27	106.22
2	A	1	J2I	C23-C7-C9	-3.08	116.97	123.15
2	A	1	J2I	C18-C20-N30	2.98	116.35	110.42
2	B	2	J2I	F35-C10-C7	2.93	121.29	117.64
2	B	2	J2I	C10-C7-C9	2.76	118.26	114.54
2	B	2	J2I	C15-N25-C14	2.76	121.81	116.21
2	A	1	J2I	C20-N30-C19	2.52	117.81	112.68
2	B	2	J2I	C20-N30-C19	2.44	117.65	112.68
2	A	1	J2I	O32-C16-C23	-2.40	117.79	121.61
2	A	1	J2I	C17-C19-N30	2.14	114.66	110.42
2	A	1	J2I	C1-C6-N28	2.05	108.24	106.73

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1	J2I	C4-C8-O33-C22
2	A	1	J2I	C3-C8-O33-C22
2	B	2	J2I	C3-C8-O33-C22
2	B	2	J2I	C4-C8-O33-C22
2	B	2	J2I	C16-C23-C7-C9

There are no ring outliers.

2 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2	J2I	1	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	J2I	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	291/301 (96%)	0.14	18 (6%) 26 23	30, 49, 99, 128	0
1	B	294/301 (97%)	0.13	14 (4%) 35 31	33, 50, 89, 114	0
All	All	585/602 (97%)	0.13	32 (5%) 30 27	30, 50, 95, 128	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	921	GLY	5.0
1	A	860	PHE	4.6
1	A	841	PRO	4.5
1	A	1132	GLY	4.2
1	A	886	HIS	3.8
1	B	918	TYR	3.8
1	B	923	ARG	3.7
1	A	920	ALA	3.6
1	B	1011	LYS	3.4
1	A	1011	LYS	3.4
1	B	842	THR	3.1
1	A	842	THR	3.1
1	A	918	TYR	3.0
1	A	922	ARG	2.9
1	B	1133	HIS	2.9
1	A	1071	GLY	2.8
1	A	887	SER	2.7
1	B	920	ALA	2.7
1	B	841	PRO	2.7
1	A	924	ASN	2.7
1	B	1071	GLY	2.6
1	B	1013	PRO	2.5
1	B	1012	GLU	2.4
1	B	838	MET	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1070	GLN	2.3
1	A	1013	PRO	2.3
1	A	919	SER	2.2
1	A	840	ASP	2.2
1	B	922	ARG	2.1
1	B	924	ASN	2.1
1	A	949	ASP	2.0
1	A	1012	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PTR	B	1008	16/17	0.90	0.11	63,78,89,90	0
1	PTR	B	1007	16/17	0.92	0.11	66,73,84,84	0
1	PTR	A	1007	16/17	0.94	0.07	43,48,65,65	0
1	PTR	A	1008	16/17	0.94	0.09	43,55,73,73	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	J2I	A	1	35/35	0.96	0.07	31,39,54,56	0
2	J2I	B	2	35/35	0.96	0.07	30,40,53,58	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around J2I A 1:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.