



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 05:58 PM UTC

PDB ID : 4Q3O / pdb_00004q3o
Title : Crystal structure of MGS-MT1, an alpha/beta hydrolase enzyme from a Lake Matapan deep-sea metagenome library
Authors : Stogios, P.J.; Xu, X.; Cui, H.; Alcaide, M.; Ferrer, M.; Savchenko, A.
Deposited on : 2014-04-11
Resolution : 1.74 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1187 (1.74-1.74)
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)
RSRZ outliers	180081	1188 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	348	2% 82% 9% 9%
1	B	348	2% 83% 8% 9%
1	C	348	0% 82% 8% 9%
1	D	348	0% 82% 8% 9%
1	E	348	0% 84% 7% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	348	 84% 7% 9%
1	G	348	 84% 7% 9%
1	H	348	 81% 9% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CL	A	401	-	-	X	-
2	CL	C	401	-	-	X	-
2	CL	E	401	-	-	X	-
3	MES	A	404	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 23319 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MGS-MT1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	317	Total	C	N	O	S	0	5	0
			2501	1608	424	463	6			
1	B	317	Total	C	N	O	S	0	4	0
			2489	1602	418	463	6			
1	C	316	Total	C	N	O	S	0	4	0
			2486	1604	417	459	6			
1	D	316	Total	C	N	O	S	0	6	0
			2498	1610	423	459	6			
1	E	316	Total	C	N	O	S	0	5	0
			2490	1603	420	461	6			
1	F	316	Total	C	N	O	S	0	3	0
			2477	1595	417	459	6			
1	G	316	Total	C	N	O	S	0	4	0
			2487	1603	420	458	6			
1	H	316	Total	C	N	O	S	0	1	0
			2471	1590	417	457	7			

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cl	0	0
			2	2		
2	B	1	Total	Cl	0	0
			1	1		
2	C	2	Total	Cl	0	0
			2	2		
2	D	1	Total	Cl	0	0
			1	1		
2	E	2	Total	Cl	0	0
			2	2		
2	F	1	Total	Cl	0	0
			1	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total	Cl	0	0
			1	1		

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	G	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	G	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	H	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	H	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	E	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	E	1	Total	C	O	0	0
			6	3	3		
4	F	1	Total	C	O	0	0
			6	3	3		
4	F	1	Total	C	O	0	0
			6	3	3		
4	G	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	391	Total	O	0	16
			407	407		
5	B	435	Total	O	0	15
			450	450		
5	C	408	Total	O	0	18
			428	428		
5	D	391	Total	O	0	10
			401	401		
5	E	390	Total	O	0	15
			405	405		
5	F	384	Total	O	0	7
			391	391		
5	G	354	Total	O	0	12
			366	366		
5	H	332	Total	O	0	8
			340	340		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MGS-MT1




- Molecule 1: MGS-MT1



- Molecule 1: MGS-MT1





- Molecule 1: MGS-MT1










- Molecule 1: MGS-MT1






- Molecule 1: MGS-MT1



- Molecule 1: MGS-MT1



- Molecule 1: MGS-MT1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	189.45Å 131.40Å 112.44Å 90.00° 103.55° 90.00°	Depositor
Resolution (Å)	19.72 – 1.74 19.72 – 1.74	Depositor EDS
% Data completeness (in resolution range)	99.5 (19.72-1.74) 93.1 (19.72-1.74)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 1.74Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9pre_1669)	Depositor
R, R_{free}	0.145 , 0.185 0.146 , 0.186	Depositor DCC
R_{free} test set	13735 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	15.6	Xtrriage
Anisotropy	0.198	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 68.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	23319	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 48.90 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.0541e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MES, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	0/2579	0.92	1/3512 (0.0%)
1	B	0.92	1/2564 (0.0%)	0.95	2/3493 (0.1%)
1	C	0.86	3/2562 (0.1%)	0.94	2/3491 (0.1%)
1	D	0.86	0/2579	0.94	0/3512
1	E	0.85	0/2568	0.90	0/3498
1	F	0.86	0/2549	0.92	3/3473 (0.1%)
1	G	0.80	1/2562 (0.0%)	0.89	0/3490
1	H	0.78	0/2537	0.91	1/3457 (0.0%)
All	All	0.85	5/20500 (0.0%)	0.92	9/27926 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	222	THR	CA-CB	6.72	1.58	1.52
1	C	49	TYR	CA-C	-6.51	1.44	1.52
1	G	222	THR	CA-CB	6.43	1.58	1.52
1	C	222	THR	CA-CB	5.58	1.57	1.52
1	C	49	TYR	CA-CB	-5.14	1.46	1.53

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	60	PHE	N-CA-C	7.84	118.96	108.38
1	C	49	TYR	CA-C-N	-6.33	113.16	119.99
1	C	49	TYR	C-N-CA	-6.33	113.16	119.99
1	B	137	SER	CA-C-N	-5.65	115.25	122.26
1	B	137	SER	C-N-CA	-5.65	115.25	122.26

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	60	PHE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2501	0	2512	25	0
1	B	2489	0	2495	18	0
1	C	2486	0	2495	21	0
1	D	2498	0	2521	24	0
1	E	2490	0	2501	16	0
1	F	2477	0	2484	16	0
1	G	2487	0	2503	18	0
1	H	2471	0	2472	22	0
2	A	2	0	0	2	0
2	B	1	0	0	1	0
2	C	2	0	0	2	0
2	D	1	0	0	0	0
2	E	2	0	0	3	0
2	F	1	0	0	0	0
2	G	1	0	0	1	0
3	A	24	0	24	9	0
3	B	12	0	12	1	0
3	C	24	0	24	1	0
3	D	12	0	12	0	0
3	G	24	0	24	6	0
3	H	24	0	24	2	0
4	A	18	0	24	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	6	0	8	0	0
4	C	18	0	24	1	0
4	D	30	0	40	0	0
4	E	12	0	16	0	0
4	F	12	0	16	0	0
4	G	6	0	8	0	0
5	A	407	0	0	3	0
5	B	450	0	0	3	0
5	C	428	0	0	7	0
5	D	401	0	0	9	0
5	E	405	0	0	6	0
5	F	391	0	0	8	0
5	G	366	0	0	6	0
5	H	340	0	0	5	0
All	All	23319	0	20239	166	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 166 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:404:MES:H71	1:B:37:SER:HB2	1.42	0.98
2:A:401:CL:CL	5:A:859:HOH:O	2.25	0.90
2:E:401:CL:CL	5:E:827:HOH:O	2.30	0.86
2:C:401:CL:CL	5:C:739:HOH:O	2.32	0.83
1:C:78:MET:SD	5:C:719:HOH:O	2.36	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	320/348 (92%)	309 (97%)	11 (3%)	0	100	100
1	B	319/348 (92%)	309 (97%)	10 (3%)	0	100	100
1	C	318/348 (91%)	309 (97%)	9 (3%)	0	100	100
1	D	320/348 (92%)	308 (96%)	12 (4%)	0	100	100
1	E	319/348 (92%)	307 (96%)	12 (4%)	0	100	100
1	F	317/348 (91%)	302 (95%)	14 (4%)	1 (0%)	36	23
1	G	318/348 (91%)	304 (96%)	14 (4%)	0	100	100
1	H	315/348 (90%)	301 (96%)	14 (4%)	0	100	100
All	All	2546/2784 (92%)	2449 (96%)	96 (4%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	184	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/291 (93%)	266 (98%)	5 (2%)	51	30
1	B	270/291 (93%)	269 (100%)	1 (0%)	84	79
1	C	269/291 (92%)	267 (99%)	2 (1%)	76	67
1	D	271/291 (93%)	268 (99%)	3 (1%)	65	50
1	E	270/291 (93%)	267 (99%)	3 (1%)	65	50
1	F	268/291 (92%)	265 (99%)	3 (1%)	65	50
1	G	269/291 (92%)	268 (100%)	1 (0%)	84	79
1	H	266/291 (91%)	261 (98%)	5 (2%)	50	28
All	All	2154/2328 (92%)	2131 (99%)	23 (1%)	68	50

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	55	GLU
1	G	318	LEU
1	F	181	GLN
1	H	48	GLN
1	C	184	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	186	ASN
1	H	174	HIS
1	F	187	HIS
1	H	67	GLN
1	E	333	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 37 ligands modelled in this entry, 10 are monoatomic - leaving 27 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	407	-	5,5,5	0.93	0	5,5,5	0.77	0
4	GOL	C	405	-	5,5,5	0.28	0	5,5,5	0.53	0
3	MES	G	403	-	12,12,12	1.93	1 (8%)	15,16,16	2.14	4 (26%)
4	GOL	D	405	-	5,5,5	0.37	0	5,5,5	0.50	0
3	MES	A	404	-	12,12,12	2.42	1 (8%)	15,16,16	2.78	6 (40%)
3	MES	C	402	-	12,12,12	2.24	1 (8%)	15,16,16	2.01	5 (33%)
4	GOL	F	403	-	5,5,5	0.42	0	5,5,5	0.88	0
4	GOL	C	406	-	5,5,5	0.47	0	5,5,5	0.51	0
4	GOL	G	404	-	5,5,5	0.46	0	5,5,5	1.01	0
3	MES	C	403	-	12,12,12	1.87	1 (8%)	15,16,16	1.84	3 (20%)
3	MES	H	401	-	12,12,12	2.31	2 (16%)	15,16,16	2.23	5 (33%)
3	MES	H	402	-	12,12,12	2.08	1 (8%)	15,16,16	2.66	7 (46%)
3	MES	A	403	-	12,12,12	2.17	2 (16%)	15,16,16	2.23	5 (33%)
4	GOL	E	404	-	5,5,5	0.50	0	5,5,5	0.56	0
4	GOL	E	403	-	5,5,5	0.35	0	5,5,5	0.42	0
4	GOL	C	404	-	5,5,5	0.45	0	5,5,5	0.32	0
4	GOL	D	406	-	5,5,5	0.31	0	5,5,5	0.22	0
4	GOL	D	402	-	5,5,5	0.49	0	5,5,5	0.91	0
4	GOL	F	402	-	5,5,5	0.20	0	5,5,5	0.86	0
3	MES	D	401	-	12,12,12	1.97	1 (8%)	15,16,16	2.55	4 (26%)
4	GOL	A	405	-	5,5,5	0.51	0	5,5,5	0.51	0
3	MES	B	402	-	12,12,12	1.94	1 (8%)	15,16,16	2.01	4 (26%)
4	GOL	D	403	-	5,5,5	0.31	0	5,5,5	1.72	1 (20%)
4	GOL	D	404	-	5,5,5	0.39	0	5,5,5	0.60	0
3	MES	G	402	-	12,12,12	2.19	2 (16%)	15,16,16	2.34	5 (33%)
4	GOL	B	403	-	5,5,5	0.35	0	5,5,5	0.94	0
4	GOL	A	406	-	5,5,5	0.39	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	407	-	-	0/4/4/4	-
4	GOL	C	405	-	-	0/4/4/4	-
3	MES	G	403	-	-	2/6/14/14	0/1/1/1
4	GOL	D	405	-	-	2/4/4/4	-
3	MES	A	404	-	-	1/6/14/14	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MES	C	402	-	-	4/6/14/14	0/1/1/1
4	GOL	F	403	-	-	0/4/4/4	-
4	GOL	C	406	-	-	1/4/4/4	-
4	GOL	G	404	-	-	1/4/4/4	-
3	MES	C	403	-	-	4/6/14/14	0/1/1/1
3	MES	H	401	-	-	1/6/14/14	0/1/1/1
3	MES	H	402	-	-	5/6/14/14	0/1/1/1
3	MES	A	403	-	-	4/6/14/14	0/1/1/1
4	GOL	E	404	-	-	2/4/4/4	-
4	GOL	E	403	-	-	3/4/4/4	-
4	GOL	C	404	-	-	2/4/4/4	-
4	GOL	D	406	-	-	0/4/4/4	-
4	GOL	D	402	-	-	3/4/4/4	-
4	GOL	F	402	-	-	2/4/4/4	-
3	MES	D	401	-	-	5/6/14/14	0/1/1/1
4	GOL	A	405	-	-	2/4/4/4	-
3	MES	B	402	-	-	4/6/14/14	0/1/1/1
4	GOL	D	403	-	-	4/4/4/4	-
4	GOL	D	404	-	-	2/4/4/4	-
3	MES	G	402	-	-	2/6/14/14	0/1/1/1
4	GOL	B	403	-	-	0/4/4/4	-
4	GOL	A	406	-	-	4/4/4/4	-

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	404	MES	C8-S	-8.03	1.66	1.77
3	C	402	MES	C8-S	-7.51	1.67	1.77
3	H	401	MES	C8-S	-7.29	1.67	1.77
3	H	402	MES	C8-S	-6.95	1.67	1.77
3	G	402	MES	C8-S	-6.89	1.67	1.77

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	401	MES	C5-N4-C3	6.50	122.85	108.84
3	A	404	MES	C5-N4-C3	6.24	122.29	108.84
3	H	402	MES	O3S-S-C8	6.17	118.07	106.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	404	MES	O3S-S-C8	6.09	117.92	106.00
3	A	403	MES	O2S-S-C8	5.71	115.36	106.73

There are no chirality outliers.

5 of 60 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	404	MES	N4-C7-C8-S
3	B	402	MES	C8-C7-N4-C5
3	C	402	MES	C8-C7-N4-C3
3	C	402	MES	C7-C8-S-O1S
3	C	402	MES	C7-C8-S-O3S

There are no ring outliers.

11 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	403	MES	2	0
3	A	404	MES	7	0
3	C	403	MES	1	0
3	H	401	MES	1	0
3	H	402	MES	1	0
3	A	403	MES	2	0
4	C	404	GOL	1	0
4	A	405	GOL	1	0
3	B	402	MES	1	0
3	G	402	MES	4	0
4	A	406	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	317/348 (91%)	-0.47	6 (1%) 66 73	8, 16, 44, 103	5 (1%)
1	B	317/348 (91%)	-0.62	6 (1%) 66 73	6, 14, 32, 94	4 (1%)
1	C	316/348 (90%)	-0.60	4 (1%) 75 82	7, 15, 33, 86	4 (1%)
1	D	316/348 (90%)	-0.60	3 (0%) 81 87	5, 16, 31, 71	6 (1%)
1	E	316/348 (90%)	-0.51	3 (0%) 81 87	7, 16, 35, 83	5 (1%)
1	F	316/348 (90%)	-0.48	2 (0%) 85 90	8, 16, 49, 88	3 (0%)
1	G	316/348 (90%)	-0.40	5 (1%) 70 77	8, 19, 48, 84	4 (1%)
1	H	316/348 (90%)	-0.16	3 (0%) 81 87	11, 22, 49, 105	1 (0%)
All	All	2530/2784 (90%)	-0.48	32 (1%) 75 82	5, 17, 42, 105	32 (1%)

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	51	THR	5.9
1	E	49	TYR	5.8
1	H	60	PHE	5.5
1	D	50	PRO	4.9
1	F	60	PHE	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	D	405	6/6	0.57	0.19	79,80,80,81	0
4	GOL	A	406	6/6	0.74	0.15	41,55,58,62	0
4	GOL	E	403	6/6	0.74	0.14	46,57,58,62	6
4	GOL	D	403	6/6	0.75	0.14	42,46,58,62	0
2	CL	C	401	1/1	0.77	0.16	54,54,54,54	0
4	GOL	B	403	6/6	0.79	0.14	29,40,42,48	6
4	GOL	C	404	6/6	0.81	0.16	38,52,58,61	0
4	GOL	D	404	6/6	0.81	0.13	35,52,59,60	0
4	GOL	F	402	6/6	0.81	0.12	34,53,60,62	6
4	GOL	C	405	6/6	0.82	0.11	27,36,39,43	6
3	MES	B	402	12/12	0.82	0.16	18,27,32,34	12
4	GOL	E	404	6/6	0.83	0.13	36,44,52,56	0
2	CL	B	401	1/1	0.83	0.13	55,55,55,55	0
4	GOL	A	405	6/6	0.84	0.13	36,42,47,47	6
3	MES	D	401	12/12	0.85	0.13	18,25,37,38	12
2	CL	E	401	1/1	0.85	0.14	55,55,55,55	0
3	MES	C	403	12/12	0.85	0.13	19,26,34,36	12
3	MES	A	404	12/12	0.86	0.17	12,34,44,45	12
3	MES	G	402	12/12	0.87	0.20	14,52,59,60	12
3	MES	H	401	12/12	0.88	0.18	10,40,52,54	12
4	GOL	D	402	6/6	0.88	0.12	34,51,54,54	0
2	CL	A	401	1/1	0.89	0.13	45,45,45,45	1
2	CL	G	401	1/1	0.89	0.12	53,53,53,53	0
3	MES	G	403	12/12	0.90	0.16	26,52,60,62	12
2	CL	F	401	1/1	0.91	0.11	49,49,49,49	0
4	GOL	D	406	6/6	0.94	0.08	20,23,33,41	0
3	MES	C	402	12/12	0.94	0.12	19,35,49,49	12
4	GOL	C	406	6/6	0.94	0.07	27,29,31,40	6
3	MES	H	402	12/12	0.94	0.12	19,37,51,52	12
4	GOL	G	404	6/6	0.94	0.08	15,22,36,43	0
4	GOL	A	407	6/6	0.95	0.08	13,17,26,41	6
3	MES	A	403	12/12	0.95	0.14	13,37,55,58	12
4	GOL	F	403	6/6	0.96	0.07	14,16,26,33	0
2	CL	E	402	1/1	0.99	0.03	14,14,14,14	0
2	CL	D	407	1/1	0.99	0.04	13,13,13,13	0
2	CL	A	402	1/1	1.00	0.04	13,13,13,13	0
2	CL	C	407	1/1	1.00	0.02	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.