



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 06:06 PM UTC

PDB ID : 2Q44 / pdb_00002q44
Title : Ensemble refinement of the protein crystal structure of gene product from Arabidopsis thaliana At1g77540
Authors : Levin, E.J.; Kondrashov, D.A.; Wesenberg, G.E.; Phillips Jr., G.N.; Center for Eukaryotic Structural Genomics (CESG)
Deposited on : 2007-05-31
Resolution : 1.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1380 (1.16-1.12)
Clashscore	190562	1393 (1.16-1.12)
Ramachandran outliers	187476	1369 (1.16-1.12)
Sidechain outliers	187428	1369 (1.16-1.12)
RSRZ outliers	180081	1379 (1.16-1.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1-A	103	85% 32% 50% 8% 8%
1	2-A	103	32% 50% 10% 8%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1797 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein At1g77540.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	1-A	95	Total	C	N	O	S	0	0	0
			768	494	134	136	4			
1	2-A	95	Total	C	N	O	S	0	0	0
			768	494	134	136	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP Q9CAQ2

- Molecule 2 is BROMIDE ION (CCD ID: HOH) (formula: H₂O).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	1-A	11	Total	Br	0	0
			11	11		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	1-A	125	Total	O	0	0
			125	125		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Uncharacterized protein At1g77540



• Molecule 1: Uncharacterized protein At1g77540



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	27.35Å 60.60Å 29.42Å 90.00° 91.50° 90.00°	Depositor
Resolution (Å)	20.29 – 1.15 20.29 – 1.15	Depositor EDS
% Data completeness (in resolution range)	95.8 (20.29-1.15) 95.9 (20.29-1.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 1.15Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.165 , 0.190 0.172 , 0.201	Depositor DCC
R_{free} test set	1658 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	11.5	Xtriage
Anisotropy	0.153	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.25 , 52.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.047 for h,-k,-l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	1797	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	1-A	2.95	75/793 (9.5%)	2.04	33/1074 (3.1%)
1	2-A	2.95	75/793 (9.5%)	2.04	33/1074 (3.1%)
All	All	2.95	150/1586 (9.5%)	2.04	66/2148 (3.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	1-A	0	9
1	2-A	0	10
All	All	0	19

All (150) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	1-A	80	SER	CA-CB	-12.79	1.33	1.53
1	2-A	80	SER	CA-CB	-12.79	1.33	1.53
1	1-A	83	PHE	CG-CD2	11.91	1.63	1.38
1	2-A	83	PHE	CG-CD2	11.91	1.63	1.38
1	1-A	98	VAL	CA-CB	11.67	1.68	1.54
1	2-A	98	VAL	CA-CB	11.67	1.68	1.54
1	1-A	76	CYS	N-CA	-10.56	1.32	1.46
1	2-A	76	CYS	N-CA	-10.56	1.32	1.46
1	1-A	46	SER	CA-CB	-9.99	1.37	1.53
1	2-A	46	SER	CA-CB	-9.99	1.37	1.53
1	1-A	34	GLY	C-O	9.30	1.37	1.24
1	2-A	34	GLY	C-O	9.30	1.37	1.24
1	1-A	83	PHE	CG-CD1	-9.14	1.19	1.38
1	2-A	83	PHE	CG-CD1	-9.14	1.19	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	1-A	30	MET	SD-CE	-9.12	1.56	1.79
1	2-A	30	MET	SD-CE	-9.12	1.56	1.79
1	1-A	83	PHE	CB-CG	-9.09	1.29	1.50
1	2-A	83	PHE	CB-CG	-9.09	1.29	1.50
1	1-A	35	LYS	C-O	8.93	1.35	1.24
1	2-A	35	LYS	C-O	8.93	1.35	1.24
1	1-A	66	SER	CA-C	-8.90	1.41	1.52
1	2-A	66	SER	CA-C	-8.90	1.41	1.52
1	1-A	11	ASN	CA-C	8.76	1.64	1.53
1	2-A	11	ASN	CA-C	8.76	1.64	1.53
1	1-A	74	PRO	CA-C	8.39	1.60	1.53
1	2-A	74	PRO	CA-C	8.39	1.60	1.53
1	1-A	91	LYS	CE-NZ	8.15	1.73	1.49
1	2-A	91	LYS	CE-NZ	8.15	1.73	1.49
1	1-A	35	LYS	CG-CD	8.07	1.76	1.52
1	2-A	35	LYS	CG-CD	8.07	1.76	1.52
1	1-A	15	ARG	C-N	-7.97	1.22	1.33
1	2-A	15	ARG	C-N	-7.97	1.22	1.33
1	1-A	44	VAL	CA-CB	7.89	1.60	1.54
1	2-A	44	VAL	CA-CB	7.89	1.60	1.54
1	1-A	99	PHE	CA-C	7.84	1.69	1.52
1	2-A	99	PHE	CA-C	7.84	1.69	1.52
1	1-A	30	MET	CG-SD	-7.84	1.61	1.80
1	2-A	30	MET	CG-SD	-7.84	1.61	1.80
1	1-A	99	PHE	C-O	7.74	1.39	1.23
1	2-A	99	PHE	C-O	7.74	1.39	1.23
1	1-A	71	SER	CA-CB	7.69	1.64	1.53
1	2-A	71	SER	CA-CB	7.69	1.64	1.53
1	1-A	15	ARG	CZ-NH1	7.55	1.43	1.32
1	2-A	15	ARG	CZ-NH1	7.55	1.43	1.32
1	1-A	48	LYS	CB-CG	7.41	1.74	1.52
1	2-A	48	LYS	CB-CG	7.41	1.74	1.52
1	1-A	72	ILE	C-O	7.38	1.31	1.24
1	2-A	72	ILE	C-O	7.38	1.31	1.24
1	1-A	76	CYS	CA-CB	7.17	1.65	1.53
1	2-A	76	CYS	CA-CB	7.17	1.65	1.53
1	1-A	31	ARG	C-N	7.17	1.42	1.33
1	2-A	31	ARG	C-N	7.17	1.42	1.33
1	1-A	54	ALA	C-O	7.09	1.32	1.24
1	2-A	54	ALA	C-O	7.09	1.32	1.24
1	1-A	38	ASP	CG-OD2	7.08	1.38	1.25
1	2-A	38	ASP	CG-OD2	7.08	1.38	1.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	1-A	73	ILE	CA-C	7.00	1.58	1.53
1	2-A	73	ILE	CA-C	7.00	1.58	1.53
1	1-A	31	ARG	CZ-NH2	6.79	1.42	1.33
1	2-A	31	ARG	CZ-NH2	6.79	1.42	1.33
1	1-A	12	GLU	N-CA	-6.75	1.37	1.46
1	2-A	12	GLU	N-CA	-6.75	1.37	1.46
1	1-A	56	HIS	C-O	-6.70	1.15	1.24
1	2-A	56	HIS	C-O	-6.70	1.15	1.24
1	1-A	96	SER	CB-OG	-6.64	1.28	1.42
1	2-A	96	SER	CB-OG	-6.64	1.28	1.42
1	1-A	22	HIS	CG-ND1	-6.50	1.31	1.38
1	2-A	22	HIS	CG-ND1	-6.50	1.31	1.38
1	1-A	21	ASP	CG-OD2	6.47	1.37	1.25
1	2-A	21	ASP	CG-OD2	6.47	1.37	1.25
1	1-A	79	VAL	N-CA	-6.39	1.38	1.46
1	2-A	79	VAL	N-CA	-6.39	1.38	1.46
1	1-A	95	HIS	C-O	6.35	1.31	1.23
1	2-A	95	HIS	C-O	6.35	1.31	1.23
1	1-A	21	ASP	CA-C	6.25	1.64	1.52
1	2-A	21	ASP	CA-C	6.25	1.64	1.52
1	1-A	37	MET	SD-CE	-6.22	1.64	1.79
1	2-A	37	MET	SD-CE	-6.22	1.64	1.79
1	1-A	7	LYS	CA-C	6.18	1.60	1.52
1	1-A	39	LEU	C-O	-6.18	1.16	1.24
1	2-A	7	LYS	CA-C	6.18	1.60	1.52
1	2-A	39	LEU	C-O	-6.18	1.16	1.24
1	1-A	20	GLU	C-O	6.17	1.31	1.24
1	2-A	20	GLU	C-O	6.17	1.31	1.24
1	1-A	8	ILE	C-N	-6.14	1.25	1.33
1	2-A	8	ILE	C-N	-6.14	1.25	1.33
1	1-A	22	HIS	ND1-CE1	6.13	1.38	1.32
1	2-A	22	HIS	ND1-CE1	6.13	1.38	1.32
1	1-A	67	SER	CA-CB	-6.11	1.43	1.53
1	2-A	67	SER	CA-CB	-6.11	1.43	1.53
1	1-A	96	SER	CA-CB	6.01	1.63	1.53
1	2-A	96	SER	CA-CB	6.01	1.63	1.53
1	1-A	13	GLY	C-N	5.92	1.41	1.33
1	2-A	13	GLY	C-N	5.92	1.41	1.33
1	1-A	44	VAL	CA-C	-5.87	1.48	1.52
1	2-A	44	VAL	CA-C	-5.87	1.48	1.52
1	1-A	55	SER	C-N	5.86	1.41	1.33
1	2-A	55	SER	C-N	5.86	1.41	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	1-A	5	PRO	CG-CD	5.78	1.70	1.50
1	2-A	5	PRO	CG-CD	5.78	1.70	1.50
1	1-A	82	THR	C-N	5.77	1.41	1.33
1	2-A	82	THR	C-N	5.77	1.41	1.33
1	1-A	96	SER	C-N	-5.75	1.25	1.33
1	2-A	96	SER	C-N	-5.75	1.25	1.33
1	1-A	97	GLU	C-N	5.71	1.41	1.33
1	2-A	97	GLU	C-N	5.71	1.41	1.33
1	1-A	28	TYR	C-N	5.71	1.40	1.33
1	2-A	28	TYR	C-N	5.71	1.40	1.33
1	1-A	47	PHE	CD1-CE1	5.69	1.55	1.38
1	2-A	47	PHE	CD1-CE1	5.69	1.55	1.38
1	1-A	36	VAL	C-O	-5.62	1.18	1.24
1	2-A	36	VAL	C-O	-5.62	1.18	1.24
1	1-A	58	CYS	C-O	-5.51	1.17	1.24
1	2-A	58	CYS	C-O	-5.51	1.17	1.24
1	1-A	41	HIS	C-N	5.49	1.41	1.33
1	2-A	41	HIS	C-N	5.49	1.41	1.33
1	1-A	43	TYR	CZ-OH	-5.48	1.26	1.38
1	2-A	43	TYR	CZ-OH	-5.48	1.26	1.38
1	1-A	90	TRP	CA-CB	5.42	1.63	1.53
1	2-A	90	TRP	CA-CB	5.42	1.63	1.53
1	1-A	9	VAL	CB-CG1	-5.40	1.34	1.52
1	2-A	9	VAL	CB-CG1	-5.40	1.34	1.52
1	1-A	19	THR	C-O	5.39	1.31	1.23
1	2-A	19	THR	C-O	5.39	1.31	1.23
1	1-A	71	SER	C-O	5.39	1.30	1.23
1	2-A	71	SER	C-O	5.39	1.30	1.23
1	1-A	8	ILE	CB-CG1	5.36	1.64	1.53
1	2-A	8	ILE	CB-CG1	5.36	1.64	1.53
1	1-A	8	ILE	C-O	5.29	1.30	1.23
1	2-A	8	ILE	C-O	5.29	1.30	1.23
1	1-A	32	ASN	CA-C	5.25	1.59	1.53
1	2-A	32	ASN	CA-C	5.25	1.59	1.53
1	1-A	47	PHE	CA-CB	5.24	1.62	1.53
1	2-A	47	PHE	CA-CB	5.24	1.62	1.53
1	1-A	32	ASN	CA-CB	-5.22	1.46	1.53
1	2-A	32	ASN	CA-CB	-5.22	1.46	1.53
1	1-A	35	LYS	CA-C	5.21	1.59	1.52
1	2-A	35	LYS	CA-C	5.21	1.59	1.52
1	1-A	12	GLU	CA-CB	5.12	1.61	1.53
1	2-A	12	GLU	CA-CB	5.12	1.61	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	1-A	72	ILE	CB-CG1	5.09	1.63	1.53
1	2-A	72	ILE	CB-CG1	5.09	1.63	1.53
1	1-A	50	GLY	CA-C	-5.07	1.44	1.51
1	2-A	50	GLY	CA-C	-5.07	1.44	1.51
1	1-A	88	PRO	CG-CD	5.07	1.68	1.50
1	2-A	88	PRO	CG-CD	5.07	1.68	1.50
1	1-A	66	SER	C-O	5.05	1.29	1.24
1	2-A	66	SER	C-O	5.05	1.29	1.24
1	1-A	92	PRO	C-O	5.03	1.30	1.24
1	2-A	92	PRO	C-O	5.03	1.30	1.24

All (66) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	1-A	39	LEU	CA-C-O	8.30	132.38	120.51
1	2-A	39	LEU	CA-C-O	8.30	132.38	120.51
1	1-A	13	GLY	CA-C-N	-7.70	110.51	122.37
1	1-A	13	GLY	C-N-CA	-7.70	110.51	122.37
1	2-A	13	GLY	CA-C-N	-7.70	110.51	122.37
1	2-A	13	GLY	C-N-CA	-7.70	110.51	122.37
1	1-A	31	ARG	CD-NE-CZ	-7.33	114.14	124.40
1	2-A	31	ARG	CD-NE-CZ	-7.33	114.14	124.40
1	1-A	50	GLY	O-C-N	-7.08	114.64	122.50
1	2-A	50	GLY	O-C-N	-7.08	114.64	122.50
1	1-A	8	ILE	O-C-N	6.53	130.74	123.10
1	2-A	8	ILE	O-C-N	6.53	130.74	123.10
1	1-A	78	TYR	CA-C-O	-6.51	113.99	120.82
1	2-A	78	TYR	CA-C-O	-6.51	113.99	120.82
1	1-A	8	ILE	CA-C-N	6.46	131.86	122.69
1	1-A	8	ILE	C-N-CA	6.46	131.86	122.69
1	2-A	8	ILE	CA-C-N	6.46	131.86	122.69
1	2-A	8	ILE	C-N-CA	6.46	131.86	122.69
1	1-A	96	SER	N-CA-C	-6.26	104.90	112.54
1	2-A	96	SER	N-CA-C	-6.26	104.90	112.54
1	1-A	25	PHE	CA-C-N	6.12	131.68	122.98
1	1-A	25	PHE	C-N-CA	6.12	131.68	122.98
1	2-A	25	PHE	CA-C-N	6.12	131.68	122.98
1	2-A	25	PHE	C-N-CA	6.12	131.68	122.98
1	1-A	98	VAL	O-C-N	5.96	128.10	121.90
1	2-A	98	VAL	O-C-N	5.96	128.10	121.90
1	1-A	7	LYS	O-C-N	5.90	129.94	123.04
1	2-A	7	LYS	O-C-N	5.90	129.94	123.04

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	1-A	90	TRP	CZ3-CH2-CZ2	5.82	129.06	121.50
1	2-A	90	TRP	CZ3-CH2-CZ2	5.82	129.06	121.50
1	1-A	31	ARG	CA-C-O	5.79	128.57	121.56
1	2-A	31	ARG	CA-C-O	5.79	128.57	121.56
1	1-A	41	HIS	CE1-NE2-CD2	-5.74	103.26	109.00
1	2-A	41	HIS	CE1-NE2-CD2	-5.74	103.26	109.00
1	1-A	23	GLU	CG-CD-OE2	-5.65	105.40	118.40
1	2-A	23	GLU	CG-CD-OE2	-5.65	105.40	118.40
1	1-A	5	PRO	N-CA-CB	5.62	109.18	103.00
1	2-A	5	PRO	N-CA-CB	5.62	109.18	103.00
1	1-A	78	TYR	N-CA-C	-5.60	105.08	111.07
1	2-A	78	TYR	N-CA-C	-5.60	105.08	111.07
1	1-A	66	SER	O-C-N	-5.48	116.31	122.12
1	2-A	66	SER	O-C-N	-5.48	116.31	122.12
1	1-A	81	ASP	CB-CG-OD2	5.46	130.95	118.40
1	2-A	81	ASP	CB-CG-OD2	5.46	130.95	118.40
1	1-A	26	ILE	N-CA-CB	5.40	119.16	111.82
1	2-A	26	ILE	N-CA-CB	5.40	119.16	111.82
1	1-A	27	GLU	O-C-N	5.37	130.03	123.21
1	2-A	27	GLU	O-C-N	5.37	130.03	123.21
1	1-A	63	GLU	OE1-CD-OE2	5.34	135.71	122.90
1	2-A	63	GLU	OE1-CD-OE2	5.34	135.71	122.90
1	1-A	38	ASP	OD1-CG-OD2	5.18	135.33	122.90
1	2-A	38	ASP	OD1-CG-OD2	5.18	135.33	122.90
1	1-A	17	PHE	CA-C-O	5.15	126.17	120.71
1	2-A	17	PHE	CA-C-O	5.15	126.17	120.71
1	1-A	90	TRP	CA-C-O	5.13	125.68	119.11
1	2-A	90	TRP	CA-C-O	5.13	125.68	119.11
1	1-A	22	HIS	O-C-N	5.13	128.41	122.00
1	2-A	22	HIS	O-C-N	5.13	128.41	122.00
1	1-A	95	HIS	ND1-CG-CD2	5.07	111.17	106.10
1	2-A	95	HIS	ND1-CG-CD2	5.07	111.17	106.10
1	1-A	90	TRP	CB-CG-CD2	-5.05	119.73	126.80
1	2-A	90	TRP	CB-CG-CD2	-5.05	119.73	126.80
1	1-A	25	PHE	CE1-CZ-CE2	5.03	129.06	120.00
1	2-A	25	PHE	CE1-CZ-CE2	5.03	129.06	120.00
1	1-A	39	LEU	O-C-N	-5.02	115.92	122.59
1	2-A	39	LEU	O-C-N	-5.02	115.92	122.59

There are no chirality outliers.

All (19) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	1-A	13	GLY	Mainchain
1	1-A	14	LYS	Mainchain
1	1-A	15	ARG	Mainchain
1	1-A	21	ASP	Sidechain
1	1-A	28	TYR	Mainchain
1	1-A	30	MET	Mainchain
1	1-A	31	ARG	Sidechain
1	1-A	79	VAL	Mainchain
1	1-A	97	GLU	Mainchain
1	2-A	15	ARG	Mainchain
1	2-A	28	TYR	Mainchain
1	2-A	30	MET	Mainchain
1	2-A	31	ARG	Sidechain
1	2-A	79	VAL	Mainchain
1	2-A	8	ILE	Mainchain
1	2-A	81	ASP	Sidechain
1	2-A	82	THR	Mainchain
1	2-A	93	LEU	Mainchain
1	2-A	98	VAL	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1-A	768	0	747	0	0
1	2-A	768	0	749	0	0
2	1-A	11	0	0	0	0
3	1-A	125	0	0	0	0
3	2-A	125	0	0	0	0
All	All	1797	0	1496	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1-A	93/103 (90%)	88 (95%)	4 (4%)	1 (1%)	11	0
1	2-A	93/103 (90%)	88 (95%)	5 (5%)	0	100	100
All	All	186/206 (90%)	176 (95%)	9 (5%)	1 (0%)	24	5

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	1-A	79	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1-A	86/93 (92%)	83 (96%)	3 (4%)	32	2
1	2-A	86/93 (92%)	83 (96%)	3 (4%)	32	2
All	All	172/186 (92%)	166 (96%)	6 (4%)	32	2

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	1-A	31	ARG
1	1-A	74	PRO
1	1-A	99	PHE
1	2-A	31	ARG
1	2-A	74	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	2-A	99	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	1-A	11	ASN
1	1-A	64	HIS
1	2-A	56	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	2-A	1
1	1-A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
2	A	55:SER	C	56:HIS	N	1.20
1	A	41:HIS	C	42:THR	N	1.16

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	1-A	95/103 (92%)	3.51	88 (92%) 0 0	3, 6, 12, 18	95 (100%)
1	2-A	0/103	-	-	-	-
All	All	95/206 (46%)	3.51	88 (92%) 0 0	3, 6, 12, 18	95 (100%)

All (88) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	1-A	99	PHE	12.3
1	1-A	33	ASN	9.0
1	1-A	98	VAL	7.9
1	1-A	5	PRO	6.8
1	1-A	15	ARG	6.2
1	1-A	47	PHE	5.9
1	1-A	50	GLY	5.4
1	1-A	13	GLY	5.1
1	1-A	6	PRO	5.1
1	1-A	36	VAL	4.9
1	1-A	70	ILE	4.9
1	1-A	73	ILE	4.9
1	1-A	8	ILE	4.7
1	1-A	90	TRP	4.7
1	1-A	67	SER	4.5
1	1-A	76	CYS	4.5
1	1-A	29	LYS	4.4
1	1-A	93	LEU	4.4
1	1-A	10	TRP	4.4
1	1-A	78	TYR	4.2
1	1-A	94	ILE	4.1
1	1-A	35	LYS	4.0
1	1-A	95	HIS	4.0
1	1-A	77	SER	4.0
1	1-A	65	ALA	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	1-A	31	ARG	3.8
1	1-A	40	VAL	3.8
1	1-A	44	VAL	3.8
1	1-A	45	PRO	3.7
1	1-A	39	LEU	3.6
1	1-A	49	ARG	3.6
1	1-A	20	GLU	3.6
1	1-A	32	ASN	3.6
1	1-A	82	THR	3.6
1	1-A	89	SER	3.6
1	1-A	96	SER	3.6
1	1-A	7	LYS	3.5
1	1-A	51	LEU	3.5
1	1-A	14	LYS	3.5
1	1-A	9	VAL	3.5
1	1-A	27	GLU	3.5
1	1-A	60	ALA	3.4
1	1-A	34	GLY	3.4
1	1-A	28	TYR	3.3
1	1-A	72	ILE	3.3
1	1-A	53	LEU	3.3
1	1-A	55	SER	3.3
1	1-A	86	ARG	3.3
1	1-A	79	VAL	3.2
1	1-A	37	MET	3.2
1	1-A	54	ALA	3.1
1	1-A	24	ALA	3.1
1	1-A	58	CYS	3.1
1	1-A	12	GLU	3.1
1	1-A	92	PRO	3.0
1	1-A	17	PHE	3.0
1	1-A	97	GLU	3.0
1	1-A	52	GLY	3.0
1	1-A	69	SER	2.9
1	1-A	25	PHE	2.9
1	1-A	81	ASP	2.9
1	1-A	19	THR	2.9
1	1-A	43	TYR	2.9
1	1-A	84	LEU	2.8
1	1-A	62	PHE	2.8
1	1-A	74	PRO	2.7
1	1-A	85	PRO	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	1-A	22	HIS	2.7
1	1-A	83	PHE	2.6
1	1-A	59	VAL	2.6
1	1-A	88	PRO	2.6
1	1-A	91	LYS	2.5
1	1-A	30	MET	2.5
1	1-A	75	SER	2.4
1	1-A	56	HIS	2.4
1	1-A	64	HIS	2.4
1	1-A	38	ASP	2.4
1	1-A	48	LYS	2.4
1	1-A	26	ILE	2.4
1	1-A	21	ASP	2.4
1	1-A	46	SER	2.4
1	1-A	42	THR	2.3
1	1-A	23	GLU	2.3
1	1-A	61	ALA	2.3
1	1-A	11	ASN	2.2
1	1-A	68	HIS	2.2
1	1-A	80	SER	2.2
1	1-A	57	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BR	1-A	200	1/1	0.81	0.26	44,44,44,44	1
3	BR	1-A	209	1/1	0.81	0.24	49,49,49,49	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BR	1-A	210	1/1	0.92	0.20	27,27,27,27	1
3	BR	1-A	204	1/1	0.95	0.28	22,22,22,22	1
3	BR	1-A	201	1/1	0.95	0.36	22,22,22,22	1
3	BR	1-A	202	1/1	0.95	0.35	21,21,21,21	1
3	BR	1-A	205	1/1	0.97	0.46	20,20,20,20	1
3	BR	1-A	207	1/1	0.97	0.30	19,19,19,19	1
3	BR	1-A	203	1/1	0.98	0.39	21,21,21,21	1
3	BR	1-A	208	1/1	0.98	0.41	20,20,20,20	1
3	BR	1-A	206	1/1	0.99	0.21	11,11,11,11	1

6.5 Other polymers [i](#)

There are no such residues in this entry.