



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:59 PM UTC

PDB ID : 2Q4A / pdb_00002q4a
Title : Ensemble refinement of the protein crystal structure of gene product from Arabidopsis thaliana At3g21360
Authors : Levin, E.J.; Kondrashov, D.A.; Wesenberg, G.E.; Phillips Jr., G.N.; Center for Eukaryotic Structural Genomics (CESG)
Deposited on : 2007-05-31
Resolution : 2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1-A	330	25% 97%
1	1-B	330	16% 97%
1	2-A	330	95%
1	2-B	330	95%
1	3-A	330	95%
1	3-B	330	95%
1	4-A	330	95%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	4-B	330	95%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 23772 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Clavamate synthase-like protein At3g21360.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	1-A	321	Total	C	N	O	S	0	0	0
			2565	1644	442	469	10			
1	2-A	321	Total	C	N	O	S	0	0	0
			2565	1644	442	469	10			
1	3-A	321	Total	C	N	O	S	0	0	0
			2565	1644	442	469	10			
1	4-A	321	Total	C	N	O	S	0	0	0
			2565	1644	442	469	10			
1	1-B	322	Total	C	N	O	S	0	0	0
			2576	1652	440	474	10			
1	2-B	322	Total	C	N	O	S	0	0	0
			2576	1652	440	474	10			
1	3-B	322	Total	C	N	O	S	0	0	0
			2576	1652	440	474	10			
1	4-B	322	Total	C	N	O	S	0	0	0
			2576	1652	440	474	10			

- Molecule 2 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	1-A	1	Total	Fe	0	0
			1	1		
2	2-A	1	Total	Fe	0	0
			1	1		
2	3-A	1	Total	Fe	0	0
			1	1		
2	4-A	1	Total	Fe	0	0
			1	1		
2	1-B	1	Total	Fe	0	0
			1	1		
2	2-B	1	Total	Fe	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	3-B	1	Total	Fe	0	0
			1	1		
2	4-B	1	Total	Fe	0	0
			1	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	1-A	1	Total	O	S	0	0
			5	4	1		
3	2-A	1	Total	O	S	0	0
			5	4	1		
3	3-A	1	Total	O	S	0	0
			5	4	1		
3	4-A	1	Total	O	S	0	0
			5	4	1		
3	1-A	1	Total	O	S	0	0
			5	4	1		
3	2-A	1	Total	O	S	0	0
			5	4	1		
3	3-A	1	Total	O	S	0	0
			5	4	1		
3	4-A	1	Total	O	S	0	0
			5	4	1		
3	1-A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	2-A	1	Total O S 5 4 1	0	0
3	3-A	1	Total O S 5 4 1	0	0
3	4-A	1	Total O S 5 4 1	0	0
3	1-B	1	Total O S 5 4 1	0	0
3	2-B	1	Total O S 5 4 1	0	0
3	3-B	1	Total O S 5 4 1	0	0
3	4-B	1	Total O S 5 4 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	1-A	339	Total O 339 339	0	0
4	2-A	342	Total O 342 342	0	0
4	3-A	338	Total O 338 338	0	0
4	4-A	336	Total O 336 336	0	0
4	1-B	441	Total O 441 441	0	0
4	2-B	438	Total O 438 438	0	0
4	3-B	442	Total O 442 442	0	0
4	4-B	444	Total O 444 444	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Clavamate synthase-like protein At3g21360




- Molecule 1: Clavamate synthase-like protein At3g21360




- Molecule 1: Clavamate synthase-like protein At3g21360




- Molecule 1: Clavamate synthase-like protein At3g21360





- Molecule 1: Clavamate synthase-like protein At3g21360

Chain 3-A:  95% ..




- Molecule 1: Clavamate synthase-like protein At3g21360

Chain 3-B:  95% ..





- Molecule 1: Clavamate synthase-like protein At3g21360

Chain 4-A:  95% ..



- Molecule 1: Clavamate synthase-like protein At3g21360

Chain 4-B:  95% ..



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	145.28Å 61.09Å 114.68Å 90.00° 121.68° 90.00°	Depositor
Resolution (Å)	43.63 – 2.39 43.63 – 2.39	Depositor EDS
% Data completeness (in resolution range)	96.8 (43.63-2.39) 96.6 (43.63-2.39)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.48 (at 2.39Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.171 , 0.235 0.170 , 0.231	Depositor DCC
R_{free} test set	1628 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	23.4	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.09 , 76.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	23772	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	1-A	0.40	0/2583	0.66	1/3502 (0.0%)
1	1-B	0.46	0/2651	0.69	3/3605 (0.1%)
1	2-A	0.40	0/2640	0.66	1/3588 (0.0%)
1	2-B	0.46	0/2651	0.69	3/3605 (0.1%)
1	3-A	0.40	0/2640	0.66	1/3588 (0.0%)
1	3-B	0.46	0/2651	0.69	3/3605 (0.1%)
1	4-A	0.40	0/2479	0.66	1/3367 (0.0%)
1	4-B	0.46	0/2651	0.69	3/3605 (0.1%)
All	All	0.43	0/20946	0.68	16/28465 (0.1%)

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	1-B	72	ASN	N-CA-C	6.39	120.70	112.26
1	2-B	72	ASN	N-CA-C	6.39	120.70	112.26
1	3-B	72	ASN	N-CA-C	6.39	120.70	112.26
1	4-B	72	ASN	N-CA-C	6.39	120.70	112.26
1	1-B	121	HIS	N-CA-C	-5.96	100.84	110.32
1	2-B	121	HIS	N-CA-C	-5.96	100.84	110.32
1	3-B	121	HIS	N-CA-C	-5.96	100.84	110.32
1	4-B	121	HIS	N-CA-C	-5.96	100.84	110.32
1	1-A	121	HIS	N-CA-C	-5.84	102.27	110.50
1	2-A	121	HIS	N-CA-C	-5.84	102.27	110.50
1	3-A	121	HIS	N-CA-C	-5.84	102.27	110.50
1	4-A	121	HIS	N-CA-C	-5.84	102.27	110.50
1	1-B	192	GLY	N-CA-C	5.00	120.21	114.16
1	2-B	192	GLY	N-CA-C	5.00	120.21	114.16
1	3-B	192	GLY	N-CA-C	5.00	120.21	114.16
1	4-B	192	GLY	N-CA-C	5.00	120.21	114.16

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1-A	2565	0	0	0	0
1	1-B	2576	0	0	0	0
1	2-A	2565	0	2519	0	0
1	2-B	2576	0	2528	0	0
1	3-A	2565	0	0	0	0
1	3-B	2576	0	0	0	0
1	4-A	2565	0	1822	0	0
1	4-B	2576	0	0	0	0
2	1-A	1	0	0	0	0
2	1-B	1	0	0	0	0
2	2-A	1	0	0	0	0
2	2-B	1	0	0	0	0
2	3-A	1	0	0	0	0
2	3-B	1	0	0	0	0
2	4-A	1	0	0	0	0
2	4-B	1	0	0	0	0
3	1-A	15	0	0	0	0
3	1-B	5	0	0	0	0
3	2-A	15	0	0	0	0
3	2-B	5	0	0	0	0
3	3-A	15	0	0	0	0
3	3-B	5	0	0	0	0
3	4-A	15	0	0	0	0
3	4-B	5	0	0	0	0
4	1-A	339	0	0	0	0
4	1-B	441	0	0	0	0
4	2-A	342	0	0	0	0
4	2-B	438	0	0	0	0
4	3-A	338	0	0	0	0
4	3-B	442	0	0	0	0
4	4-A	336	0	0	0	0
4	4-B	444	0	0	0	0
All	All	23772	0	6869	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). Clashscore could not be calculated for this entry.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	2-A	317/330 (96%)	293 (92%)	21 (7%)	3 (1%)	14	22
1	2-B	318/330 (96%)	298 (94%)	19 (6%)	1 (0%)	36	50
1	3-A	317/330 (96%)	285 (90%)	29 (9%)	3 (1%)	14	22
1	3-B	318/330 (96%)	296 (93%)	20 (6%)	2 (1%)	21	32
1	4-A	297/330 (90%)	273 (92%)	22 (7%)	2 (1%)	18	28
1	4-B	318/330 (96%)	295 (93%)	20 (6%)	3 (1%)	14	22
All	All	1885/1980 (95%)	1740 (92%)	131 (7%)	14 (1%)	18	28

All (14) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	4-A	92	VAL
1	4-B	14	GLN
1	4-B	247	TRP
1	3-A	220	TRP
1	3-B	146	GLU
1	4-A	70	PRO
1	2-A	172	GLU
1	4-B	221	THR
1	2-A	223	ASP
1	2-B	208	GLU
1	3-A	47	ILE
1	3-B	31	ALA
1	3-A	13	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	2-A	10	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	2-A	281/288 (98%)	276 (98%)	5 (2%)	51	73
1	2-B	284/288 (99%)	280 (99%)	4 (1%)	59	79
1	3-A	281/288 (98%)	276 (98%)	5 (2%)	51	73
1	3-B	284/288 (99%)	280 (99%)	4 (1%)	59	79
1	4-A	265/288 (92%)	260 (98%)	5 (2%)	50	71
1	4-B	284/288 (99%)	280 (99%)	4 (1%)	59	79
All	All	1679/1728 (97%)	1652 (98%)	27 (2%)	55	76

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	2-A	5	LEU
1	2-A	45	GLN
1	2-A	110	GLU
1	2-A	112	PRO
1	2-A	306	ILE
1	2-B	26	ILE
1	2-B	176	LEU
1	2-B	181	LEU
1	2-B	262	ARG
1	3-A	5	LEU
1	3-A	45	GLN
1	3-A	110	GLU
1	3-A	112	PRO
1	3-A	306	ILE
1	3-B	26	ILE
1	3-B	176	LEU
1	3-B	181	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	3-B	262	ARG
1	4-A	5	LEU
1	4-A	45	GLN
1	4-A	110	GLU
1	4-A	112	PRO
1	4-A	306	ILE
1	4-B	26	ILE
1	4-B	176	LEU
1	4-B	181	LEU
1	4-B	262	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	2-A	45	GLN
1	2-A	59	HIS
1	2-A	125	GLN
1	2-A	153	HIS
1	2-A	173	HIS
1	2-B	153	HIS
1	2-B	299	GLN
1	2-B	313	HIS
1	4-A	14	GLN
1	4-A	72	ASN
1	4-A	125	GLN
1	4-A	153	HIS
1	4-A	243	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 8 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	3-A	801	-	4,4,4	0.32	0	6,6,6	0.07	0
3	SO4	3-B	804	-	4,4,4	0.37	0	6,6,6	0.16	0
3	SO4	4-A	803	-	4,4,4	0.41	0	6,6,6	0.17	0
3	SO4	4-B	804	-	4,4,4	0.37	0	6,6,6	0.18	0
3	SO4	1-B	804	-	4,4,4	0.35	0	6,6,6	0.16	0
3	SO4	2-A	801	-	4,4,4	0.39	0	6,6,6	0.15	0
3	SO4	1-A	803	-	4,4,4	0.42	0	6,6,6	0.19	0
3	SO4	3-A	803	-	4,4,4	0.41	0	6,6,6	0.19	0
3	SO4	2-A	803	-	4,4,4	0.41	0	6,6,6	0.18	0
3	SO4	2-B	804	-	4,4,4	0.40	0	6,6,6	0.13	0
3	SO4	1-A	802	-	4,4,4	0.40	0	6,6,6	0.11	0
3	SO4	3-A	802	-	4,4,4	0.44	0	6,6,6	0.07	0
3	SO4	4-A	801	-	4,4,4	0.32	0	6,6,6	0.08	0
3	SO4	4-A	802	-	4,4,4	0.41	0	6,6,6	0.06	0
3	SO4	2-A	802	-	4,4,4	0.41	0	6,6,6	0.05	0
3	SO4	1-A	801	-	4,4,4	0.35	0	6,6,6	0.11	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	1-A	321/330 (97%)	1.42	84 (26%) ⓘ ⓘ	2, 6, 19, 22	321 (100%)
1	1-B	322/330 (97%)	1.00	54 (16%) ⓘ ⓘ	1, 5, 14, 24	322 (100%)
1	2-A	0/330	-	-	-	-
1	2-B	0/330	-	-	-	-
1	3-A	0/330	-	-	-	-
1	3-B	0/330	-	-	-	-
1	4-A	0/330	-	-	-	-
1	4-B	0/330	-	-	-	-
All	All	643/2640 (24%)	1.21	138 (21%) ⓘ ⓘ	1, 6, 17, 24	643 (100%)

All (138) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	1-A	4	LEU	6.6
1	1-A	36	PRO	5.3
1	1-A	91	TYR	5.2
1	1-A	218	LEU	5.1
1	1-B	91	TYR	4.8
1	1-A	108	ALA	4.8
1	1-B	37	ALA	4.7
1	1-A	206	LEU	4.7
1	1-A	5	LEU	4.6
1	1-A	143	CYS	4.5
1	1-A	202	HIS	4.3
1	1-A	201	THR	4.3
1	1-A	200	LEU	4.3
1	1-A	97	PRO	4.2
1	1-A	216	MET	4.2
1	1-A	221	THR	4.1
1	1-A	109	ASN	4.0
1	1-B	143	CYS	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	1-A	92	VAL	3.9
1	1-A	95	ALA	3.9
1	1-B	31	ALA	3.9
1	1-A	214	LEU	3.9
1	1-A	199	PHE	3.7
1	1-A	93	GLY	3.7
1	1-B	32	SER	3.7
1	1-B	111	SER	3.7
1	1-A	96	ALA	3.6
1	1-B	108	ALA	3.6
1	1-B	224	GLY	3.6
1	1-B	201	THR	3.5
1	1-A	224	GLY	3.5
1	1-A	283	HIS	3.5
1	1-B	262	ARG	3.5
1	1-A	163	HIS	3.5
1	1-B	202	HIS	3.5
1	1-A	162	LYS	3.4
1	1-B	329	CYS	3.4
1	1-A	94	GLY	3.4
1	1-A	217	LYS	3.3
1	1-A	37	ALA	3.3
1	1-A	219	GLU	3.2
1	1-B	109	ASN	3.1
1	1-B	203	ASP	3.1
1	1-A	207	ALA	3.1
1	1-B	221	THR	3.0
1	1-A	223	ASP	3.0
1	1-A	211	ALA	3.0
1	1-B	218	LEU	3.0
1	1-A	98	ARG	3.0
1	1-A	107	THR	3.0
1	1-A	274	GLY	2.9
1	1-B	38	LEU	2.9
1	1-A	171	GLU	2.9
1	1-A	213	ASP	2.9
1	1-B	85	GLY	2.9
1	1-A	220	TRP	2.9
1	1-A	52	HIS	2.9
1	1-A	179	ARG	2.9
1	1-B	223	ASP	2.9
1	1-A	189	SER	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	1-A	110	GLU	2.8
1	1-B	171	GLU	2.8
1	1-B	186	ASP	2.8
1	1-B	261	LYS	2.8
1	1-A	282	VAL	2.8
1	1-A	99	THR	2.8
1	1-B	207	ALA	2.8
1	1-B	216	MET	2.8
1	1-A	82	GLU	2.7
1	1-A	167	VAL	2.7
1	1-A	186	ASP	2.7
1	1-B	266	ARG	2.7
1	1-A	212	VAL	2.7
1	1-A	319	ASP	2.7
1	1-A	208	GLU	2.7
1	1-B	217	LYS	2.7
1	1-B	68	GLY	2.7
1	1-A	19	SER	2.7
1	1-B	219	GLU	2.6
1	1-B	105	VAL	2.6
1	1-B	107	THR	2.6
1	1-A	209	GLN	2.6
1	1-A	190	PRO	2.6
1	1-B	181	LEU	2.6
1	1-A	187	PRO	2.6
1	1-A	168	GLN	2.5
1	1-A	203	ASP	2.5
1	1-A	188	SER	2.5
1	1-A	305	LEU	2.5
1	1-B	180	VAL	2.4
1	1-A	128	GLU	2.4
1	1-A	146	GLU	2.4
1	1-A	222	GLU	2.4
1	1-A	228	THR	2.4
1	1-B	58	LEU	2.4
1	1-B	222	GLU	2.4
1	1-B	28	PRO	2.4
1	1-B	205	ASN	2.4
1	1-A	144	GLY	2.4
1	1-B	2	ALA	2.4
1	1-A	193	ARG	2.4
1	1-A	140	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	1-A	165	GLU	2.4
1	1-B	194	GLY	2.3
1	1-B	274	GLY	2.3
1	1-A	181	LEU	2.3
1	1-A	180	VAL	2.3
1	1-B	258	TRP	2.3
1	1-A	281	ILE	2.3
1	1-A	284	ASP	2.2
1	1-B	133	LEU	2.2
1	1-B	184	ASP	2.2
1	1-B	18	GLU	2.2
1	1-B	209	GLN	2.2
1	1-B	213	ASP	2.2
1	1-B	70	PRO	2.2
1	1-B	35	ILE	2.2
1	1-B	60	GLU	2.2
1	1-A	215	GLY	2.2
1	1-A	225	GLY	2.2
1	1-A	285	CYS	2.2
1	1-B	204	LYS	2.1
1	1-A	90	PRO	2.1
1	1-A	204	LYS	2.1
1	1-A	288	ILE	2.1
1	1-B	206	LEU	2.1
1	1-A	271	PHE	2.1
1	1-A	100	SER	2.1
1	1-A	253	ALA	2.1
1	1-A	236	ILE	2.1
1	1-A	115	GLN	2.1
1	1-B	33	ILE	2.1
1	1-B	199	PHE	2.1
1	1-B	314	SER	2.0
1	1-A	255	TYR	2.0
1	1-A	83	ALA	2.0
1	1-B	211	ALA	2.0
1	1-A	176	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	1-A	802	5/5	0.69	0.19	74,74,74,74	5
2	FE	2-A	901	1/1	-	-	36,36,36,36	1
2	FE	3-A	901	1/1	-	-	36,36,36,36	1
2	FE	4-A	901	1/1	-	-	36,36,36,36	1
3	SO4	1-A	801	5/5	0.77	0.26	73,75,75,75	5
2	FE	2-B	902	1/1	-	-	26,26,26,26	1
2	FE	3-B	902	1/1	-	-	5,5,5,5	1
2	FE	4-B	902	1/1	-	-	16,16,16,16	1
3	SO4	1-A	803	5/5	0.82	0.19	45,46,47,49	5
3	SO4	2-A	801	5/5	-	-	76,76,77,77	5
3	SO4	3-A	801	5/5	-	-	73,73,74,75	5
3	SO4	4-A	801	5/5	-	-	72,73,74,74	5
3	SO4	1-B	804	5/5	0.87	0.16	40,42,44,44	5
3	SO4	2-A	802	5/5	-	-	75,75,75,76	5
3	SO4	3-A	802	5/5	-	-	76,76,77,77	5
3	SO4	4-A	802	5/5	-	-	74,74,75,75	5
2	FE	1-B	902	1/1	0.88	0.10	26,26,26,26	1
3	SO4	2-A	803	5/5	-	-	47,48,49,50	5
3	SO4	3-A	803	5/5	-	-	46,47,48,49	5
3	SO4	4-A	803	5/5	-	-	42,43,44,44	5
2	FE	1-A	901	1/1	0.91	0.07	34,34,34,34	1
3	SO4	2-B	804	5/5	-	-	46,47,47,48	5
3	SO4	3-B	804	5/5	-	-	41,43,45,45	5
3	SO4	4-B	804	5/5	-	-	40,42,44,44	5

6.5 Other polymers [i](#)

There are no such residues in this entry.