



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:06 AM UTC

PDB ID : 2Q4L / pdb_00002q4l
Title : Ensemble refinement of the crystal structure of GALT-like protein from Arabidopsis thaliana At5g18200
Authors : Levin, E.J.; Kondrashov, D.A.; Wesenberg, G.E.; Phillips Jr., G.N.; Center for Eukaryotic Structural Genomics (CESG)
Deposited on : 2007-05-31
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1-A	351	17% 81%5%14%
1	1-B	351	15% 81%.15%
1	2-A	351	81%.14%
1	2-B	351	81%.15%
1	3-A	351	81%.14%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	3-B	351	 80%5%15%
1	4-A	351	 81%5%14%
1	4-B	351	 81%•15%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 19940 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable galactose-1-phosphate uridyl transferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	1-A	301	Total	C	N	O	S	0	0	0
			2388	1536	403	436	13			
1	2-A	301	Total	C	N	O	S	0	0	0
			2388	1536	403	436	13			
1	3-A	301	Total	C	N	O	S	0	0	0
			2388	1536	403	436	13			
1	4-A	301	Total	C	N	O	S	0	0	0
			2388	1536	403	436	13			
1	1-B	299	Total	C	N	O	S	0	0	0
			2368	1524	397	434	13			
1	2-B	299	Total	C	N	O	S	0	0	0
			2368	1524	397	434	13			
1	3-B	299	Total	C	N	O	S	0	0	0
			2368	1524	397	434	13			
1	4-B	299	Total	C	N	O	S	0	0	0
			2368	1524	397	434	13			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	1-A	2	Total	Zn	0	0
			2	2		
2	2-A	2	Total	Zn	0	0
			2	2		
2	3-A	2	Total	Zn	0	0
			2	2		
2	4-A	2	Total	Zn	0	0
			2	2		
2	1-B	2	Total	Zn	0	0
			2	2		
2	2-B	2	Total	Zn	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	3-B	2	Total 2	Zn 2	0	0
2	4-B	2	Total 2	Zn 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	1-A	109	Total 109	O 109	0	0
3	2-A	114	Total 114	O 114	0	0
3	3-A	112	Total 112	O 112	0	0
3	4-A	114	Total 114	O 114	0	0
3	1-B	116	Total 116	O 116	0	0
3	2-B	111	Total 111	O 111	0	0
3	3-B	113	Total 113	O 113	0	0
3	4-B	111	Total 111	O 111	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Probable galactose-1-phosphate uridyl transferase



- Molecule 1: Probable galactose-1-phosphate uridyl transferase




- Molecule 1: Probable galactose-1-phosphate uridyl transferase




- Molecule 1: Probable galactose-1-phosphate uridyl transferase

Chain 2-B:  81% 15%




- Molecule 1: Probable galactose-1-phosphate uridyl transferase

Chain 3-A:  81% 14%




- Molecule 1: Probable galactose-1-phosphate uridyl transferase

Chain 3-B:  80% 5% 15%




- Molecule 1: Probable galactose-1-phosphate uridyl transferase

Chain 4-A:  81% 5% 14%



- Molecule 1: Probable galactose-1-phosphate uridyl transferase

Chain 4-B:  81% 15%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.98Å 95.53Å 110.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.26 – 2.30 27.26 – 2.30	Depositor EDS
% Data completeness (in resolution range)	89.8 (27.26-2.30) 89.8 (27.26-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.79 (at 2.31Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.190 , 0.270 0.198 , 0.270	Depositor DCC
R_{free} test set	2581 reflections (9.37%)	wwPDB-VP
Wilson B-factor (Å ²)	39.9	Xtriage
Anisotropy	0.447	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.08 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	19940	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	1-A	0.27	0/2454	0.59	1/3331 (0.0%)
1	1-B	0.26	0/2433	0.63	1/3306 (0.0%)
1	2-A	0.27	0/2454	0.59	1/3331 (0.0%)
1	2-B	0.26	0/2433	0.63	1/3306 (0.0%)
1	3-A	0.27	0/2454	0.59	1/3331 (0.0%)
1	3-B	0.26	0/2433	0.63	1/3306 (0.0%)
1	4-A	0.27	0/2454	0.59	1/3331 (0.0%)
1	4-B	0.26	0/2433	0.63	1/3306 (0.0%)
All	All	0.27	0/19548	0.61	8/26548 (0.0%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	1-B	98	ALA	N-CA-C	-5.27	105.21	111.69
1	2-B	98	ALA	N-CA-C	-5.27	105.21	111.69
1	3-B	98	ALA	N-CA-C	-5.27	105.21	111.69
1	4-B	98	ALA	N-CA-C	-5.27	105.21	111.69
1	1-A	330	GLY	N-CA-C	-5.15	108.14	115.30
1	2-A	330	GLY	N-CA-C	-5.15	108.14	115.30
1	3-A	330	GLY	N-CA-C	-5.15	108.14	115.30
1	4-A	330	GLY	N-CA-C	-5.15	108.14	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1-A	2388	0	2353	0	0
1	1-B	2368	0	2325	0	0
1	2-A	2388	0	2355	0	0
1	2-B	2368	0	2326	0	0
1	3-A	2388	0	2353	0	0
1	3-B	2368	0	2325	0	0
1	4-A	2388	0	2354	0	0
1	4-B	2368	0	2325	0	0
2	1-A	2	0	0	0	0
2	1-B	2	0	0	0	0
2	2-A	2	0	0	0	0
2	2-B	2	0	0	0	0
2	3-A	2	0	0	0	0
2	3-B	2	0	0	0	0
2	4-A	2	0	0	0	0
2	4-B	2	0	0	0	0
3	1-A	109	0	0	0	0
3	1-B	116	0	0	0	0
3	2-A	114	0	0	0	0
3	2-B	111	0	0	0	0
3	3-A	112	0	0	0	0
3	3-B	113	0	0	0	0
3	4-A	114	0	0	0	0
3	4-B	111	0	0	0	0
All	All	19940	0	18716	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1-A	293/351 (84%)	253 (86%)	33 (11%)	7 (2%)	4	4
1	1-B	289/351 (82%)	255 (88%)	28 (10%)	6 (2%)	5	4
1	2-A	293/351 (84%)	262 (89%)	26 (9%)	5 (2%)	7	7
1	2-B	289/351 (82%)	249 (86%)	36 (12%)	4 (1%)	9	9
1	3-A	293/351 (84%)	262 (89%)	26 (9%)	5 (2%)	7	7
1	3-B	289/351 (82%)	256 (89%)	25 (9%)	8 (3%)	4	2
1	4-A	293/351 (84%)	254 (87%)	32 (11%)	7 (2%)	4	4
1	4-B	289/351 (82%)	255 (88%)	30 (10%)	4 (1%)	9	9
All	All	2328/2808 (83%)	2046 (88%)	236 (10%)	46 (2%)	6	5

All (46) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	1-B	83	HIS
1	1-B	104	GLU
1	3-A	137	LEU
1	3-B	142	PRO
1	1-A	50	SER
1	1-A	83	HIS
1	1-A	93	GLU
1	1-A	284	GLN
1	1-B	176	ALA
1	2-A	104	GLU
1	2-A	286	ASN
1	3-A	83	HIS
1	3-B	83	HIS
1	3-B	104	GLU
1	3-B	255	HIS
1	4-A	134	SER
1	4-A	160	ALA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	4-A	299	LEU
1	1-A	104	GLU
1	1-B	222	LYS
1	2-A	32	ASN
1	2-A	83	HIS
1	2-A	217	CYS
1	2-B	235	VAL
1	2-B	264	ASP
1	3-A	104	GLU
1	4-A	221	ALA
1	4-B	154	LYS
1	1-A	70	GLU
1	2-B	83	HIS
1	4-A	120	GLY
1	4-B	104	GLU
1	4-B	260	HIS
1	1-A	181	SER
1	1-B	64	PRO
1	3-A	132	VAL
1	3-B	217	CYS
1	4-B	120	GLY
1	3-B	120	GLY
1	3-B	220	GLU
1	4-A	239	PRO
1	1-B	120	GLY
1	2-B	120	GLY
1	4-A	235	VAL
1	3-A	131	PRO
1	3-B	143	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1-A	270/312 (86%)	260 (96%)	10 (4%)	30	45
1	1-B	269/312 (86%)	261 (97%)	8 (3%)	36	53

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	2-A	270/312 (86%)	260 (96%)	10 (4%)	30	45
1	2-B	269/312 (86%)	261 (97%)	8 (3%)	36	53
1	3-A	270/312 (86%)	260 (96%)	10 (4%)	30	45
1	3-B	269/312 (86%)	261 (97%)	8 (3%)	36	53
1	4-A	270/312 (86%)	260 (96%)	10 (4%)	30	45
1	4-B	269/312 (86%)	261 (97%)	8 (3%)	36	53
All	All	2156/2496 (86%)	2084 (97%)	72 (3%)	33	50

All (72) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	1-A	25	LEU
1	1-A	83	HIS
1	1-A	91	VAL
1	1-A	166	ASN
1	1-A	185	SER
1	1-A	219	CYS
1	1-A	237	VAL
1	1-A	258	HIS
1	1-A	293	MET
1	1-A	348	VAL
1	1-B	155	ARG
1	1-B	163	ASP
1	1-B	190	MET
1	1-B	258	HIS
1	1-B	262	LEU
1	1-B	285	LEU
1	1-B	292	TYR
1	1-B	293	MET
1	2-A	25	LEU
1	2-A	83	HIS
1	2-A	91	VAL
1	2-A	166	ASN
1	2-A	185	SER
1	2-A	219	CYS
1	2-A	237	VAL
1	2-A	258	HIS
1	2-A	293	MET
1	2-A	348	VAL
1	2-B	155	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	2-B	163	ASP
1	2-B	190	MET
1	2-B	258	HIS
1	2-B	262	LEU
1	2-B	285	LEU
1	2-B	292	TYR
1	2-B	293	MET
1	3-A	25	LEU
1	3-A	83	HIS
1	3-A	91	VAL
1	3-A	166	ASN
1	3-A	185	SER
1	3-A	219	CYS
1	3-A	237	VAL
1	3-A	258	HIS
1	3-A	293	MET
1	3-A	348	VAL
1	3-B	155	ARG
1	3-B	163	ASP
1	3-B	190	MET
1	3-B	258	HIS
1	3-B	262	LEU
1	3-B	285	LEU
1	3-B	292	TYR
1	3-B	293	MET
1	4-A	25	LEU
1	4-A	83	HIS
1	4-A	91	VAL
1	4-A	166	ASN
1	4-A	185	SER
1	4-A	219	CYS
1	4-A	237	VAL
1	4-A	258	HIS
1	4-A	293	MET
1	4-A	348	VAL
1	4-B	155	ARG
1	4-B	163	ASP
1	4-B	190	MET
1	4-B	258	HIS
1	4-B	262	LEU
1	4-B	285	LEU
1	4-B	292	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	4-B	293	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (42) such sidechains are listed below:

Mol	Chain	Res	Type
1	1-A	94	ASN
1	1-A	158	GLN
1	1-A	162	HIS
1	1-A	166	ASN
1	1-A	174	GLN
1	1-A	291	ASN
1	1-B	86	ASN
1	1-B	133	HIS
1	1-B	136	GLN
1	1-B	162	HIS
1	1-B	174	GLN
1	1-B	286	ASN
1	2-A	32	ASN
1	2-A	158	GLN
1	2-A	305	GLN
1	2-B	136	GLN
1	2-B	169	GLN
1	2-B	173	ASN
1	2-B	174	GLN
1	2-B	286	ASN
1	3-A	83	HIS
1	3-A	161	GLN
1	3-A	169	GLN
1	3-A	260	HIS
1	3-A	279	GLN
1	3-A	284	GLN
1	3-A	291	ASN
1	3-B	21	GLN
1	3-B	158	GLN
1	3-B	161	GLN
1	3-B	166	ASN
1	3-B	279	GLN
1	3-B	284	GLN
1	4-A	174	GLN
1	4-A	314	GLN
1	4-B	136	GLN
1	4-B	161	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	4-B	174	GLN
1	4-B	233	HIS
1	4-B	260	HIS
1	4-B	291	ASN
1	4-B	318	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 16 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	1-A	301/351 (85%)	1.27	60 (19%) 3 3	6, 10, 19, 23	301 (100%)
1	1-B	299/351 (85%)	1.25	51 (17%) 4 5	5, 10, 18, 21	299 (100%)
1	2-A	0/351	-	-	-	-
1	2-B	0/351	-	-	-	-
1	3-A	0/351	-	-	-	-
1	3-B	0/351	-	-	-	-
1	4-A	0/351	-	-	-	-
1	4-B	0/351	-	-	-	-
All	All	600/2808 (21%)	1.26	111 (18%) 3 4	5, 10, 19, 23	600 (100%)

All (111) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	1-A	135	ILE	5.7
1	1-B	130	SER	5.6
1	1-A	292	TYR	5.5
1	1-A	67	ILE	5.1
1	1-B	105	THR	4.4
1	1-A	87	TRP	4.3
1	1-A	137	LEU	4.3
1	1-B	182	MET	4.2
1	1-A	140	ILE	4.0
1	1-B	67	ILE	3.9
1	1-B	148	ILE	3.9
1	1-B	143	VAL	3.9
1	1-A	228	ILE	3.7
1	1-B	49	LYS	3.7
1	1-B	83	HIS	3.7
1	1-B	104	GLU	3.6
1	1-B	240	PHE	3.6
1	1-A	257	SER	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	1-B	46	THR	3.4
1	1-A	227	VAL	3.4
1	1-A	48	PHE	3.4
1	1-A	190	MET	3.4
1	1-A	83	HIS	3.4
1	1-B	176	ALA	3.4
1	1-A	70	GLU	3.3
1	1-B	25	LEU	3.3
1	1-B	48	PHE	3.3
1	1-A	95	LEU	3.3
1	1-B	325	PHE	3.3
1	1-B	340	ASP	3.3
1	1-A	226	PHE	3.2
1	1-B	80	VAL	3.2
1	1-A	166	ASN	3.2
1	1-B	87	TRP	3.1
1	1-B	66	CYS	3.1
1	1-B	191	ALA	3.1
1	1-B	350	LEU	3.1
1	1-A	233	HIS	3.0
1	1-A	96	TYR	3.0
1	1-A	162	HIS	3.0
1	1-B	65	PHE	3.0
1	1-A	73	CYS	2.9
1	1-A	351	THR	2.9
1	1-A	85	PRO	2.9
1	1-B	138	SER	2.9
1	1-B	261	HIS	2.9
1	1-A	337	PHE	2.8
1	1-B	147	ASP	2.8
1	1-A	336	VAL	2.8
1	1-A	105	THR	2.7
1	1-A	50	SER	2.7
1	1-B	343	LYS	2.6
1	1-A	262	LEU	2.6
1	1-B	275	LYS	2.6
1	1-A	348	VAL	2.6
1	1-A	313	LEU	2.6
1	1-A	350	LEU	2.5
1	1-B	103	LEU	2.5
1	1-A	23	PRO	2.5
1	1-B	64	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	1-B	328	GLY	2.5
1	1-A	86	ASN	2.5
1	1-B	88	LYS	2.5
1	1-B	62	SER	2.5
1	1-B	72	GLU	2.5
1	1-A	39	PRO	2.5
1	1-B	81	PRO	2.5
1	1-A	183	SER	2.4
1	1-B	181	SER	2.4
1	1-B	208	ASP	2.4
1	1-A	134	SER	2.4
1	1-A	182	MET	2.4
1	1-A	254	ASP	2.4
1	1-A	224	LYS	2.4
1	1-B	281	ILE	2.4
1	1-A	215	LYS	2.3
1	1-B	282	ALA	2.3
1	1-B	135	ILE	2.3
1	1-A	65	PHE	2.3
1	1-A	163	ASP	2.3
1	1-B	292	TYR	2.3
1	1-A	130	SER	2.3
1	1-A	234	PHE	2.3
1	1-B	134	SER	2.3
1	1-B	21	GLN	2.3
1	1-B	119	VAL	2.3
1	1-B	277	MET	2.2
1	1-A	181	SER	2.2
1	1-B	276	LEU	2.2
1	1-A	64	PRO	2.2
1	1-B	146	GLY	2.2
1	1-A	251	ILE	2.2
1	1-A	62	SER	2.2
1	1-A	132	VAL	2.2
1	1-A	189	MET	2.2
1	1-A	345	MET	2.2
1	1-A	49	LYS	2.2
1	1-A	136	GLN	2.2
1	1-B	238	ALA	2.2
1	1-A	332	TYR	2.1
1	1-A	186	HIS	2.1
1	1-B	95	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	1-B	133	HIS	2.1
1	1-A	298	PRO	2.1
1	1-A	216	CYS	2.0
1	1-A	199	VAL	2.0
1	1-B	144	GLY	2.0
1	1-A	297	SER	2.0
1	1-B	184	HIS	2.0
1	1-A	335	PRO	2.0
1	1-A	222	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	1-A	401	1/1	0.71	0.13	42,42,42,42	1
2	ZN	2-A	400	1/1	-	-	52,52,52,52	1
2	ZN	3-A	400	1/1	-	-	55,55,55,55	1
2	ZN	4-A	400	1/1	-	-	49,49,49,49	1
2	ZN	1-B	402	1/1	0.84	0.10	60,60,60,60	1
2	ZN	2-A	401	1/1	-	-	25,25,25,25	1
2	ZN	3-A	401	1/1	-	-	40,40,40,40	1
2	ZN	4-A	401	1/1	-	-	32,32,32,32	1
2	ZN	1-A	400	1/1	0.95	0.04	53,53,53,53	1
2	ZN	2-B	402	1/1	-	-	55,55,55,55	1
2	ZN	3-B	402	1/1	-	-	53,53,53,53	1
2	ZN	4-B	402	1/1	-	-	60,60,60,60	1
2	ZN	1-B	403	1/1	0.95	0.04	36,36,36,36	1
2	ZN	2-B	403	1/1	-	-	26,26,26,26	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	3-B	403	1/1	-	-	32,32,32,32	1
2	ZN	4-B	403	1/1	-	-	35,35,35,35	1

6.5 Other polymers [i](#)

There are no such residues in this entry.