



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 11:27 AM UTC

PDB ID : 5Q4V / pdb_00005q4v
Title : PanDDA analysis group deposition – Crystal Structure of DCLRE1A after initial refinement with no ligand modelled (structure 96)
Authors : Newman, J.A.; Aitkenhead, H.; Lee, S.Y.; Kupinska, K.; Burgess-Brown, N.; Tallon, R.; Krojer, T.; von Delft, F.; Arrowsmith, C.H.; Edwards, A.; Bountra, C.; Gileadi, O.
Deposited on : 2017-05-25
Resolution : 1.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	47% 87% 12% ..

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3044 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DCLRE1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	341	2726	1765	450	491	20	0	4	0

- Molecule 2 is MALONATE ION (CCD ID: MLI) (formula: $C_3H_2O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	7	3	4	0	0

- Molecule 3 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ni	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	310	Total 310	O 310	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.87Å 58.04Å 114.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.26 – 1.59 57.26 – 1.59	Depositor EDS
% Data completeness (in resolution range)	99.9 (57.26-1.59) 99.9 (57.26-1.59)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 1.59Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.297 , 0.344 0.309 , 0.354	Depositor DCC
R_{free} test set	2280 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	21.2	Xtriage
Anisotropy	0.252	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 34.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3044	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	0/2802	1.08	4/3809 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	860	ASN	N-CA-C	6.55	118.11	110.97
1	A	804	LEU	CA-C-N	6.16	125.64	119.24
1	A	804	LEU	C-N-CA	6.16	125.64	119.24
1	A	830	GLN	N-CA-C	6.01	117.99	108.79

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2726	0	2676	23	0
2	A	7	0	2	0	0
3	A	1	0	0	0	0
4	A	310	0	0	10	0
All	All	3044	0	2678	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:909:GLN:NE2	4:A:1201:HOH:O	2.04	0.91
1:A:1032:ARG:NH2	4:A:1208:HOH:O	2.34	0.58
1:A:1032:ARG:CZ	4:A:1225:HOH:O	2.55	0.55
1:A:973:ILE:HG23	4:A:1447:HOH:O	2.07	0.54
1:A:834:MET:HE2	1:A:1013:ILE:HD11	1.91	0.53
1:A:755:THR:O	1:A:759:LEU:HG	2.09	0.53
1:A:742:SER:O	1:A:765:VAL:HG22	2.09	0.53
1:A:743:LYS:HG3	1:A:744:HIS:CE1	2.50	0.47
1:A:815:ASP:OD1	4:A:1202:HOH:O	2.19	0.47
1:A:1032:ARG:NE	4:A:1225:HOH:O	2.48	0.46
1:A:717:PHE:CE1	1:A:728:TYR:HB3	2.53	0.44
1:A:859:ILE:HD11	1:A:889:ALA:HB1	2.00	0.44
1:A:860:ASN:O	1:A:861:THR:C	2.61	0.44
1:A:874:VAL:HG11	1:A:890:ILE:HG21	2.01	0.43
1:A:937:ILE:HG21	1:A:960:ARG:HG2	2.00	0.43
1:A:856:ARG:HE	1:A:856:ARG:HA	1.83	0.43
1:A:760:LYS:O	1:A:764[A]:HIS:HA	2.19	0.42
1:A:875:VAL:HG23	1:A:956:ILE:HG23	2.01	0.42
1:A:702:PRO:HG2	1:A:705:LYS:HG3	2.01	0.42
1:A:973:ILE:HG22	4:A:1236:HOH:O	2.19	0.41
1:A:935:MET:HG2	4:A:1480:HOH:O	2.20	0.41
1:A:972:ARG:HB3	4:A:1236:HOH:O	2.21	0.41
1:A:980:THR:HG23	4:A:1303:HOH:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/343 (100%)	320 (93%)	23 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	299/305 (98%)	290 (97%)	9 (3%)	36	13

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	816	PHE
1	A	820	PRO
1	A	829	ASP
1	A	830	GLN
1	A	879	TYR
1	A	893	VAL
1	A	928	LEU
1	A	975	ASP
1	A	1029	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	744	HIS
1	A	851	GLN
1	A	909	GLN
1	A	966	HIS
1	A	979	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MLI	A	1101	3	6,6,6	1.21	1 (16%)	7,7,7	1.28	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLI	A	1101	3	-	3/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	MLI	O7-C2	-2.22	1.23	1.30

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	MLI	O7-C2-C1	2.34	121.76	114.51

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1101	MLI	C3-C1-C2-O6
2	A	1101	MLI	C3-C1-C2-O7
2	A	1101	MLI	C2-C1-C3-O8

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	341/343 (99%)	2.76	162 (47%) 0 0	13, 33, 92, 145	4 (1%)

All (162) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	956	ILE	11.2
1	A	984	ILE	11.2
1	A	939	PHE	10.8
1	A	973	ILE	10.3
1	A	745	PHE	9.8
1	A	970	PHE	9.6
1	A	942	LEU	9.4
1	A	765	VAL	9.3
1	A	986	ILE	9.3
1	A	974	ALA	9.2
1	A	937	ILE	9.1
1	A	987	TYR	9.0
1	A	977	ILE	8.7
1	A	781	VAL	8.4
1	A	769	TYR	8.2
1	A	862	ALA	8.1
1	A	982	GLY	7.7
1	A	747	PHE	7.7
1	A	946	LEU	7.6
1	A	971	THR	7.6
1	A	866	VAL	7.5
1	A	770	ILE	7.3
1	A	780	ILE	7.3
1	A	976	VAL	7.3
1	A	744	HIS	7.2
1	A	985	SER	7.2
1	A	980	THR	7.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	966	HIS	7.0
1	A	951	GLY	6.9
1	A	957	LEU	6.9
1	A	873	LEU	6.8
1	A	868	LEU	6.7
1	A	928	LEU	6.7
1	A	722	VAL	6.6
1	A	978	PRO	6.5
1	A	750	TYR	6.5
1	A	941	GLY	6.4
1	A	954	ASN	6.3
1	A	768	GLN	6.2
1	A	981	LYS	6.2
1	A	763	LEU	6.0
1	A	746	THR	6.0
1	A	953	TYR	6.0
1	A	711	GLY	5.8
1	A	950	GLY	5.8
1	A	968	ASN	5.7
1	A	865	ALA	5.7
1	A	967	SER	5.7
1	A	964	TRP	5.7
1	A	863	PHE	5.6
1	A	764[A]	HIS	5.6
1	A	972	ARG	5.5
1	A	741	LEU	5.4
1	A	879	TYR	5.3
1	A	759	LEU	5.3
1	A	867	THR	5.3
1	A	874	VAL	5.3
1	A	959	PHE	5.3
1	A	988	GLY	5.2
1	A	721	VAL	5.2
1	A	893	VAL	5.2
1	A	779	CYS	5.1
1	A	963	GLY	5.0
1	A	983	ASN	4.9
1	A	940	LYS	4.8
1	A	989	ILE	4.8
1	A	965	THR	4.8
1	A	935	MET	4.7
1	A	748	PRO	4.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	938	ASN	4.7
1	A	943	GLN	4.6
1	A	784	VAL	4.6
1	A	949	CYS	4.6
1	A	894	LEU	4.6
1	A	871	HIS	4.5
1	A	857	PHE	4.5
1	A	724	GLY	4.5
1	A	856	ARG	4.5
1	A	945	HIS	4.5
1	A	742	SER	4.4
1	A	870	PRO	4.4
1	A	861	THR	4.4
1	A	936	GLN	4.3
1	A	955	GLN	4.3
1	A	749	VAL	4.3
1	A	975	ASP	4.3
1	A	771	HIS	4.2
1	A	961	PRO	4.2
1	A	709	GLY	4.1
1	A	979	GLN	4.1
1	A	826	LEU	4.1
1	A	927	SER	4.0
1	A	712	PHE	4.0
1	A	782	ASN	4.0
1	A	740	GLY	3.9
1	A	848	PHE	3.9
1	A	743	LYS	3.9
1	A	766	GLN	3.9
1	A	804	LEU	3.9
1	A	876[A]	CYS	3.8
1	A	727	ALA	3.8
1	A	864	GLU	3.7
1	A	932	LEU	3.7
1	A	872	ALA	3.7
1	A	717	PHE	3.7
1	A	828	ALA	3.7
1	A	700	THR	3.6
1	A	739	ALA	3.6
1	A	958	ALA	3.6
1	A	777	THR	3.6
1	A	726	THR	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	767	GLU	3.6
1	A	710	THR	3.5
1	A	869	ASN	3.5
1	A	860	ASN	3.5
1	A	944	SER	3.5
1	A	948	LYS	3.5
1	A	962	THR	3.5
1	A	960	ARG	3.4
1	A	952	LYS	3.4
1	A	969	LYS	3.4
1	A	725	CYS	3.3
1	A	925	CYS	3.3
1	A	772	PRO	3.2
1	A	875	VAL	3.2
1	A	719	TYR	3.2
1	A	773	LEU	3.2
1	A	775	LEU	3.1
1	A	708	PRO	3.0
1	A	934	MET	3.0
1	A	878	THR	3.0
1	A	703	PHE	3.0
1	A	738	TYR	3.0
1	A	880	SER	3.0
1	A	847	THR	2.9
1	A	720	GLY	2.9
1	A	723	GLU	2.8
1	A	783	GLY	2.8
1	A	947	LYS	2.8
1	A	926	SER	2.8
1	A	704	TYR	2.7
1	A	716	ALA	2.7
1	A	895	GLY	2.6
1	A	929	VAL	2.6
1	A	774	PRO	2.6
1	A	707	ILE	2.5
1	A	827	LEU	2.5
1	A	805	PRO	2.5
1	A	829	ASP	2.4
1	A	729	PHE	2.3
1	A	912	ASN	2.3
1	A	736	ASP	2.3
1	A	787	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	859	ILE	2.2
1	A	931	LEU	2.2
1	A	990	PRO	2.2
1	A	1025	SER	2.2
1	A	778	GLU	2.2
1	A	852	GLN	2.1
1	A	933	PRO	2.0
1	A	917	ASN	2.0
1	A	1040	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MLI	A	1101	7/7	0.91	0.09	20,26,29,30	0
3	NI	A	1102	1/1	0.97	0.05	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.