



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 08:27 PM UTC

PDB ID : 2Q60 / pdb_00002q60
Title : Crystal structure of the ligand binding domain of polyandrocarpa misakiensis rxr in tetramer in absence of ligand
Authors : Borel, F.; De Groot, A.; Juillan-Binard, C.; De Rosny, E.; Laudet, V.; Pebay-Peyroula, E.; Fontecilla-Camps, J.-C.; Ferrer, J.-L.
Deposited on : 2007-06-04
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	9% 59% 15% • 23%
1	B	258	6% 62% 14% • 23%
1	C	258	5% 62% 14% • 21%
1	D	258	14% 64% 10% • 24%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6256 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Retinoid X receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	198	Total	C	N	O	S	0	0	0
			1505	963	261	274	7			
1	B	198	Total	C	N	O	S	0	0	0
			1522	975	266	272	9			
1	C	203	Total	C	N	O	S	0	0	0
			1566	1001	276	281	8			
1	D	197	Total	C	N	O	S	0	0	0
			1488	941	265	274	8			

There are 100 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	80	MET	-	expression tag	UNP Q9UAF1
A	81	GLY	-	expression tag	UNP Q9UAF1
A	82	SER	-	expression tag	UNP Q9UAF1
A	83	SER	-	expression tag	UNP Q9UAF1
A	84	HIS	-	expression tag	UNP Q9UAF1
A	85	HIS	-	expression tag	UNP Q9UAF1
A	86	HIS	-	expression tag	UNP Q9UAF1
A	87	HIS	-	expression tag	UNP Q9UAF1
A	88	HIS	-	expression tag	UNP Q9UAF1
A	89	HIS	-	expression tag	UNP Q9UAF1
A	90	SER	-	expression tag	UNP Q9UAF1
A	91	SER	-	expression tag	UNP Q9UAF1
A	92	GLY	-	expression tag	UNP Q9UAF1
A	93	LEU	-	expression tag	UNP Q9UAF1
A	94	VAL	-	expression tag	UNP Q9UAF1
A	95	PRO	-	expression tag	UNP Q9UAF1
A	96	ARG	-	expression tag	UNP Q9UAF1
A	97	GLY	-	expression tag	UNP Q9UAF1
A	98	SER	-	expression tag	UNP Q9UAF1
A	99	HIS	-	expression tag	UNP Q9UAF1
A	100	MET	-	expression tag	UNP Q9UAF1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	101	ALA	-	expression tag	UNP Q9UAF1
A	102	SER	-	expression tag	UNP Q9UAF1
A	248	VAL	ILE	engineered mutation	UNP Q9UAF1
A	258	PRO	LEU	engineered mutation	UNP Q9UAF1
B	80	MET	-	expression tag	UNP Q9UAF1
B	81	GLY	-	expression tag	UNP Q9UAF1
B	82	SER	-	expression tag	UNP Q9UAF1
B	83	SER	-	expression tag	UNP Q9UAF1
B	84	HIS	-	expression tag	UNP Q9UAF1
B	85	HIS	-	expression tag	UNP Q9UAF1
B	86	HIS	-	expression tag	UNP Q9UAF1
B	87	HIS	-	expression tag	UNP Q9UAF1
B	88	HIS	-	expression tag	UNP Q9UAF1
B	89	HIS	-	expression tag	UNP Q9UAF1
B	90	SER	-	expression tag	UNP Q9UAF1
B	91	SER	-	expression tag	UNP Q9UAF1
B	92	GLY	-	expression tag	UNP Q9UAF1
B	93	LEU	-	expression tag	UNP Q9UAF1
B	94	VAL	-	expression tag	UNP Q9UAF1
B	95	PRO	-	expression tag	UNP Q9UAF1
B	96	ARG	-	expression tag	UNP Q9UAF1
B	97	GLY	-	expression tag	UNP Q9UAF1
B	98	SER	-	expression tag	UNP Q9UAF1
B	99	HIS	-	expression tag	UNP Q9UAF1
B	100	MET	-	expression tag	UNP Q9UAF1
B	101	ALA	-	expression tag	UNP Q9UAF1
B	102	SER	-	expression tag	UNP Q9UAF1
B	248	VAL	ILE	engineered mutation	UNP Q9UAF1
B	258	PRO	LEU	engineered mutation	UNP Q9UAF1
C	80	MET	-	expression tag	UNP Q9UAF1
C	81	GLY	-	expression tag	UNP Q9UAF1
C	82	SER	-	expression tag	UNP Q9UAF1
C	83	SER	-	expression tag	UNP Q9UAF1
C	84	HIS	-	expression tag	UNP Q9UAF1
C	85	HIS	-	expression tag	UNP Q9UAF1
C	86	HIS	-	expression tag	UNP Q9UAF1
C	87	HIS	-	expression tag	UNP Q9UAF1
C	88	HIS	-	expression tag	UNP Q9UAF1
C	89	HIS	-	expression tag	UNP Q9UAF1
C	90	SER	-	expression tag	UNP Q9UAF1
C	91	SER	-	expression tag	UNP Q9UAF1
C	92	GLY	-	expression tag	UNP Q9UAF1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	93	LEU	-	expression tag	UNP Q9UAF1
C	94	VAL	-	expression tag	UNP Q9UAF1
C	95	PRO	-	expression tag	UNP Q9UAF1
C	96	ARG	-	expression tag	UNP Q9UAF1
C	97	GLY	-	expression tag	UNP Q9UAF1
C	98	SER	-	expression tag	UNP Q9UAF1
C	99	HIS	-	expression tag	UNP Q9UAF1
C	100	MET	-	expression tag	UNP Q9UAF1
C	101	ALA	-	expression tag	UNP Q9UAF1
C	102	SER	-	expression tag	UNP Q9UAF1
C	248	VAL	ILE	engineered mutation	UNP Q9UAF1
C	258	PRO	LEU	engineered mutation	UNP Q9UAF1
D	80	MET	-	expression tag	UNP Q9UAF1
D	81	GLY	-	expression tag	UNP Q9UAF1
D	82	SER	-	expression tag	UNP Q9UAF1
D	83	SER	-	expression tag	UNP Q9UAF1
D	84	HIS	-	expression tag	UNP Q9UAF1
D	85	HIS	-	expression tag	UNP Q9UAF1
D	86	HIS	-	expression tag	UNP Q9UAF1
D	87	HIS	-	expression tag	UNP Q9UAF1
D	88	HIS	-	expression tag	UNP Q9UAF1
D	89	HIS	-	expression tag	UNP Q9UAF1
D	90	SER	-	expression tag	UNP Q9UAF1
D	91	SER	-	expression tag	UNP Q9UAF1
D	92	GLY	-	expression tag	UNP Q9UAF1
D	93	LEU	-	expression tag	UNP Q9UAF1
D	94	VAL	-	expression tag	UNP Q9UAF1
D	95	PRO	-	expression tag	UNP Q9UAF1
D	96	ARG	-	expression tag	UNP Q9UAF1
D	97	GLY	-	expression tag	UNP Q9UAF1
D	98	SER	-	expression tag	UNP Q9UAF1
D	99	HIS	-	expression tag	UNP Q9UAF1
D	100	MET	-	expression tag	UNP Q9UAF1
D	101	ALA	-	expression tag	UNP Q9UAF1
D	102	SER	-	expression tag	UNP Q9UAF1
D	248	VAL	ILE	engineered mutation	UNP Q9UAF1
D	258	PRO	LEU	engineered mutation	UNP Q9UAF1

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	49	Total O 49 49	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	42	Total 42	O 42	0	0
2	C	47	Total 47	O 47	0	0
2	D	37	Total 37	O 37	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Retinoid X receptor




• Molecule 1: Retinoid X receptor






• Molecule 1: Retinoid X receptor





● Molecule 1: Retinoid X receptor



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	82.26Å 96.12Å 151.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.22 – 2.90 48.22 – 2.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.22-2.90) 99.4 (48.22-2.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.34 (at 2.91Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.269 , 0.325 0.269 , 0.331	Depositor DCC
R_{free} test set	1361 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	58.6	Xtriage
Anisotropy	0.415	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 86.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	6256	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.89 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.7349e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/1537	0.78	2/2085 (0.1%)
1	B	0.44	0/1554	0.74	0/2103
1	C	0.45	0/1599	0.77	2/2163 (0.1%)
1	D	0.46	0/1519	0.77	1/2057 (0.0%)
All	All	0.46	0/6209	0.76	5/8408 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	D	0	1
All	All	0	2

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	108	MET	CA-C-N	5.33	126.50	119.84
1	A	108	MET	C-N-CA	5.33	126.50	119.84
1	C	320	THR	CA-C-N	5.22	139.52	127.00
1	C	320	THR	C-N-CA	5.22	139.52	127.00
1	D	149	ARG	N-CA-C	5.04	115.95	108.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	320	THR	Peptide
1	D	320	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1505	0	1471	22	0
1	B	1522	0	1497	16	0
1	C	1566	0	1539	19	0
1	D	1488	0	1431	12	0
2	A	49	0	0	1	0
2	B	42	0	0	0	0
2	C	47	0	0	0	0
2	D	37	0	0	0	0
All	All	6256	0	5938	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 66 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:190:HIS:CG	1:D:242:LEU:HD22	2.38	0.58
1:A:235:MET:HE1	1:A:293:LEU:CD2	2.35	0.56
1:D:235:MET:HE1	1:D:293:LEU:HD11	1.86	0.56
1:D:235:MET:HE1	1:D:293:LEU:CD1	2.35	0.56
1:C:164:PHE:HE1	1:C:175:LEU:HD11	1.70	0.55

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/258 (75%)	161 (83%)	21 (11%)	12 (6%)	1	3
1	B	194/258 (75%)	173 (89%)	14 (7%)	7 (4%)	2	11
1	C	199/258 (77%)	175 (88%)	16 (8%)	8 (4%)	2	9
1	D	193/258 (75%)	173 (90%)	14 (7%)	6 (3%)	3	14
All	All	780/1032 (76%)	682 (87%)	65 (8%)	33 (4%)	2	9

5 of 33 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	147	ALA
1	A	203	SER
1	A	319	ASP
1	A	321	PRO
1	B	120	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/231 (68%)	149 (94%)	9 (6%)	18	49
1	B	157/231 (68%)	145 (92%)	12 (8%)	12	36
1	C	163/231 (71%)	153 (94%)	10 (6%)	17	46
1	D	154/231 (67%)	141 (92%)	13 (8%)	10	31
All	All	632/924 (68%)	588 (93%)	44 (7%)	14	40

5 of 44 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	305	LEU
1	D	172	GLN
1	C	308	LEU
1	D	151	LEU
1	D	193	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	208	HIS
1	D	210	HIS
1	C	257	ASN
1	D	286	GLN
1	D	172	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	198/258 (76%)	0.85	24 (12%) 8 7	31, 59, 80, 85	1 (0%)
1	B	198/258 (76%)	0.94	15 (7%) 20 16	43, 66, 78, 81	0
1	C	203/258 (78%)	0.82	13 (6%) 25 20	41, 62, 73, 77	0
1	D	197/258 (76%)	1.22	37 (18%) 3 3	46, 67, 85, 94	0
All	All	796/1032 (77%)	0.96	89 (11%) 10 9	31, 64, 82, 94	1 (0%)

The worst 5 of 89 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	210	HIS	7.8
1	D	318	GLY	4.6
1	D	152	VAL	4.0
1	D	198	SER	3.7
1	A	152	VAL	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.