



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:30 AM UTC

PDB ID : 2Q61 / pdb_00002q61
Title : Crystal Structure of PPARgamma ligand binding domain bound to partial agonist SR145
Authors : Bruning, J.B.; Nettles, K.W.
Deposited on : 2007-06-04
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	3% 75% 15% • 7%
1	B	274	10% 75% 18% • 5%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4373 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Peroxisome Proliferator-Activated Receptor gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	256	Total	C	N	O	S	0	8	0
			2072	1343	335	383	11			
1	B	259	Total	C	N	O	S	0	8	0
			2091	1358	343	379	11			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	204	SER	-	expression tag	UNP P37231
B	204	SER	-	expression tag	UNP P37231

- Molecule 2 is 1-BENZYL-5-CHLORO-3-(PHENYLTHIO)-1H-INDOLE-2-CARBOXYLIC ACID (CCD ID: SF1) (formula: C₂₂H₁₆ClNO₂S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	S	0	0
			27	22	1	1	2	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	B	1	Total	C	Cl	N	O	S	0	0
			27	22	1	1	2	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	79	Total	O	0	0
			79	79		
3	B	77	Total	O	0	0
			77	77		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peroxisome Proliferator-Activated Receptor gamma



- Molecule 1: Peroxisome Proliferator-Activated Receptor gamma



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	90.13Å 62.39Å 117.93Å 90.00° 100.98° 90.00°	Depositor
Resolution (Å)	9.97 – 2.20 9.97 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.4 (9.97-2.20) 98.2 (9.97-2.20)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.04 (at 2.19Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.188 , 0.229 0.186 , 0.223	Depositor DCC
R_{free} test set	1650 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	31.4	Xtriage
Anisotropy	0.080	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 62.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4373	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	2/2130 (0.1%)	0.92	0/2872
1	B	0.86	4/2148 (0.2%)	0.92	0/2893
All	All	0.80	6/4278 (0.1%)	0.92	0/5765

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	461	THR	C-N	-18.34	1.10	1.33
1	B	460	GLU	C-N	-12.90	1.15	1.33
1	A	462	ASP	CG-OD2	9.01	1.42	1.25
1	A	462	ASP	CG-OD1	8.43	1.41	1.25
1	B	244	LYS	N-CA	8.27	1.61	1.46
1	B	244	LYS	CA-CB	7.65	1.68	1.53

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2072	0	2135	47	0
1	B	2091	0	2149	40	0
2	A	27	0	15	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	27	0	15	6	0
3	A	79	0	0	0	0
3	B	77	0	0	6	0
All	All	4373	0	4314	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:357:ARG:HD2	1:A:358:LYS:O	1.60	1.01
1:A:357:ARG:HG3	1:A:357:ARG:HH11	1.24	1.00
1:A:437:GLN:HG2	1:B:439[A]:MET:HE1	1.54	0.90
1:B:449:HIS:HD2	3:B:5058:HOH:O	1.60	0.83
1:A:357:ARG:CD	1:A:358:LYS:O	2.26	0.83
1:A:444:GLN:HG3	1:B:443[A]:ARG:NH2	1.99	0.78
1:A:276:GLU:OE2	1:A:357:ARG:NH1	2.23	0.71
1:B:334:MET:HE3	1:B:371:ALA:HB2	1.75	0.68
1:A:357:ARG:HG3	1:A:357:ARG:NH1	2.05	0.66
1:B:331:ALA:HA	1:B:334:MET:HE3	1.78	0.66
1:A:292:ALA:O	1:A:296[A]:ILE:HG12	1.96	0.66
1:B:334:MET:HE1	3:B:5019:HOH:O	1.95	0.65
1:B:362:ASP:HB2	3:B:5046:HOH:O	1.95	0.65
1:B:363:PHE:HB2	1:B:452:LEU:HD21	1.77	0.64
1:A:226:PHE:HE1	1:A:296[A]:ILE:HD13	1.62	0.64
1:B:279:ILE:HG23	1:B:463:MET:HE2	1.79	0.64
1:B:466:HIS:N	3:B:5054:HOH:O	2.30	0.64
1:A:396:ASP:OD2	1:B:438:LYS:HE2	1.99	0.62
1:A:288[B]:ARG:HE	2:A:7001:SF1:CAS	2.14	0.59
1:A:370:PHE:CZ	1:A:442:LEU:HD13	2.36	0.59
1:A:437:GLN:HG2	1:B:439[A]:MET:CE	2.31	0.59
1:B:330:LEU:O	1:B:334:MET:HG3	2.03	0.59
1:A:454:GLN:O	1:A:458:LYS:HG2	2.04	0.56
2:A:7001:SF1:OAA	2:A:7001:SF1:HAQ1	2.06	0.56
1:B:449:HIS:CD2	3:B:5058:HOH:O	2.46	0.56
1:A:363:PHE:HB2	1:A:452:LEU:CD2	2.38	0.54
1:A:228:LEU:HB2	1:A:343:GLU:HG2	1.90	0.53
1:A:357:ARG:HH11	1:A:357:ARG:CG	2.07	0.53
1:B:334:MET:CE	1:B:371:ALA:HB2	2.38	0.53
1:A:255:LEU:HD22	1:A:277:VAL:CG2	2.38	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:288[B]:ARG:HE	2:B:5001:SF1:CAS	2.22	0.52
1:B:328:THR:OG1	1:B:442:LEU:HD11	2.10	0.51
1:A:255:LEU:HD22	1:A:277:VAL:HG21	1.93	0.51
1:B:462:ASP:O	1:B:463:MET:HG2	2.11	0.51
1:B:334:MET:CE	3:B:5019:HOH:O	2.58	0.50
1:A:252:MET:SD	1:A:277:VAL:HG11	2.51	0.50
1:A:282:PHE:CD2	1:A:463:MET:HE1	2.46	0.50
1:A:226:PHE:CE1	1:A:296[A]:ILE:HD13	2.45	0.50
1:A:439:MET:HE1	1:B:437:GLN:HB2	1.92	0.50
1:A:325:ILE:HD11	1:A:392:ILE:HG13	1.93	0.50
1:B:276:GLU:HG2	1:B:279:ILE:HG12	1.93	0.49
1:B:348:MET:CE	2:B:5001:SF1:HAH	2.42	0.49
1:B:323:HIS:CE1	1:B:476:LEU:HD12	2.48	0.49
1:A:363:PHE:CB	1:A:452:LEU:HD21	2.42	0.49
1:A:282:PHE:HD2	1:A:463:MET:HE1	1.77	0.49
1:A:439:MET:HE1	1:B:437:GLN:CB	2.43	0.49
1:A:444:GLN:HG3	1:B:443[A]:ARG:CZ	2.41	0.49
1:A:360:PHE:O	1:A:363:PHE:HD2	1.97	0.48
2:B:5001:SF1:HAQ1	2:B:5001:SF1:OAB	2.14	0.47
1:A:357:ARG:HD2	1:A:358:LYS:C	2.35	0.47
1:B:325:ILE:HD11	1:B:392:ILE:HG13	1.97	0.47
1:A:288[A]:ARG:HH22	1:A:295:GLU:CD	2.23	0.46
1:A:276:GLU:CD	1:A:357:ARG:HH12	2.24	0.46
1:B:266:HIS:CD2	1:B:280:ARG:HG2	2.52	0.45
1:B:411:ASP:OD2	1:B:415:GLN:NE2	2.46	0.45
1:A:370:PHE:CB	1:A:445:ILE:HD11	2.48	0.44
1:A:446:VAL:O	1:A:450:VAL:HG13	2.17	0.44
1:A:404:LYS:HE2	1:A:408:ASP:OD1	2.17	0.44
1:A:439:MET:CE	1:B:437:GLN:HB2	2.48	0.44
1:A:348:MET:HE3	1:A:353:LEU:HD21	1.99	0.44
1:B:325:ILE:HD11	1:B:392:ILE:CG1	2.48	0.44
1:A:433:ALA:O	1:A:437:GLN:HG3	2.18	0.43
2:B:5001:SF1:OAA	2:B:5001:SF1:SAR	2.76	0.43
1:A:318:LEU:O	1:A:322:VAL:HB	2.18	0.43
1:A:363:PHE:CB	1:A:452:LEU:CD2	2.96	0.43
1:A:379:LEU:HD11	1:A:435:LEU:HD13	2.01	0.43
1:A:363:PHE:HB3	1:A:452:LEU:HD21	1.99	0.43
1:B:226:PHE:CG	1:B:329:MET:HE1	2.54	0.42
1:B:334:MET:HE3	1:B:371:ALA:CB	2.44	0.42
1:B:348:MET:HE3	2:B:5001:SF1:HAH	2.00	0.42
1:A:437:GLN:O	1:A:440:THR:HB	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:249:ILE:HD12	1:B:341:ILE:HD11	2.02	0.42
1:B:252:MET:SD	1:B:277:VAL:HG11	2.59	0.41
1:B:436:LEU:O	1:B:439[A]:MET:HE2	2.20	0.41
1:B:255:LEU:HD21	1:B:277:VAL:HG13	2.03	0.41
2:B:5001:SF1:CAZ	2:B:5001:SF1:HAJ	2.50	0.41
1:A:474:LYS:HA	1:A:475:ASP:HA	1.81	0.41
1:B:331:ALA:HA	1:B:334:MET:CE	2.47	0.40
1:B:447:THR:O	1:B:450:VAL:HG22	2.21	0.40
1:A:354:LYS:HA	1:A:361:GLY:O	2.21	0.40
1:A:226:PHE:CG	1:A:329:MET:HE1	2.56	0.40
1:A:439:MET:CE	1:B:437:GLN:CB	2.99	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/274 (95%)	255 (98%)	5 (2%)	0	100	100
1	B	259/274 (94%)	251 (97%)	7 (3%)	1 (0%)	30	34
All	All	519/548 (95%)	506 (98%)	12 (2%)	1 (0%)	43	51

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	264	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/247 (94%)	219 (95%)	12 (5%)	21	26
1	B	230/247 (93%)	212 (92%)	18 (8%)	11	13
All	All	461/494 (93%)	431 (94%)	30 (6%)	16	18

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	277	VAL
1	A	322	VAL
1	A	342[A]	SER
1	A	342[B]	SER
1	A	343	GLU
1	A	357	ARG
1	A	441	ASP
1	A	442	LEU
1	A	450	VAL
1	A	461	THR
1	A	463	MET
1	A	474	LYS
1	B	216	LYS
1	B	234	ARG
1	B	253	ASN
1	B	275	LYS
1	B	276	GLU
1	B	323	HIS
1	B	342	SER
1	B	345	GLN
1	B	357	ARG
1	B	362	ASP
1	B	363	PHE
1	B	369	GLU
1	B	428	SER
1	B	443[A]	ARG
1	B	443[B]	ARG
1	B	452	LEU
1	B	468	LEU
1	B	474	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	283	GLN
1	A	430	GLN
1	A	449	HIS
1	A	451	GLN
1	B	253	ASN
1	B	283	GLN
1	B	294	GLN
1	B	312	ASN
1	B	323	HIS
1	B	412	ASN
1	B	449	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SF1	A	7001	-	30,30,30	1.58	5 (16%)	40,42,42	1.17	3 (7%)
2	SF1	B	5001	-	30,30,30	1.53	5 (16%)	40,42,42	1.22	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SF1	A	7001	-	-	0/12/12/12	0/4/4/4
2	SF1	B	5001	-	-	2/12/12/12	0/4/4/4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	7001	SF1	CAX-NBA	-5.41	1.32	1.39
2	B	5001	SF1	CAX-NBA	-5.12	1.32	1.39
2	B	5001	SF1	CAQ-CAU	3.13	1.57	1.51
2	A	7001	SF1	CAQ-CAU	3.01	1.56	1.51
2	B	5001	SF1	CAY-CAW	-2.93	1.39	1.44
2	A	7001	SF1	CAY-CAW	-2.78	1.39	1.44
2	A	7001	SF1	CAZ-NBA	-2.67	1.34	1.39
2	B	5001	SF1	OAB-CAS	-2.31	1.24	1.30
2	B	5001	SF1	CAZ-NBA	-2.28	1.35	1.39
2	A	7001	SF1	OAB-CAS	-2.28	1.24	1.30

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	5001	SF1	CAS-CAX-NBA	3.80	127.32	122.59
2	A	7001	SF1	CAU-CAQ-NBA	-3.68	107.48	113.46
2	B	5001	SF1	CAU-CAQ-NBA	-3.43	107.88	113.46
2	A	7001	SF1	CAV-SAR-CAW	2.76	107.51	103.43
2	B	5001	SF1	CAO-CAZ-CAY	-2.38	119.30	121.95
2	A	7001	SF1	CAS-CAX-NBA	2.31	125.47	122.59

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	5001	SF1	OAB-CAS-CAX-NBA
2	B	5001	SF1	OAA-CAS-CAX-NBA


There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	7001	SF1	2	0
2	B	5001	SF1	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	460:GLU	C	461:THR	N	1.15
1	B	461:THR	C	462:ASP	N	1.10

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	256/274 (93%)	0.04	9 (3%) 47 44	9, 20, 42, 54	14 (5%)
1	B	259/274 (94%)	0.47	28 (10%) 11 8	9, 22, 44, 59	8 (3%)
All	All	515/548 (93%)	0.26	37 (7%) 21 18	9, 21, 43, 59	22 (4%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	475	ASP	4.5
1	A	239	GLY	3.6
1	B	241	THR	3.5
1	B	267	ILE	3.3
1	B	454	GLN	3.3
1	B	305	GLY	3.2
1	A	260	ASP	3.1
1	B	262	ILE	3.1
1	B	284	GLY	3.0
1	B	282	PHE	3.0
1	B	247	PHE	2.9
1	A	287	PHE	2.9
1	B	363	PHE	2.9
1	B	263	LYS	2.8
1	B	306	PHE	2.8
1	A	209	ALA	2.8
1	B	352	PHE	2.7
1	B	475	ASP	2.7
1	A	231	ALA	2.6
1	A	299	TYR	2.5
1	B	276	GLU	2.5
1	B	351	GLU	2.5
1	A	472	ILE	2.4
1	B	256	MET	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	249	ILE	2.3
1	B	358	LYS	2.3
1	B	303	ILE	2.3
1	B	253	ASN	2.3
1	B	245	SER	2.2
1	B	362	ASP	2.2
1	B	254	SER	2.2
1	A	242	THR	2.1
1	B	240	LYS	2.1
1	B	360	PHE	2.1
1	B	451	GLN	2.1
1	B	304	PRO	2.0
1	B	261	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SF1	B	5001	27/27	0.88	0.09	31,34,41,42	0
2	SF1	A	7001	27/27	0.93	0.06	29,35,39,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around SF1 B 5001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.