



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 06:35 PM UTC

PDB ID : 4Q6R / pdb_00004q6r
Title : Crystal structure of human sphingosine-1-phosphate lyase in complex with inhibitor 6-[(2R)-4-(4-benzyl-7-chlorophthalazin-1-yl)-2-methylpiperazin-1-yl]pyridine-3-carbonitrile
Authors : Srinivas, H.
Deposited on : 2014-04-23
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	511	
1	B	511	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SIN	A	901	-	X	-	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7336 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sphingosine-1-phosphate lyase 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	443	Total	C	N	O	P	S	0	0	0
			3449	2214	581	631	1	22			
1	B	444	Total	C	N	O	P	S	0	0	0
			3458	2220	583	632	1	22			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	58	GLY	-	expression tag	UNP O95470
A	59	PRO	-	expression tag	UNP O95470
A	60	GLY	-	expression tag	UNP O95470
A	61	SER	-	expression tag	UNP O95470
B	58	GLY	-	expression tag	UNP O95470
B	59	PRO	-	expression tag	UNP O95470
B	60	GLY	-	expression tag	UNP O95470
B	61	SER	-	expression tag	UNP O95470

- Molecule 2 is SUCCINIC ACID (CCD ID: SIN) (formula: C₄H₆O₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			8	4	4		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 6-[(2R)-4-(4-benzyl-7-chlorophthalazin-1-yl)-2-methylpiperazin-1-yl]pyridine-3-carbonitrile (CCD ID: 30J) (formula: $C_{26}H_{23}ClN_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total 33	C 26	Cl 1	N 6	0	0
4	B	1	Total 33	C 26	Cl 1	N 6	0	0

- Molecule 5 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	186	Total 186	O 186	0	0
6	B	158	Total 158	O 158	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Sphingosine-1-phosphate lyase 1



• Molecule 1: Sphingosine-1-phosphate lyase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	127.98Å 130.53Å 68.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.66 – 2.40 19.66 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.66-2.40) 99.7 (19.66-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.48 (at 2.41Å)	Xtriage
Refinement program	BUSTER 2.9.3	Depositor
R, R_{free}	0.178 , 0.206 0.182 , 0.208	Depositor DCC
R_{free} test set	2272 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	35.5	Xtriage
Anisotropy	1.064	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.010 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7336	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, 30J, SIN, LLP, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	4/3505 (0.1%)	1.32	5/4740 (0.1%)
1	B	0.84	5/3514 (0.1%)	1.32	7/4751 (0.1%)
All	All	0.84	9/7019 (0.1%)	1.32	12/9491 (0.1%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	265	MET	SD-CE	-9.11	1.56	1.79
1	A	265	MET	SD-CE	-7.61	1.60	1.79
1	A	254	MET	SD-CE	-6.91	1.62	1.79
1	B	298	VAL	CA-CB	6.67	1.57	1.54
1	B	254	MET	SD-CE	-6.65	1.62	1.79
1	B	216	MET	SD-CE	-5.94	1.64	1.79
1	B	409	MET	SD-CE	-5.94	1.64	1.79
1	A	216	MET	SD-CE	-5.68	1.65	1.79
1	A	298	VAL	CA-CB	5.24	1.56	1.54

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	394	GLY	N-CA-C	-9.73	101.36	110.21
1	B	394	GLY	N-CA-C	-8.73	102.27	110.21
1	B	232	THR	CB-CA-C	6.77	121.67	110.97
1	B	297	PRO	CA-C-N	5.74	125.94	120.43
1	B	297	PRO	C-N-CA	5.74	125.94	120.43
1	A	319	GLY	N-CA-C	5.31	120.48	113.37
1	A	297	PRO	CA-C-N	5.26	125.48	120.43
1	A	297	PRO	C-N-CA	5.26	125.48	120.43
1	B	153	GLU	CA-C-N	5.17	127.97	120.79
1	B	153	GLU	C-N-CA	5.17	127.97	120.79

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	252	PHE	N-CA-C	5.10	119.48	113.16
1	B	319	GLY	N-CA-C	5.01	120.09	113.37

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3449	0	3459	39	0
1	B	3458	0	3472	32	0
2	A	8	0	5	0	0
3	A	6	0	8	0	0
4	A	33	0	23	4	0
4	B	33	0	23	6	0
5	B	5	0	0	0	0
6	A	186	0	0	2	0
6	B	158	0	0	2	0
All	All	7336	0	6990	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:903:30J:H17	4:A:903:30J:H1	1.28	1.15
1:A:361:SER:HB3	6:A:1182:HOH:O	1.56	1.05
4:A:903:30J:H17	4:A:903:30J:C3	2.00	0.91
1:B:218:CYS:HB3	1:B:254:MET:HE1	1.58	0.83
1:A:218:CYS:HB3	1:A:254:MET:HE1	1.59	0.82
1:A:545:PHE:CD2	4:B:602:30J:H23	2.25	0.71
4:A:903:30J:H1	4:A:903:30J:C30	2.17	0.70
1:A:289:GLN:HE22	1:A:292:HIS:HD2	1.39	0.70
1:B:289:GLN:HE22	1:B:292:HIS:HD2	1.41	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:234:GLU:HG2	1:A:255:LYS:HB3	1.77	0.66
1:B:488:THR:H	1:B:491:HIS:HD2	1.45	0.65
4:A:903:30J:C3	4:A:903:30J:C30	2.74	0.65
1:B:234:GLU:HG2	1:B:255:LYS:HB3	1.79	0.64
1:A:216:MET:HE2	1:A:391:THR:HA	1.83	0.60
1:B:216:MET:HE2	1:B:391:THR:HA	1.83	0.60
1:A:179:PRO:HD2	1:B:137:ASP:HA	1.84	0.59
1:A:210:GLY:HA3	1:A:353:LLP:H5'2	1.85	0.58
1:B:210:GLY:HA3	1:B:353:LLP:H5'2	1.84	0.58
1:B:212:GLU:HG3	1:B:392:ILE:HD12	1.86	0.58
1:A:488:THR:H	1:A:491:HIS:HD2	1.52	0.57
1:A:252:PHE:HB2	1:A:254:MET:HE2	1.87	0.57
1:A:212:GLU:HG3	1:A:392:ILE:HD12	1.87	0.57
1:A:490:LEU:HD23	1:A:493:ARG:NH2	2.21	0.56
1:A:488:THR:H	1:A:491:HIS:CD2	2.24	0.56
1:B:252:PHE:HB2	1:B:254:MET:HE2	1.87	0.55
1:A:358:PRO:HG3	1:A:403:ALA:HB3	1.89	0.55
1:A:241:ALA:O	1:A:258:ARG:NH2	2.38	0.55
1:A:357:ALA:HB3	6:A:1182:HOH:O	2.07	0.55
1:B:358:PRO:HG3	1:B:403:ALA:HB3	1.89	0.54
1:B:241:ALA:O	1:B:258:ARG:NH2	2.38	0.53
1:A:545:PHE:CG	4:B:602:30J:H23	2.44	0.51
1:A:484:HIS:C	1:A:484:HIS:CD2	2.89	0.51
1:A:289:GLN:HE22	1:A:292:HIS:CD2	2.25	0.50
1:B:133:TYR:O	1:B:136:MET:HG2	2.11	0.50
1:B:484:HIS:C	1:B:484:HIS:CD2	2.90	0.50
1:B:289:GLN:HE22	1:B:292:HIS:CD2	2.26	0.49
1:A:182:ARG:NH1	1:B:551:SER:O	2.45	0.49
1:B:488:THR:H	1:B:491:HIS:CD2	2.28	0.48
1:A:188:ILE:HG22	1:A:363:LEU:HD21	1.95	0.48
4:B:602:30J:H17	4:B:602:30J:C3	2.43	0.47
1:B:352:HIS:HD2	1:B:359:LYS:NZ	2.13	0.47
1:A:352:HIS:HD2	1:A:359:LYS:NZ	2.13	0.47
1:A:474:LEU:HD22	1:A:485:PHE:HD1	1.80	0.47
1:B:140:TRP:CE3	1:B:149:VAL:HG11	2.50	0.47
1:A:551:SER:O	1:B:182:ARG:NH1	2.48	0.46
1:A:133:TYR:O	1:A:136:MET:HG2	2.15	0.46
1:B:474:LEU:HD22	1:B:485:PHE:HD1	1.81	0.46
1:A:150:TYR:HA	1:A:488:THR:HB	1.97	0.46
1:B:188:ILE:HG22	1:B:363:LEU:HD21	1.97	0.46
1:B:434:LEU:HD13	1:B:453:LEU:HD21	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:233:PRO:HB3	1:A:281:ALA:HB2	1.98	0.45
4:B:602:30J:H17	4:B:602:30J:H1	1.97	0.45
1:B:498:ILE:HG12	6:B:718:HOH:O	2.16	0.44
1:B:233:PRO:HB3	1:B:281:ALA:HB2	1.99	0.44
1:B:210:GLY:HA3	1:B:353:LLP:C5'	2.48	0.44
1:B:239:GLN:HG3	6:B:826:HOH:O	2.18	0.44
1:A:116:LEU:HD21	1:B:416:TYR:HD2	1.83	0.44
1:A:522:MET:HG3	1:A:526:TYR:CE2	2.54	0.43
1:B:267:VAL:HG12	1:B:297:PRO:HD2	2.01	0.43
1:B:285:CYS:HB2	1:B:296:ASP:CB	2.48	0.43
1:A:240:SER:HB3	1:A:261:LEU:HD11	2.01	0.43
1:A:285:CYS:HB2	1:A:296:ASP:CB	2.49	0.42
1:B:474:LEU:HD22	1:B:485:PHE:CD1	2.54	0.42
1:A:273:ARG:HA	1:A:276:ILE:HD12	2.02	0.42
1:A:474:LEU:HD22	1:A:485:PHE:CD1	2.54	0.42
1:A:259:VAL:HG23	1:A:267:VAL:HG23	2.00	0.42
1:A:545:PHE:CD2	4:B:602:30J:C39	3.01	0.42
4:B:602:30J:H4	4:B:602:30J:H19	1.71	0.42
1:A:462:ARG:CZ	1:A:511:ILE:HG23	2.50	0.42
1:A:210:GLY:HA3	1:A:353:LLP:C5'	2.48	0.41
1:B:522:MET:HG3	1:B:526:TYR:CE2	2.54	0.41
1:A:434:LEU:HD13	1:A:453:LEU:HD21	2.02	0.41
1:A:263:LYS:O	1:A:265:MET:HE2	2.21	0.41
1:B:150:TYR:HA	1:B:488:THR:HB	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	440/511 (86%)	429 (98%)	10 (2%)	1 (0%)	43 58

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	441/511 (86%)	429 (97%)	11 (2%)	1 (0%)	43 58
All	All	881/1022 (86%)	858 (97%)	21 (2%)	2 (0%)	43 58

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	139	PHE
1	B	320	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	366/429 (85%)	358 (98%)	8 (2%)	45 67
1	B	367/429 (86%)	358 (98%)	9 (2%)	42 64
All	All	733/858 (85%)	716 (98%)	17 (2%)	44 66

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	128	GLU
1	A	182	ARG
1	A	239	GLN
1	A	289	GLN
1	A	324	VAL
1	A	442	VAL
1	A	484	HIS
1	A	490	LEU
1	B	128	GLU
1	B	182	ARG
1	B	239	GLN
1	B	259	VAL
1	B	289	GLN
1	B	324	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	442	VAL
1	B	484	HIS
1	B	528	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	197	ASN
1	A	292	HIS
1	A	352	HIS
1	A	383	GLN
1	A	422	GLN
1	A	436	ASN
1	A	478	GLN
1	A	491	HIS
1	A	499	GLN
1	A	536	ASN
1	B	292	HIS
1	B	352	HIS
1	B	422	GLN
1	B	436	ASN
1	B	491	HIS
1	B	536	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	A	353	1	23,24,25	2.04	5 (21%)	25,32,34	1.44	4 (16%)
1	LLP	B	353	1	23,24,25	2.23	6 (26%)	25,32,34	1.57	6 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	353	1	-	4/16/17/19	0/1/1/1
1	LLP	B	353	1	-	4/16/17/19	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	353	LLP	C4'-NZ	7.33	1.51	1.27
1	A	353	LLP	C4'-NZ	6.83	1.50	1.27
1	B	353	LLP	C4-C4'	4.53	1.56	1.46
1	A	353	LLP	C4-C4'	3.72	1.54	1.46
1	B	353	LLP	C3-C2	3.43	1.44	1.41
1	B	353	LLP	C6-C5	3.09	1.43	1.37
1	A	353	LLP	C6-C5	2.98	1.43	1.37
1	A	353	LLP	C4-C5	2.69	1.45	1.42
1	B	353	LLP	C4-C3	2.51	1.45	1.41
1	B	353	LLP	C4-C5	2.31	1.45	1.42
1	A	353	LLP	C4-C3	2.12	1.44	1.41

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	353	LLP	CE-NZ-C4'	-3.66	107.01	118.72
1	B	353	LLP	CE-NZ-C4'	-3.65	107.02	118.72
1	B	353	LLP	OP4-P-OP1	-3.06	98.16	106.44
1	A	353	LLP	OP4-P-OP1	-2.42	99.90	106.44
1	A	353	LLP	OP3-P-OP4	2.31	112.69	106.67
1	A	353	LLP	C3-C4-C5	-2.19	116.52	118.28
1	B	353	LLP	C6-N1-C2	2.16	123.11	119.20
1	B	353	LLP	OP3-P-OP4	2.13	112.23	106.67
1	B	353	LLP	C3-C4-C5	-2.13	116.57	118.28
1	B	353	LLP	OP2-P-OP4	2.07	112.06	106.67

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	353	LLP	C4-C4'-NZ-CE
1	A	353	LLP	C6-C5-C5'-OP4
1	A	353	LLP	O-C-CA-CB
1	B	353	LLP	C4-C4'-NZ-CE
1	B	353	LLP	O-C-CA-CB
1	B	353	LLP	C6-C5-C5'-OP4
1	B	353	LLP	C3-C4-C4'-NZ
1	A	353	LLP	C3-C4-C4'-NZ

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	353	LLP	2	0
1	B	353	LLP	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	30J	A	903	-	35,37,37	1.43	6 (17%)	45,52,52	2.01	11 (24%)
4	30J	B	602	-	35,37,37	1.61	6 (17%)	45,52,52	1.81	12 (26%)
2	SIN	A	901	-	7,7,7	2.69	3 (42%)	8,8,8	2.02	3 (37%)
3	GOL	A	902	-	5,5,5	0.54	0	5,5,5	0.48	0
5	PO4	B	601	-	4,4,4	2.30	1 (25%)	6,6,6	0.93	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	30J	B	602	-	-	2/14/27/27	0/5/5/5
4	30J	A	903	-	-	2/14/27/27	0/5/5/5
2	SIN	A	901	-	-	4/5/5/5	-
3	GOL	A	902	-	-	2/4/4/4	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	602	30J	C9-C10	-5.34	1.37	1.43
4	A	903	30J	C9-C10	-5.22	1.37	1.43
2	A	901	SIN	O3-C4	4.34	1.36	1.22
2	A	901	SIN	C3-C4	4.25	1.60	1.50
5	B	601	PO4	P-O1	3.57	1.58	1.50
4	B	602	30J	C10-C5	-3.18	1.37	1.43
4	B	602	30J	C6-C5	-2.97	1.38	1.43
4	A	903	30J	C33-N35	-2.87	1.43	1.47
4	A	903	30J	C6-C5	-2.60	1.38	1.43
2	A	901	SIN	O2-C1	2.32	1.38	1.30
4	A	903	30J	C10-C5	-2.29	1.39	1.43
4	B	602	30J	C39-N29	-2.28	1.42	1.46
4	B	602	30J	C3-C5	-2.14	1.37	1.42
4	A	903	30J	C3-C5	-2.13	1.37	1.42
4	A	903	30J	C11-C10	-2.04	1.38	1.42
4	B	602	30J	C9-N8	-2.01	1.29	1.33

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	602	30J	C5-C6-N7	-5.29	118.63	126.80
4	A	903	30J	C33-C30-N29	4.93	118.16	110.86
4	A	903	30J	C5-C6-N7	-4.89	119.25	126.80
4	A	903	30J	C11-C10-C9	-4.42	119.09	123.22
4	A	903	30J	C39-N29-C30	4.33	124.79	112.53
4	B	602	30J	C36-N35-C33	4.26	121.99	114.63
4	B	602	30J	C9-N8-N7	3.43	122.82	119.97
4	A	903	30J	C9-N8-N7	3.40	122.80	119.97
4	A	903	30J	C36-N35-C33	3.24	120.24	114.63
4	B	602	30J	C39-N29-C30	3.22	121.65	112.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	602	30J	C10-C9-N8	-3.21	119.05	122.31
2	A	901	SIN	C3-C2-C1	3.13	121.97	113.67
4	A	903	30J	C10-C9-N8	-3.11	119.14	122.31
2	A	901	SIN	O4-C4-O3	-3.09	115.39	123.33
2	A	901	SIN	O4-C4-C3	2.94	123.30	114.00
4	A	903	30J	C18-C15-C9	-2.94	106.00	113.00
4	A	903	30J	C11-C10-C5	2.74	121.82	118.42
4	B	602	30J	C44-N43-C42	2.72	121.86	117.41
4	A	903	30J	C44-N43-C42	2.63	121.72	117.41
4	B	602	30J	C46-C44-N43	-2.50	119.93	123.50
4	B	602	30J	C11-C10-C9	-2.47	120.92	123.22
4	A	903	30J	C46-C44-N43	-2.44	120.01	123.50
4	B	602	30J	C6-C5-C10	2.20	119.10	116.37
4	B	602	30J	C11-C13-C2	-2.19	116.88	119.22
4	B	602	30J	C18-C15-C9	-2.10	108.00	113.00
4	B	602	30J	C5-C6-N29	2.07	124.73	121.18

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	902	GOL	O1-C1-C2-O2
3	A	902	GOL	O1-C1-C2-C3
4	B	602	30J	N7-C6-N29-C39
4	A	903	30J	N7-C6-N29-C39
2	A	901	SIN	C2-C3-C4-O3
2	A	901	SIN	C2-C3-C4-O4
4	A	903	30J	C5-C6-N29-C39
4	B	602	30J	C5-C6-N29-C39
2	A	901	SIN	O2-C1-C2-C3
2	A	901	SIN	O1-C1-C2-C3


There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	903	30J	4	0
4	B	602	30J	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	442/511 (86%)	-0.38	2 (0%) 87 85	31, 43, 64, 92	0
1	B	443/511 (86%)	-0.10	7 (1%) 70 66	30, 44, 90, 114	0
All	All	885/1022 (86%)	-0.24	9 (1%) 79 76	30, 43, 82, 114	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	139	PHE	2.3
1	A	537	MET	2.3
1	B	528	MET	2.3
1	B	537	MET	2.3
1	B	137	ASP	2.2
1	B	531	THR	2.2
1	A	139	PHE	2.1
1	B	533	VAL	2.0
1	B	494	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	B	353	24/25	0.92	0.11	47,53,55,57	0
1	LLP	A	353	24/25	0.97	0.06	31,35,37,37	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	902	6/6	0.79	0.17	67,69,69,70	0
2	SIN	A	901	8/8	0.82	0.15	47,52,59,61	0
5	PO4	B	601	5/5	0.83	0.15	98,104,105,105	0
4	30J	A	903	33/33	0.89	0.10	58,64,70,72	0
4	30J	B	602	33/33	0.93	0.08	36,41,48,51	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 30J A 903:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 30J B 602:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.