



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2026 – 05:29 PM UTC

PDB ID : 7Q63 / pdb_00007q63
Title : The tandem SH2 domains of SYK
Authors : Bradshaw, W.J.; Katis, V.L.; Chen, Z.; Bountra, C.; von Delft, F.; Gileadi, O.; Brennan, P.E.
Deposited on : 2021-11-05
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	265	 12% 87% 10% ..
1	BBB	265	 6% 89% 8% ..
1	CCC	265	 17% 77% 8% 15%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	BBB	305	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6380 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase SYK.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	259	Total	C	N	O	S	0	7	0
			2102	1329	378	388	7			
1	BBB	259	Total	C	N	O	S	0	4	0
			2081	1318	373	383	7			
1	CCC	225	Total	C	N	O	S	0	6	0
			1834	1166	330	330	8			

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
AAA	5	SER	-	expression tag	UNP P43405
BBB	5	SER	-	expression tag	UNP P43405
CCC	5	SER	-	expression tag	UNP P43405

- Molecule 2 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	AAA	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is THIOCYANATE ION (CCD ID: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	AAA	1	Total	C	N	S	0	0
			3	1	1	1		
3	AAA	1	Total	C	N	S	0	0
			3	1	1	1		
3	AAA	1	Total	C	N	S	0	0
			3	1	1	1		
3	BBB	1	Total	C	N	S	0	0
			3	1	1	1		
3	BBB	1	Total	C	N	S	0	0
			3	1	1	1		
3	BBB	1	Total	C	N	S	0	0
			3	1	1	1		
3	BBB	1	Total	C	N	S	0	0
			3	1	1	1		
3	CCC	1	Total	C	N	S	0	0
			3	1	1	1		
3	CCC	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	AAA	1	Total Cl 1 1	0	0

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	BBB	1	Total C O 4 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	AAA	134	Total O 134 134	0	0
6	BBB	122	Total O 122 122	0	0
6	CCC	68	Total O 68 68	0	1


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tyrosine-protein kinase SYK



- Molecule 1: Tyrosine-protein kinase SYK



- Molecule 1: Tyrosine-protein kinase SYK



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	85.90Å 152.58Å 146.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	76.29 – 1.90 76.29 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (76.29-1.90) 100.0 (76.29-1.90)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.25 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.217 , 0.264 0.224 , 0.268	Depositor DCC
R_{free} test set	1972 reflections (2.59%)	wwPDB-VP
Wilson B-factor (Å ²)	40.9	Xtriage
Anisotropy	0.471	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 37.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.011 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.025 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6380	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, SCN, PEG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.00	0/2152	1.28	1/2900 (0.0%)
1	BBB	1.03	1/2131 (0.0%)	1.29	2/2873 (0.1%)
1	CCC	1.00	0/1895	1.25	1/2550 (0.0%)
All	All	1.01	1/6178 (0.0%)	1.27	4/8323 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	BBB	243	HIS	CE1-NE2	7.11	1.39	1.32

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	197	ARG	NE-CZ-NH2	5.42	124.07	119.20
1	CCC	121	GLU	CB-CA-C	-5.21	100.22	110.11
1	BBB	70	LEU	CA-C-N	5.02	127.71	120.38
1	BBB	70	LEU	C-N-CA	5.02	127.71	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2102	0	2084	19	0
1	BBB	2081	0	2066	16	0
1	CCC	1834	0	1821	15	0
2	AAA	7	0	10	2	0
3	AAA	9	0	0	0	0
3	BBB	12	0	0	0	0
3	CCC	6	0	0	0	0
4	AAA	1	0	0	0	0
5	BBB	4	0	6	6	0
6	AAA	134	0	0	2	0
6	BBB	122	0	0	0	0
6	CCC	68	0	0	2	0
All	All	6380	0	5987	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:CCC:246:TYR:HE2	6:CCC:445:HOH:O	1.63	0.81
1:AAA:79[B]:GLY:O	1:AAA:80[B]:ARG:HB3	1.81	0.80
1:BBB:181:ILE:HG22	1:BBB:259:CYS:SG	2.29	0.73
1:AAA:142:ALA:O	1:AAA:146:ALA:N	2.24	0.69
1:CCC:163:HIS:HA	1:CCC:166[B]:MET:HE2	1.76	0.67
1:CCC:206[B]:CYS:SG	1:CCC:213:VAL:CG1	2.83	0.66
1:BBB:243:HIS:HB2	5:BBB:305:EDO:H22	1.78	0.64
1:CCC:246:TYR:CE2	6:CCC:445:HOH:O	2.42	0.63
1:AAA:67:GLU:OE2	6:AAA:401:HOH:O	2.15	0.63
1:AAA:142:ALA:HB1	1:AAA:146:ALA:HB2	1.82	0.61
1:AAA:142:ALA:HA	1:AAA:145:GLN:HB2	1.86	0.57
1:BBB:234:PHE:HE2	5:BBB:305:EDO:H22	1.70	0.57
1:AAA:220:LYS:HG2	1:AAA:224:GLY:HA2	1.88	0.56
1:AAA:202:SER:OG	6:AAA:402:HOH:O	2.19	0.54
1:CCC:206[B]:CYS:SG	1:CCC:213:VAL:HG11	2.47	0.54
1:CCC:11:ASN:HB3	1:CCC:16:PHE:CD2	2.43	0.53
1:AAA:117:THR:HB	1:AAA:121:GLU:HB3	1.90	0.52
1:BBB:243:HIS:HB2	5:BBB:305:EDO:C2	2.40	0.51
1:BBB:45:ARG:HB3	5:BBB:305:EDO:C2	2.41	0.50
1:BBB:260:GLN:OE1	1:CCC:31:GLN:HG2	2.13	0.49
1:CCC:119:PRO:C	1:CCC:121:GLU:H	2.21	0.48
1:BBB:220:LYS:HG2	1:BBB:224:GLY:HA2	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:260:GLN:OE1	1:CCC:31:GLN:CG	2.63	0.46
1:CCC:163:HIS:O	1:CCC:166[B]:MET:HE2	2.16	0.45
1:CCC:180:GLN:O	1:CCC:184:ILE:HG13	2.15	0.45
1:BBB:123:LEU:HD13	1:BBB:127:LEU:HD11	1.99	0.44
1:AAA:63:HIS:O	2:AAA:301:PEG:H32	2.17	0.44
1:AAA:120:PHE:CZ	1:AAA:124:LYS:HG3	2.52	0.44
1:AAA:151:LYS:N	1:AAA:152:PRO:HD2	2.33	0.44
1:BBB:234:PHE:CE2	5:BBB:305:EDO:H22	2.52	0.44
1:AAA:221:ASP:OD2	1:AAA:233:LYS:NZ	2.51	0.44
1:AAA:144:GLU:O	1:AAA:148:ILE:HG12	2.18	0.43
1:AAA:133:LYS:O	1:AAA:137:ASN:HA	2.19	0.43
1:AAA:223:THR:HG22	1:AAA:223:THR:O	2.18	0.43
1:BBB:45:ARG:HB3	5:BBB:305:EDO:H21	1.99	0.43
1:BBB:221:ASP:OD2	1:BBB:233:LYS:NZ	2.52	0.43
1:CCC:220:LYS:HG2	1:CCC:224:GLY:HA2	1.99	0.43
1:AAA:63:HIS:HB3	2:AAA:301:PEG:H22	2.01	0.43
1:BBB:71:ASN:OD1	1:BBB:71:ASN:N	2.40	0.42
1:CCC:62:HIS:ND1	1:CCC:100:VAL:HG11	2.34	0.42
1:CCC:182:VAL:HG21	1:CCC:206[B]:CYS:SG	2.60	0.42
1:AAA:139:GLN:C	1:AAA:143:LEU:HD22	2.44	0.42
1:BBB:133:LYS:O	1:BBB:137:ASN:HA	2.20	0.41
1:AAA:26:GLU:HG2	1:AAA:61:ALA:CB	2.51	0.41
1:BBB:151:LYS:N	1:BBB:152:PRO:HD2	2.36	0.41
1:BBB:193:LEU:C	1:BBB:193:LEU:HD12	2.46	0.41
1:AAA:193:LEU:HD12	1:AAA:193:LEU:C	2.46	0.40
1:CCC:92:HIS:HA	1:CCC:95:GLU:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	264/265 (100%)	249 (94%)	12 (4%)	3 (1%)	11	4
1	BBB	261/265 (98%)	253 (97%)	6 (2%)	2 (1%)	16	8
1	CCC	227/265 (86%)	222 (98%)	4 (2%)	1 (0%)	30	22
All	All	752/795 (95%)	724 (96%)	22 (3%)	6 (1%)	24	8

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	80[A]	ARG
1	AAA	80[B]	ARG
1	CCC	120	PHE
1	AAA	143	LEU
1	BBB	79[A]	GLY
1	BBB	79[B]	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	222/224 (99%)	222 (100%)	0	100	100
1	BBB	220/224 (98%)	217 (99%)	3 (1%)	59	59
1	CCC	197/224 (88%)	196 (100%)	1 (0%)	81	84
All	All	639/672 (95%)	635 (99%)	4 (1%)	78	81

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	BBB	71	ASN
1	BBB	80	ARG
1	BBB	135	THR
1	CCC	181	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SCN	AAA	302	-	1,2,2	0.25	0	0,1,1	-	-
3	SCN	AAA	304	-	1,2,2	0.67	0	0,1,1	-	-
3	SCN	BBB	302	-	1,2,2	0.98	0	0,1,1	-	-
3	SCN	CCC	301	-	1,2,2	0.18	0	0,1,1	-	-
3	SCN	BBB	303	-	1,2,2	0.56	0	0,1,1	-	-
3	SCN	BBB	304	-	1,2,2	0.74	0	0,1,1	-	-
3	SCN	CCC	302	-	1,2,2	0.79	0	0,1,1	-	-
5	EDO	BBB	305	-	3,3,3	0.45	0	2,2,2	1.06	0
2	PEG	AAA	301	-	6,6,6	0.48	0	5,5,5	0.27	0
3	SCN	AAA	303	-	1,2,2	0.65	0	0,1,1	-	-
3	SCN	BBB	301	-	1,2,2	0.56	0	0,1,1	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	AAA	301	-	-	3/4/4/4	-
5	EDO	BBB	305	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	AAA	301	PEG	O2-C3-C4-O4
2	AAA	301	PEG	C1-C2-O2-C3
2	AAA	301	PEG	C4-C3-O2-C2
5	BBB	305	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	BBB	305	EDO	6	0
2	AAA	301	PEG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	259/265 (97%)	0.71	32 (12%) 8 8	20, 46, 116, 157	7 (2%)
1	BBB	259/265 (97%)	0.64	15 (5%) 29 30	29, 47, 73, 91	4 (1%)
1	CCC	225/265 (84%)	1.23	46 (20%) 3 2	28, 56, 105, 129	6 (2%)
All	All	743/795 (93%)	0.84	93 (12%) 8 8	20, 50, 98, 157	17 (2%)

All (93) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	143	LEU	5.9
1	AAA	148	ILE	5.7
1	CCC	127	LEU	5.0
1	CCC	120	PHE	5.0
1	CCC	128	ILE	4.9
1	CCC	131	TYR	4.5
1	CCC	162	ALA	4.5
1	CCC	122	ASP	4.3
1	CCC	123	LEU	4.1
1	AAA	147	ILE	4.1
1	CCC	70	LEU	4.1
1	BBB	78[A]	GLY	4.0
1	AAA	78[A]	GLY	3.9
1	CCC	264	THR	3.8
1	AAA	140	GLY	3.7
1	CCC	161	THR	3.6
1	AAA	7	ALA	3.6
1	AAA	262	ILE	3.5
1	CCC	83	ALA	3.5
1	AAA	181	ILE	3.4
1	CCC	85	PRO	3.4
1	AAA	136	TRP	3.3
1	CCC	89	CYS	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	BBB	262	ILE	3.1
1	CCC	12	HIS	3.0
1	CCC	91	TYR	2.9
1	BBB	263	GLY	2.9
1	AAA	123	LEU	2.9
1	CCC	164	GLU	2.8
1	AAA	142	ALA	2.8
1	AAA	146	ALA	2.7
1	AAA	199	ASN	2.7
1	CCC	93	SER	2.7
1	BBB	123	LEU	2.7
1	CCC	199	ASN	2.7
1	AAA	175	ARG	2.6
1	CCC	88	LEU	2.6
1	BBB	79[A]	GLY	2.6
1	CCC	126	ASN	2.6
1	AAA	127	LEU	2.6
1	AAA	132	VAL	2.6
1	AAA	135	THR	2.6
1	CCC	66	ILE	2.5
1	CCC	223	THR	2.5
1	AAA	6	MET	2.5
1	AAA	128	ILE	2.5
1	CCC	11	ASN	2.5
1	CCC	39	TYR	2.5
1	CCC	201	GLY	2.5
1	CCC	101	CYS	2.4
1	BBB	59	ARG	2.4
1	AAA	157	LEU	2.4
1	CCC	232[A]	LYS	2.4
1	BBB	136	TRP	2.4
1	CCC	100	VAL	2.4
1	CCC	184	ILE	2.4
1	CCC	86	ALA	2.4
1	CCC	112	GLY	2.4
1	BBB	70	LEU	2.3
1	CCC	222	LYS	2.3
1	AAA	138	LEU	2.3
1	CCC	102	LEU	2.3
1	CCC	163	HIS	2.3
1	AAA	184	ILE	2.3
1	CCC	55	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	CCC	113	VAL	2.3
1	CCC	73	THR	2.3
1	CCC	107	PHE	2.3
1	BBB	6	MET	2.3
1	CCC	13	LEU	2.3
1	BBB	112	GLY	2.2
1	BBB	7	ALA	2.2
1	AAA	129	ARG	2.2
1	AAA	141	GLN	2.2
1	CCC	181	ILE	2.2
1	AAA	223	THR	2.2
1	CCC	33	GLY	2.2
1	CCC	72	GLY	2.2
1	AAA	131	TYR	2.1
1	AAA	246	TYR	2.1
1	CCC	262	ILE	2.1
1	AAA	5	SER	2.1
1	BBB	33	GLY	2.1
1	CCC	263	GLY	2.1
1	AAA	120	PHE	2.1
1	AAA	112	GLY	2.1
1	AAA	200	ASN	2.1
1	CCC	103	LEU	2.1
1	BBB	77[A]	ALA	2.0
1	BBB	60	LYS	2.0
1	AAA	263	GLY	2.0
1	BBB	19	ASN	2.0
1	CCC	197	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	BBB	305	4/4	0.83	0.19	48,63,64,74	0
2	PEG	AAA	301	7/7	0.85	0.14	42,53,70,70	0
3	SCN	BBB	301	3/3	0.87	0.15	92,92,96,97	0
3	SCN	BBB	304	3/3	0.93	0.19	56,56,60,67	0
3	SCN	AAA	303	3/3	0.93	0.12	57,57,73,76	0
3	SCN	BBB	302	3/3	0.94	0.19	43,43,59,66	0
3	SCN	CCC	301	3/3	0.95	0.10	38,38,49,52	0
3	SCN	CCC	302	3/3	0.95	0.11	56,56,69,72	0
3	SCN	AAA	304	3/3	0.95	0.21	51,51,68,71	0
3	SCN	BBB	303	3/3	0.97	0.15	49,49,54,64	0
4	CL	AAA	305	1/1	0.98	0.06	56,56,56,56	0
3	SCN	AAA	302	3/3	0.98	0.08	43,43,49,50	3

6.5 Other polymers [i](#)

There are no such residues in this entry.