



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 04:45 AM UTC

PDB ID : 7Q6C / pdb_00007q6c
Title : complement C6 FIM1-2 bound to CP010 antibody
Authors : Olesen, H.G.; Andersen, G.R.
Deposited on : 2021-11-06
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	166	12% 90% 7% ..
2	H	219	8% 95% 5%
3	K	122	11% 99% .
4	L	219	% 98% .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 10943 atoms, of which 5289 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Complement component C6.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	163	Total	C	H	N	O	S	0	0	0
			2436	766	1181	219	249	21			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	855	ALA	ASN	engineered mutation	UNP P13671

- Molecule 2 is a protein called CP010 heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	H	219	Total	C	H	N	O	S	0	0	0
			3248	1038	1597	275	331	7			

- Molecule 3 is a protein called kappa specific nanobody.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	K	122	Total	C	H	N	O	S	0	0	0
			1821	582	881	165	189	4			

- Molecule 4 is a protein called CP010 light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	L	219	Total	C	H	N	O	S	0	0	0
			3303	1050	1627	279	341	6			

- Molecule 5 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			7	2	3	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	21	Total	O	0	0
			21	21		
6	H	41	Total	O	0	0
			41	41		
6	K	12	Total	O	0	0
			12	12		
6	L	54	Total	O	0	0
			54	54		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Complement component C6

Chain A: 





- Molecule 2: CP010 heavy chain

Chain H: 





- Molecule 3: kappa specific nanobody

Chain K: 



- Molecule 4: CP010 light chain

Chain L: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	161.06Å 63.53Å 128.59Å 90.00° 128.36° 90.00°	Depositor
Resolution (Å)	44.45 – 2.29 44.45 – 2.29	Depositor EDS
% Data completeness (in resolution range)	97.9 (44.45-2.29) 97.9 (44.45-2.29)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.26 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.208 , 0.233 0.211 , 0.236	Depositor DCC
R_{free} test set	1988 reflections (4.32%)	wwPDB-VP
Wilson B-factor (Å ²)	49.6	Xtriage
Anisotropy	0.210	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 32.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.002 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10943	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/1278	0.48	0/1718
2	H	0.38	0/1691	0.53	0/2305
3	K	0.24	0/959	0.41	0/1296
4	L	0.24	0/1712	0.46	0/2330
All	All	0.30	0/5640	0.48	0/7649

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1255	1181	1181	4	0
2	H	1651	1597	1598	5	0
3	K	940	881	881	0	0
4	L	1676	1627	1628	2	0
5	A	4	3	3	0	0
6	A	21	0	0	1	0
6	H	41	0	0	1	0
6	K	12	0	0	0	0
6	L	54	0	0	1	0
All	All	5654	5289	5291	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:899:LYS:HG3	1:A:927:LEU:HD11	1.85	0.59
2:H:12:VAL:HG11	2:H:86:LEU:HD13	1.86	0.58
2:H:149:ASP:OD1	2:H:176:GLN:NE2	2.42	0.52
2:H:67:ARG:NH2	2:H:90:ASP:OD2	2.41	0.52
4:L:163:ASN:ND2	6:L:302:HOH:O	2.36	0.50
4:L:41:TYR:HE1	4:L:94:MET:HE3	1.77	0.50
1:A:914:VAL:HG22	1:A:924:MET:HE1	1.94	0.49
2:H:215:ARG:NH1	6:H:306:HOH:O	2.47	0.46
1:A:893:ASN:HB3	1:A:911:ILE:HG22	1.98	0.44
2:H:155:VAL:CG2	2:H:183:LEU:HD21	2.48	0.43
1:A:817:LYS:NZ	6:A:1101:HOH:O	2.51	0.42

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	161/166 (97%)	150 (93%)	10 (6%)	1 (1%)	21	27
2	H	217/219 (99%)	210 (97%)	7 (3%)	0	100	100
3	K	120/122 (98%)	116 (97%)	4 (3%)	0	100	100
4	L	217/219 (99%)	213 (98%)	4 (2%)	0	100	100
All	All	715/726 (98%)	689 (96%)	25 (4%)	1 (0%)	48	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	891	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/148 (99%)	140 (96%)	6 (4%)	27	41
2	H	185/185 (100%)	183 (99%)	2 (1%)	65	81
3	K	98/98 (100%)	97 (99%)	1 (1%)	68	82
4	L	193/193 (100%)	191 (99%)	2 (1%)	68	82
All	All	622/624 (100%)	611 (98%)	11 (2%)	51	70

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	890	LYS
1	A	894	GLN
1	A	898	VAL
1	A	923	LYS
1	A	924	MET
1	A	925	GLU
2	H	64	VAL
2	H	132	CYS
3	K	16	VAL
4	L	38	LEU
4	L	168	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	82	GLN
4	L	50	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ACT	A	1001	-	3,3,3	1.07	0	3,3,3	1.71	2 (66%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1001	ACT	O-C-CH3	-2.11	113.86	122.53
5	A	1001	ACT	OXT-C-O	2.04	129.60	122.03

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	163/166 (98%)	0.87	20 (12%) 8 9	53, 74, 114, 139	0
2	H	219/219 (100%)	0.47	17 (7%) 19 21	43, 59, 98, 178	0
3	K	122/122 (100%)	0.60	13 (10%) 11 12	43, 65, 102, 141	0
4	L	219/219 (100%)	0.06	2 (0%) 81 82	40, 56, 82, 121	0
All	All	723/726 (99%)	0.46	52 (7%) 21 23	40, 62, 102, 178	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	934	ALA	5.5
2	H	218	SER	5.3
3	K	126	GLU	5.1
2	H	219	SER	4.5
1	A	926	ILE	4.4
4	L	219	CYS	4.3
1	A	904	THR	3.9
4	L	163	ASN	3.9
2	H	216	VAL	3.9
1	A	772	HIS	3.9
2	H	134	ARG	3.8
2	H	217	GLU	3.4
1	A	894	GLN	3.3
2	H	215	ARG	3.3
2	H	196	THR	3.2
1	A	889	PHE	3.2
2	H	136	THR	3.2
3	K	6	VAL	3.1
3	K	124	SER	2.9
2	H	212	VAL	2.9
2	H	135	SER	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	210	THR	2.7
2	H	214	LYS	2.7
1	A	893	ASN	2.6
2	H	213	ASP	2.6
1	A	857	THR	2.6
1	A	891	GLY	2.6
3	K	125	SER	2.6
1	A	923	LYS	2.6
1	A	901	GLY	2.5
3	K	21	SER	2.5
1	A	900	MET	2.5
1	A	892	GLY	2.4
3	K	22	LEU	2.4
1	A	933	LEU	2.4
2	H	195	GLY	2.3
3	K	58	SER	2.3
3	K	5	GLN	2.3
1	A	782	SER	2.2
1	A	785	ILE	2.2
2	H	145	CYS	2.2
2	H	137	SER	2.1
1	A	902	SER	2.1
3	K	45	PRO	2.1
1	A	775	LEU	2.1
1	A	834	ILE	2.1
3	K	18	PRO	2.1
3	K	102	ILE	2.0
3	K	70	GLY	2.0
1	A	784	CYS	2.0
3	K	17	GLN	2.0
2	H	211	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	ACT	A	1001	4/4	0.70	0.30	53,57,64,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.