



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:36 PM UTC

PDB ID : 7Q6S / pdb_00007q6s
Title : Keap1 compound complex
Authors : Johansson, P.
Deposited on : 2021-11-09
Resolution : 2.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3689 (2.16-2.12)
Clashscore	190562	3812 (2.16-2.12)
Ramachandran outliers	187476	3773 (2.16-2.12)
Sidechain outliers	187428	3772 (2.16-2.12)
RSRZ outliers	180081	3691 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	4% 85% 10% ••
1	B	295	7% 86% 10% •

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4602 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Kelch-like ECH-associated protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	285	Total	C	N	O	S	0	0	0
			2186	1359	397	415	15			
1	B	285	Total	C	N	O	S	0	0	0
			2186	1359	397	415	15			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	315	PRO	-	expression tag	UNP Q14145
A	316	ARG	-	expression tag	UNP Q14145
A	317	GLY	-	expression tag	UNP Q14145
A	318	SER	-	expression tag	UNP Q14145
A	319	ALA	-	expression tag	UNP Q14145
A	320	GLY	-	expression tag	UNP Q14145
A	540	ALA	GLU	conflict	UNP Q14145
A	542	ALA	GLU	conflict	UNP Q14145
B	315	PRO	-	expression tag	UNP Q14145
B	316	ARG	-	expression tag	UNP Q14145
B	317	GLY	-	expression tag	UNP Q14145
B	318	SER	-	expression tag	UNP Q14145
B	319	ALA	-	expression tag	UNP Q14145
B	320	GLY	-	expression tag	UNP Q14145
B	540	ALA	GLU	conflict	UNP Q14145
B	542	ALA	GLU	conflict	UNP Q14145

- Molecule 2 is (5S,8R)-16-(2,1,3-benzoxadiazol-4-yl)-8-[[[(2S)-1-ethanoylpyrrolidin-2-yl]carbamoylamino]-N,N-dimethyl-7,11-bis(oxidanylidene)-10-oxa-3-thia-6-azabicyclo[10.4.0]hexadeca-1(16),12,14-triene-5-carboxamide (CCD ID: 91M) (formula: C₂₉H₃₂N₆O₇S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			43	29	6	7	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	100	Total	O	0	0
			100	100		
4	B	86	Total	O	0	0
			86	86		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kelch-like ECH-associated protein 1



- Molecule 1: Kelch-like ECH-associated protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	75.40Å 75.57Å 202.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	101.06 – 2.14 101.06 – 2.14	Depositor EDS
% Data completeness (in resolution range)	66.0 (101.06-2.14) 66.0 (101.06-2.14)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.36 (at 2.14Å)	Xtriage
Refinement program	BUSTER 2.11.8 (24-FEB-2021)	Depositor
R, R_{free}	0.211 , 0.232 0.197 , 0.217	Depositor DCC
R_{free} test set	2122 reflections (3.30%)	wwPDB-VP
Wilson B-factor (Å ²)	62.3	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.021 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4602	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 91M, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/2239	0.99	2/3048 (0.1%)
1	B	0.71	0/2239	1.02	2/3048 (0.1%)
All	All	0.71	0/4478	1.01	4/6096 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	385	ASP	CA-CB-CG	7.34	119.94	112.60
1	B	573	ASP	CA-CB-CG	5.56	118.16	112.60
1	A	484	LEU	N-CA-C	5.41	118.30	109.59
1	B	438	ASN	N-CA-C	-5.08	107.14	113.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2186	0	2081	18	0
1	B	2186	0	2081	15	0
2	A	43	0	0	1	0
3	A	1	0	0	1	0
4	A	100	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	86	0	0	0	0
All	All	4602	0	4162	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:508:SER:OG	2:A:701:91M:N22	2.26	0.68
1:A:362:ARG:HD2	1:A:378:GLY:O	2.05	0.56
1:A:483:ARG:HB3	1:A:506:ILE:CG2	2.36	0.56
1:A:361:PRO:O	1:A:362:ARG:HG3	2.09	0.52
1:A:336:ARG:CZ	1:B:336:ARG:HD3	2.42	0.50
1:A:369:VAL:HG21	1:A:609:THR:HB	1.95	0.48
1:B:369:VAL:HG21	1:B:609:THR:HB	1.96	0.48
1:B:416:ILE:HD11	1:B:427:ALA:HB1	1.95	0.48
1:A:515:LEU:HD22	1:A:566:ILE:HG13	1.97	0.47
1:B:515:LEU:HD22	1:B:566:ILE:HG13	1.97	0.46
1:A:415:ARG:NH1	3:A:702:CL:CL	2.86	0.45
1:B:562:HIS:CE1	1:B:563:GLN:HE21	2.34	0.45
1:A:471:LEU:HD13	1:A:489:CYS:SG	2.57	0.45
1:A:566:ILE:HB	1:A:584:TYR:HB3	1.98	0.45
1:A:575:HIS:ND1	1:B:385:ASP:OD1	2.50	0.45
1:B:566:ILE:HB	1:B:584:TYR:HB3	1.98	0.45
1:B:557:LEU:H	1:B:557:LEU:HD23	1.82	0.44
1:A:416:ILE:HD11	1:A:427:ALA:HB1	2.00	0.44
1:B:362:ARG:HD2	1:B:378:GLY:O	2.17	0.44
1:A:385:ASP:O	1:B:415:ARG:NH2	2.51	0.43
1:B:335:PHE:C	1:B:337:GLN:H	2.26	0.43
1:A:557:LEU:HD23	1:A:557:LEU:H	1.83	0.43
1:B:486:SER:HB2	1:B:499:MET:HE1	2.00	0.43
1:A:486:SER:HB2	1:A:499:MET:HE1	2.02	0.42
1:B:362:ARG:NH2	1:B:394:ASP:OD2	2.53	0.41
1:A:472:LEU:HB3	1:A:490:TYR:HB3	2.03	0.41
1:B:525:TYR:HA	1:B:529:ASP:O	2.21	0.41
1:A:335:PHE:C	1:A:337:GLN:H	2.29	0.41
1:B:471:LEU:HD13	1:B:489:CYS:SG	2.61	0.40
1:A:483:ARG:HB3	1:A:506:ILE:HG21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/295 (96%)	276 (98%)	7 (2%)	0	100	100
1	B	283/295 (96%)	277 (98%)	6 (2%)	0	100	100
All	All	566/590 (96%)	553 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/235 (97%)	220 (96%)	9 (4%)	28	26
1	B	229/235 (97%)	225 (98%)	4 (2%)	53	58
All	All	458/470 (97%)	445 (97%)	13 (3%)	38	39

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	380	ARG
1	A	385	ASP
1	A	415	ARG
1	A	461	ILE
1	A	483	ARG
1	A	514	VAL
1	A	530	GLN
1	A	539	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	593	GLU
1	B	461	ILE
1	B	514	VAL
1	B	539	VAL
1	B	593	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	552	HIS
1	A	562	HIS
1	B	382	ASN
1	B	414	ASN
1	B	562	HIS
1	B	575	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	91M	A	701	-	47,47,47	1.19	6 (12%)	61,67,67	1.73	17 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	91M	A	701	-	-	6/47/57/57	0/4/5/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	91M	C4-N2	3.74	1.39	1.34
2	A	701	91M	C41-N40	3.04	1.39	1.35
2	A	701	91M	C44-N46	2.31	1.39	1.34
2	A	701	91M	C33-N32	2.22	1.38	1.34
2	A	701	91M	C17-C25	-2.21	1.41	1.43
2	A	701	91M	C25-N24	2.14	1.34	1.31

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	91M	C13-C12-C26	-3.87	110.82	118.66
2	A	701	91M	C18-C17-C25	-3.60	114.54	117.52
2	A	701	91M	C19-C20-C21	-3.36	114.92	119.86
2	A	701	91M	C11-C12-C26	3.28	129.15	122.25
2	A	701	91M	C12-C11-C16	-3.26	116.03	118.97
2	A	701	91M	O42-C41-N40	3.18	124.12	120.45
2	A	701	91M	C15-C16-C17	-2.91	112.86	118.74
2	A	701	91M	C18-C17-C16	2.90	124.60	118.74
2	A	701	91M	C33-C35-N40	-2.87	104.66	112.50
2	A	701	91M	C43-C41-N40	-2.84	114.92	117.99
2	A	701	91M	C4-C6-N46	-2.80	101.71	108.97
2	A	701	91M	C15-C16-C11	2.61	122.93	119.02
2	A	701	91M	O28-C29-C30	2.42	115.06	108.26
2	A	701	91M	O28-C26-C12	2.33	116.81	112.24
2	A	701	91M	C35-C33-N32	-2.18	111.73	116.52
2	A	701	91M	C29-O28-C26	-2.06	112.91	116.45
2	A	701	91M	C37-C35-N40	2.05	106.03	103.02

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	701	91M	O28-C29-C30-N32
2	A	701	91M	O42-C41-N40-C39
2	A	701	91M	C43-C41-N40-C39
2	A	701	91M	C12-C26-O28-C29
2	A	701	91M	O27-C26-O28-C29
2	A	701	91M	O28-C29-C30-C44

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	91M	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	285/295 (96%)	0.54	12 (4%) 40 46	50, 62, 84, 98	0
1	B	285/295 (96%)	0.58	20 (7%) 22 26	51, 63, 83, 98	0
All	All	570/590 (96%)	0.56	32 (5%) 30 34	50, 63, 84, 98	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	325	GLY	5.3
1	A	325	GLY	4.9
1	B	526	ASP	4.8
1	B	525	TYR	4.3
1	A	575	HIS	4.2
1	B	384	PRO	4.1
1	B	528	GLN	4.0
1	B	383	SER	3.8
1	B	529	ASP	3.5
1	B	387	ASN	3.4
1	A	385	ASP	3.2
1	B	551	LYS	3.0
1	A	336	ARG	3.0
1	B	527	GLY	3.0
1	B	489	CYS	2.8
1	B	341	TYR	2.7
1	B	326	ARG	2.7
1	A	326	ARG	2.7
1	A	384	PRO	2.6
1	B	575	HIS	2.5
1	B	385	ASP	2.3
1	A	516	HIS	2.3
1	A	552	HIS	2.3
1	B	521	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	576	THR	2.2
1	A	498	ARG	2.1
1	B	606	VAL	2.1
1	B	447	ARG	2.0
1	B	588	THR	2.0
1	A	528	GLN	2.0
1	B	415	ARG	2.0
1	A	527	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	91M	A	701	43/43	0.93	0.14	77,80,91,93	0
3	CL	A	702	1/1	0.97	0.09	79,79,79,79	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 91M A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**6.5 Other polymers** ⓘ

There are no such residues in this entry.