



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 12:35 AM UTC

PDB ID : 2Q70 / pdb_00002q70
Title : Estrogen receptor alpha ligand-binding domain complexed to a benzopyran lig-
and
Authors : Wang, Y.
Deposited on : 2007-06-05
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	21% 69% 24% • 5%
1	B	248	16% 63% 21% • 15%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3641 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Estrogen receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	235	Total	C	N	O	S	0	0	0
			1870	1201	317	336	16			
1	B	211	Total	C	N	O	S	0	0	0
			1681	1077	288	301	15			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	381	SER	CYS	engineered mutation	UNP P03372
A	417	SER	CYS	engineered mutation	UNP P03372
A	530	SER	CYS	engineered mutation	UNP P03372
B	381	SER	CYS	engineered mutation	UNP P03372
B	417	SER	CYS	engineered mutation	UNP P03372
B	530	SER	CYS	engineered mutation	UNP P03372

- Molecule 2 is (3AS,4R,9BR)-2,2-DIFLUORO-4-(4-HYDROXYPHENYL)-1,2,3,3A,4,9B-HE XAHYDROCYCLOPENTA[C]CHROMEN-8-OL (CCD ID: DC8) (formula: C₁₈H₁₆F₂O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	F	O	0	0
			23	18	2	3		
2	B	1	Total	C	F	O	0	0
			23	18	2	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	26	Total	O	0	0
			26	26		
3	B	18	Total	O	0	0
			18	18		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Estrogen receptor



• Molecule 1: Estrogen receptor



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	62.56Å 97.88Å 174.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.93 – 1.95 28.93 – 1.95	Depositor EDS
% Data completeness (in resolution range)	94.3 (28.93-1.95) 94.6 (28.93-1.95)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 1.95Å)	Xtriage
Refinement program	CNX 2005	Depositor
R, R_{free}	0.260 , 0.303 0.254 , 0.294	Depositor DCC
R_{free} test set	1036 reflections (2.78%)	wwPDB-VP
Wilson B-factor (Å ²)	36.7	Xtriage
Anisotropy	0.503	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3641	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DC8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/1903	0.73	0/2571
1	B	0.44	0/1709	0.67	0/2304
All	All	0.44	0/3612	0.70	0/4875

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1870	0	1916	51	0
1	B	1681	0	1732	45	0
2	A	23	0	14	0	0
2	B	23	0	14	0	0
3	A	26	0	0	0	0
3	B	18	0	0	0	0
All	All	3641	0	3676	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:401:LYS:HD3	1:B:409:LEU:HD21	1.24	1.18
1:B:343:MET:HE2	1:B:528:MET:HE1	1.35	1.07
1:B:342:MET:HE1	1:B:410:LEU:HD13	1.50	0.93
1:A:434:ARG:HH11	1:A:437:MET:HE3	1.33	0.92
1:A:519:ASN:HD22	1:B:519:ASN:HD22	1.19	0.89
1:A:519:ASN:HD22	1:B:519:ASN:ND2	1.78	0.80
1:A:519:ASN:ND2	1:B:519:ASN:HD22	1.82	0.76
1:A:372:LEU:HD11	1:A:545:ASP:HB2	1.69	0.74
1:A:343:MET:HE2	1:A:418:VAL:HG11	1.70	0.72
1:A:415:GLY:O	1:A:421:MET:HG2	1.89	0.71
1:B:376:VAL:O	1:B:380:GLU:HG3	1.91	0.70
1:A:466:LEU:O	1:A:470:GLU:HG3	1.92	0.69
1:A:434:ARG:NH1	1:A:437:MET:HE3	2.10	0.67
1:A:522:MET:HA	1:A:522:MET:HE2	1.75	0.67
1:A:315:MET:CE	1:A:365:PRO:HG2	2.25	0.66
1:B:377:HIS:CE1	1:B:460:THR:HG23	2.30	0.66
1:A:315:MET:HE1	1:A:365:PRO:HG2	1.78	0.66
1:A:315:MET:HE2	1:A:319:LEU:HD11	1.79	0.65
1:A:539:LEU:CD2	1:A:543:MET:HE2	2.27	0.65
1:B:488:HIS:NE2	1:B:492:LYS:HE3	2.12	0.64
1:A:438:MET:HE3	1:A:506:GLN:HE21	1.63	0.62
1:B:496:THR:OG1	1:B:499:GLN:HG3	2.00	0.62
1:B:396:MET:HE3	1:B:436:ARG:HA	1.84	0.60
1:B:343:MET:HE2	1:B:528:MET:CE	2.21	0.59
1:A:379:LEU:HD12	1:A:544:LEU:HD11	1.85	0.58
1:A:311:THR:OG1	1:A:314:GLN:HG3	2.04	0.57
1:A:360:TRP:CZ2	1:A:449:LYS:HD2	2.41	0.55
1:B:438:MET:HE3	1:B:506:GLN:HE21	1.70	0.55
1:A:539:LEU:HD22	1:A:543:MET:HE2	1.89	0.55
1:A:390:GLY:O	1:A:394:ARG:HG2	2.06	0.55
1:A:412:ARG:NH1	1:A:416:LYS:HE3	2.22	0.55
1:B:416:LYS:HA	1:B:422:VAL:HG22	1.87	0.55
1:A:535:PRO:HG2	1:A:538:ASP:OD2	2.07	0.54
1:A:343:MET:CE	1:A:418:VAL:HG11	2.38	0.53
1:A:412:ARG:HH12	1:A:416:LYS:CE	2.21	0.53
1:B:308:LEU:O	1:B:481:LYS:HE2	2.09	0.53
1:A:391:LEU:HD13	1:A:404:PHE:HA	1.91	0.53
1:B:392:VAL:HG13	1:B:432:SER:HA	1.90	0.53
1:B:515:ARG:HB2	1:B:515:ARG:HH11	1.75	0.52
1:A:418:VAL:HB	1:A:421:MET:SD	2.50	0.51
1:A:513:HIS:HB2	1:B:459:TYR:CD2	2.45	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:502:GLN:O	1:A:506:GLN:HG3	2.09	0.51
1:A:315:MET:HE1	1:A:365:PRO:CG	2.40	0.51
1:A:376:VAL:HG22	1:A:544:LEU:HD12	1.92	0.51
1:B:412:ARG:HH12	1:B:416:LYS:HE2	1.75	0.51
1:B:416:LYS:HA	1:B:422:VAL:CG2	2.41	0.51
1:B:455:ASN:O	1:B:458:VAL:HG12	2.11	0.51
1:A:459:TYR:CD2	1:B:513:HIS:HB2	2.45	0.50
1:B:403:LEU:HD11	1:B:406:PRO:HA	1.94	0.50
1:A:536:LEU:O	1:A:540:LEU:HG	2.12	0.49
1:A:372:LEU:HD11	1:A:545:ASP:CB	2.39	0.49
1:B:412:ARG:NH1	1:B:416:LYS:HE2	2.28	0.49
1:A:315:MET:HE2	1:A:319:LEU:CD1	2.43	0.48
1:B:374:ASP:OD2	1:B:471:GLU:OE1	2.31	0.48
1:A:518:SER:O	1:A:522:MET:HG2	2.13	0.48
1:A:424:ILE:HA	1:A:427:MET:HE3	1.96	0.48
1:A:512:SER:OG	1:B:515:ARG:NH1	2.47	0.47
1:B:360:TRP:CZ2	1:B:449:LYS:HD2	2.51	0.46
1:A:539:LEU:O	1:A:539:LEU:HD23	2.16	0.46
1:B:310:LEU:O	1:B:481:LYS:HE3	2.16	0.46
1:B:377:HIS:HE1	1:B:457:GLY:O	1.99	0.45
1:A:424:ILE:HD13	1:A:427:MET:CE	2.46	0.45
1:B:504:LEU:O	1:B:508:LEU:HG	2.15	0.45
1:B:411:ASP:OD1	1:B:411:ASP:C	2.59	0.45
1:A:423:GLU:H	1:A:423:GLU:CD	2.24	0.45
1:B:396:MET:CE	1:B:436:ARG:HA	2.46	0.45
1:B:409:LEU:HD23	1:B:409:LEU:O	2.17	0.45
1:A:321:ASP:C	1:A:363:ARG:HH21	2.24	0.44
1:A:392:VAL:HG12	1:A:435:PHE:CD2	2.52	0.44
1:A:315:MET:HE1	1:A:365:PRO:HD2	2.00	0.44
1:B:495:LEU:HB3	1:B:499:GLN:HB2	1.99	0.44
1:B:421:MET:HG2	1:B:425:PHE:CD2	2.52	0.44
1:B:412:ARG:HH11	1:B:412:ARG:HG2	1.83	0.43
1:A:487:ILE:HD11	1:A:504:LEU:HD22	2.00	0.43
1:A:306:LEU:HD23	1:A:306:LEU:N	2.34	0.43
1:A:396:MET:O	1:A:436:ARG:HD3	2.20	0.42
1:B:409:LEU:HD23	1:B:409:LEU:C	2.45	0.42
1:B:392:VAL:HG11	1:B:431:THR:HG22	2.00	0.42
1:B:396:MET:HE3	1:B:435:PHE:C	2.44	0.42
1:A:308:LEU:O	1:A:481:LYS:HE2	2.19	0.42
1:B:377:HIS:HE1	1:B:460:THR:HG23	1.82	0.42
1:B:315:MET:SD	1:B:365:PRO:HG2	2.60	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:308:LEU:HA	1:B:481:LYS:HD3	2.02	0.41
1:A:467:LYS:O	1:A:471:GLU:HG2	2.20	0.41
1:B:392:VAL:HG13	1:B:432:SER:CA	2.49	0.41
1:A:424:ILE:HD13	1:A:427:MET:HE3	2.01	0.41
1:A:328:TYR:CE2	1:A:406:PRO:HB2	2.55	0.41
1:A:392:VAL:HG13	1:A:432:SER:HA	2.04	0.40
1:B:461:PHE:HD1	1:B:461:PHE:HA	1.75	0.40
1:B:474:HIS:O	1:B:478:VAL:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/248 (92%)	224 (98%)	4 (2%)	1 (0%)	30	21
1	B	205/248 (83%)	203 (99%)	2 (1%)	0	100	100
All	All	434/496 (88%)	427 (98%)	6 (1%)	1 (0%)	43	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	462	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/223 (94%)	203 (97%)	6 (3%)	37	29
1	B	189/223 (85%)	183 (97%)	6 (3%)	34	25
All	All	398/446 (89%)	386 (97%)	12 (3%)	36	27

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	372	LEU
1	A	413	ASN
1	A	421	MET
1	A	466	LEU
1	A	515	ARG
1	A	536	LEU
1	B	372	LEU
1	B	418	VAL
1	B	419	GLU
1	B	425	PHE
1	B	461	PHE
1	B	515	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	377	HIS
1	A	502	GLN
1	A	506	GLN
1	B	375	GLN
1	B	377	HIS
1	B	413	ASN
1	B	501	HIS
1	B	502	GLN
1	B	506	GLN
1	B	519	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	DC8	A	50	-	24,26,26	1.38	3 (12%)	34,40,40	0.84	1 (2%)
2	DC8	B	51	-	24,26,26	1.49	5 (20%)	34,40,40	0.83	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DC8	A	50	-	-	0/4/30/30	0/4/4/4
2	DC8	B	51	-	-	0/4/30/30	0/4/4/4

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	50	DC8	C8-C6	2.76	1.57	1.53
2	A	50	DC8	C14-C15	2.55	1.58	1.54
2	B	51	DC8	C8-C6	2.38	1.56	1.53
2	B	51	DC8	C18-C1	2.31	1.42	1.39
2	B	51	DC8	C14-C15	2.29	1.57	1.54
2	A	50	DC8	C18-C1	2.23	1.42	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	51	DC8	C20-C19	2.18	1.42	1.39
2	B	51	DC8	C18-C17	2.06	1.43	1.39

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	51	DC8	C14-C15-C17	-2.68	109.88	117.30
2	A	50	DC8	C14-C15-C17	-2.52	110.32	117.30

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand DC8 A 50



Ligand DC8 B 51



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	235/248 (94%)	1.30	53 (22%) 2 2	29, 44, 73, 83	0
1	B	211/248 (85%)	1.13	40 (18%) 3 3	30, 43, 66, 81	0
All	All	446/496 (89%)	1.22	93 (20%) 2 2	29, 44, 71, 83	0

All (93) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	462	LEU	6.9
1	A	536	LEU	5.5
1	A	549	LEU	5.5
1	B	306	LEU	5.5
1	A	306	LEU	5.1
1	A	525	LEU	4.9
1	B	526	TYR	4.9
1	B	528	MET	4.8
1	A	461	PHE	4.5
1	A	340	ALA	4.5
1	B	459	TYR	4.3
1	A	535	PRO	4.3
1	A	534	VAL	4.1
1	A	537	TYR	4.1
1	A	307	ALA	4.0
1	A	526	TYR	4.0
1	B	461	PHE	4.0
1	B	418	VAL	3.9
1	A	413	ASN	3.9
1	B	409	LEU	3.8
1	A	338	SER	3.7
1	A	527	SER	3.7
1	B	359	ASN	3.6
1	A	331	TYR	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	459	TYR	3.6
1	A	333	PRO	3.6
1	B	330	GLU	3.5
1	A	354	LEU	3.5
1	A	417	SER	3.4
1	A	420	GLY	3.4
1	A	542	GLU	3.4
1	A	463	SER	3.3
1	A	339	GLU	3.3
1	A	539	LEU	3.3
1	B	397	GLU	3.2
1	A	528	MET	3.2
1	B	422	VAL	3.1
1	B	460	THR	3.1
1	B	495	LEU	3.0
1	B	341	SER	3.0
1	B	525	LEU	3.0
1	A	466	LEU	2.9
1	B	342	MET	2.9
1	A	501	HIS	2.8
1	B	373	HIS	2.8
1	A	332	ASP	2.8
1	B	348	ASN	2.8
1	A	325	PRO	2.8
1	B	469	LEU	2.7
1	A	547	HIS	2.7
1	A	546	ALA	2.7
1	A	464	SER	2.6
1	B	320	LEU	2.6
1	A	492	LYS	2.6
1	B	343	MET	2.6
1	B	493	ALA	2.5
1	B	496	THR	2.5
1	A	328	TYR	2.5
1	A	465	THR	2.5
1	A	320	LEU	2.4
1	A	372	LEU	2.4
1	A	373	HIS	2.4
1	A	544	LEU	2.4
1	A	460	THR	2.4
1	B	351	ASP	2.4
1	A	439	ASN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	492	LYS	2.4
1	B	372	LEU	2.3
1	B	425	PHE	2.3
1	B	419	GLU	2.3
1	B	420	GLY	2.3
1	B	465	THR	2.3
1	B	491	ALA	2.3
1	A	497	LEU	2.3
1	A	437	MET	2.3
1	B	322	ALA	2.2
1	A	425	PHE	2.2
1	B	380	GLU	2.2
1	A	318	ALA	2.2
1	A	421	MET	2.2
1	A	541	LEU	2.2
1	A	371	THR	2.1
1	B	311	THR	2.1
1	A	491	ALA	2.1
1	B	417	SER	2.1
1	B	458	VAL	2.1
1	B	513	HIS	2.1
1	B	464	SER	2.1
1	A	419	GLU	2.1
1	A	493	ALA	2.1
1	B	358	ILE	2.0
1	A	362	LYS	2.0
1	B	326	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	DC8	B	51	23/23	0.91	0.10	37,47,51,53	0
2	DC8	A	50	23/23	0.92	0.09	32,45,49,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around DC8 A 50:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.