



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:10 AM UTC

PDB ID : 2Q72 / pdb_00002q72
Title : Crystal Structure Analysis of LeuT complexed with L-leucine, sodium, and imipramine
Authors : Singh, S.K.; Yamashita, A.; Gouaux, E.
Deposited on : 2007-06-05
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	519	14% 74% 23% ..

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4383 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Transporter.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	511	4084	2767	644	661	12	0	4	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	514	GLY	-	cloning artifact	UNP O67854
A	515	THR	-	cloning artifact	UNP O67854
A	516	LEU	-	cloning artifact	UNP O67854
A	517	VAL	-	cloning artifact	UNP O67854
A	518	PRO	-	cloning artifact	UNP O67854
A	519	ARG	-	cloning artifact	UNP O67854

- Molecule 2 is octyl beta-D-glucopyranoside (CCD ID: BOG) (formula: $C_{14}H_{28}O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			20	14	6		
2	A	1	Total	C	O	0	0
			20	14	6		
2	A	1	Total	C	O	0	0
			20	14	6		
2	A	1	Total	C	O	0	0
			20	14	6		
2	A	1	Total	C	O	0	0
			20	14	6		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Na	0	0
			2	2		

- Molecule 4 is LEUCINE (CCD ID: LEU) (formula: C₆H₁₃NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			9	6	1	2		

- Molecule 5 is 3-(5H-DIBENZO[B,F]AZEPIN-5-YL)-N,N-DIMETHYLPROPAN-1-AMINE (CCD ID: IXX) (formula: C₁₉H₂₄N₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	N	0	0
			21	19	2		
5	A	1	Total	C	N	0	0
			21	19	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	146	Total	O	0	0
			146	146		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Transporter



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	88.00Å 86.64Å 81.13Å 90.00° 95.76° 90.00°	Depositor
Resolution (Å)	27.43 – 1.70 27.43 – 1.70	Depositor EDS
% Data completeness (in resolution range)	95.4 (27.43-1.70) 95.5 (27.43-1.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 1.70Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.203 , 0.219 0.205 , 0.221	Depositor DCC
R_{free} test set	3293 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	27.4	Xtriage
Anisotropy	0.438	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 58.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4383	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, IXX, BOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/4211	0.90	16/5735 (0.3%)

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	131	PRO	N-CA-C	9.38	122.15	110.70
1	A	515	THR	N-CA-C	-7.55	103.96	113.02
1	A	187	ILE	N-CA-C	6.74	116.76	110.42
1	A	298	SER	N-CA-C	6.63	118.59	111.36
1	A	273	GLN	N-CA-C	6.05	118.77	110.24
1	A	44	MET	N-CA-C	5.92	117.73	111.28
1	A	21	ASN	N-CA-C	-5.84	104.82	111.07
1	A	455	ILE	N-CA-C	5.75	115.88	110.30
1	A	491	ILE	N-CA-C	-5.65	105.01	110.72
1	A	308	VAL	N-CA-C	5.55	115.75	110.42
1	A	77	THR	N-CA-C	5.51	121.17	113.57
1	A	136	ASP	CA-C-N	5.51	126.72	119.84
1	A	136	ASP	C-N-CA	5.51	126.72	119.84
1	A	198	ALA	N-CA-C	5.30	116.87	111.14
1	A	130	PRO	CA-C-N	-5.08	115.15	120.38
1	A	130	PRO	C-N-CA	-5.08	115.15	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4084	0	4177	77	0
2	A	100	0	140	7	0
3	A	2	0	0	0	0
4	A	9	0	10	0	0
5	A	42	0	48	2	0
6	A	146	0	0	2	0
All	All	4383	0	4375	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:396:LEU:HB2	2:A:705:BOG:H8'3	1.62	0.81
1:A:217:GLU:HG3	1:A:222:THR:HG22	1.72	0.72
1:A:45:ILE:HB	1:A:46:PRO:HD3	1.71	0.71
1:A:237:LYS:HG2	1:A:243:VAL:HG21	1.73	0.70
1:A:468:ALA:HA	1:A:472:ILE:HD12	1.74	0.68
1:A:299[A]:ILE:HB	1:A:300:PRO:HD3	1.75	0.67
1:A:33:VAL:O	1:A:37:GLU:HB2	2.01	0.61
2:A:703:BOG:H2'2	2:A:704:BOG:H1'2	1.83	0.61
1:A:128:PRO:O	1:A:130:PRO:HD3	2.00	0.60
1:A:34:GLN:HE21	5:A:801:IXX:H3	1.67	0.58
1:A:142:ARG:O	1:A:146:GLU:HG3	2.04	0.57
1:A:77:THR:OG1	1:A:97:GLY:HA3	2.05	0.56
1:A:124:VAL:HG23	1:A:126:LEU:HG	1.87	0.55
1:A:127:VAL:HB	1:A:128:PRO:HD2	1.86	0.55
1:A:155:PRO:HD3	1:A:162:LEU:HD23	1.88	0.55
1:A:512:SER:C	1:A:514:GLY:H	2.15	0.55
1:A:420:LEU:C	1:A:420:LEU:HD23	2.33	0.54
1:A:496:PHE:CZ	1:A:500:LEU:HD11	2.41	0.54
1:A:370:GLU:O	1:A:507:ARG:NE	2.41	0.54
1:A:31:PHE:HB3	1:A:32:PRO:HD3	1.88	0.54
1:A:166:LEU:O	1:A:170[B]:ILE:HG12	2.08	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:ALA:HB2	1:A:234:ASP:HB3	1.90	0.52
1:A:398:LYS:HA	6:A:977:HOH:O	2.10	0.52
1:A:409:THR:HG22	1:A:467:TRP:CZ2	2.45	0.52
1:A:299[B]:ILE:HB	1:A:300:PRO:HD3	1.91	0.51
1:A:479:THR:HB	1:A:483:VAL:HG11	1.93	0.50
1:A:415:PHE:O	1:A:419:GLU:HG2	2.11	0.50
1:A:131:PRO:HG2	1:A:139:SER:OG	2.12	0.50
1:A:480:HIS:HD2	1:A:482:THR:OG1	1.95	0.49
1:A:130:PRO:HG3	1:A:144:PHE:CE2	2.47	0.49
1:A:34:GLN:NE2	5:A:801:IXX:H3	2.27	0.49
1:A:212:ARG:HH11	1:A:212:ARG:HA	1.78	0.48
1:A:392:LEU:HD13	2:A:704:BOG:H8'1	1.95	0.48
1:A:472:ILE:N	1:A:473:PRO:CD	2.76	0.48
1:A:310:ASN:O	1:A:314:ILE:HG13	2.14	0.48
1:A:167:PHE:HB2	2:A:702:BOG:H1'1	1.95	0.47
1:A:287:GLU:O	1:A:291:VAL:HG22	2.14	0.47
1:A:130:PRO:CG	1:A:144:PHE:HE2	2.28	0.47
1:A:298:SER:HB2	1:A:322:LEU:HG	1.96	0.47
1:A:299[A]:ILE:HG12	6:A:928:HOH:O	2.14	0.47
1:A:131:PRO:HD2	1:A:139:SER:O	2.14	0.47
1:A:400:LEU:HD23	1:A:400:LEU:C	2.40	0.47
1:A:11:ARG:O	1:A:15:ILE:HG12	2.16	0.46
1:A:45:ILE:CG2	1:A:238:LEU:HD11	2.45	0.46
1:A:505:GLU:HB2	1:A:506:ARG:NH1	2.31	0.46
1:A:104:VAL:HA	1:A:107:TYR:CE2	2.51	0.46
1:A:130:PRO:HG3	1:A:144:PHE:HE2	1.80	0.46
1:A:448:TYR:O	1:A:452:MET:HG3	2.16	0.46
1:A:37:GLU:HG3	1:A:316:LYS:HG2	1.98	0.45
1:A:512:SER:C	1:A:514:GLY:N	2.75	0.45
1:A:136:ASP:HA	1:A:137:PRO:HD2	1.75	0.45
1:A:458:ALA:O	1:A:462:VAL:HG23	2.16	0.45
1:A:331:PHE:O	1:A:337:GLY:HA3	2.17	0.44
1:A:409:THR:HG22	1:A:467:TRP:CH2	2.53	0.44
1:A:421:ILE:O	1:A:425:TRP:HB2	2.17	0.44
2:A:704:BOG:H6'2	2:A:705:BOG:H7'1	2.00	0.44
1:A:289:ALA:O	1:A:293:LEU:HB2	2.18	0.44
1:A:468:ALA:CA	1:A:472:ILE:HD12	2.46	0.43
1:A:461:ALA:O	1:A:465:VAL:HG23	2.19	0.43
1:A:131:PRO:CG	1:A:139:SER:OG	2.67	0.43
1:A:280:LEU:O	1:A:284:THR:HG23	2.19	0.43
1:A:110:TYR:CZ	1:A:394:MET:HG2	2.54	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212:ARG:HH11	1:A:212:ARG:CA	2.31	0.42
2:A:703:BOG:O6	2:A:704:BOG:H62	2.19	0.42
1:A:109:VAL:HG11	1:A:176:MET:SD	2.60	0.42
1:A:24:GLY:HA2	1:A:351:ALA:HB1	2.02	0.42
1:A:66:GLY:O	1:A:267:SER:HA	2.20	0.42
1:A:234:ASP:C	1:A:236:GLU:H	2.28	0.42
1:A:505:GLU:CB	1:A:506:ARG:NH1	2.83	0.42
1:A:238:LEU:HD12	1:A:238:LEU:HA	1.92	0.41
1:A:199:MET:HB2	1:A:200:PRO:HD3	2.02	0.41
1:A:41:GLY:HA3	1:A:232:THR:O	2.20	0.41
1:A:259:PHE:CZ	1:A:412:VAL:HG11	2.55	0.41
1:A:423:PHE:O	1:A:427:PHE:HB3	2.20	0.41
1:A:391:HIS:HA	1:A:394:MET:HE3	2.03	0.41
1:A:145:LYS:HG3	1:A:325:ILE:HD13	2.02	0.40
1:A:392:LEU:CD1	2:A:704:BOG:H8'1	2.51	0.40
1:A:420:LEU:HD23	1:A:420:LEU:O	2.21	0.40
1:A:58:LEU:HD23	1:A:58:LEU:HA	1.94	0.40
1:A:45:ILE:HB	1:A:238:LEU:HD11	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	511/519 (98%)	496 (97%)	14 (3%)	1 (0%)	43 28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	235	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	413/420 (98%)	399 (97%)	14 (3%)	32	16

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	135	THR
1	A	138	ASP
1	A	139	SER
1	A	238	LEU
1	A	274	ASP
1	A	340	LEU
1	A	424	PHE
1	A	434	GLU
1	A	445	PRO
1	A	470	GLU
1	A	477	GLU
1	A	495[A]	LEU
1	A	495[B]	LEU
1	A	510	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	GLN
1	A	72	GLN
1	A	179	ASN
1	A	273	GLN
1	A	333	GLN
1	A	361	GLN
1	A	480	HIS
1	A	509	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BOG	A	701	-	20,20,20	1.10	2 (10%)	25,25,25	0.62	0
5	IXX	A	801	-	23,23,23	1.88	5 (21%)	31,31,31	1.01	1 (3%)
2	BOG	A	705	-	20,20,20	1.22	3 (15%)	25,25,25	0.83	1 (4%)
2	BOG	A	704	-	20,20,20	1.07	1 (5%)	25,25,25	0.70	0
2	BOG	A	702	-	20,20,20	1.22	2 (10%)	25,25,25	1.04	1 (4%)
2	BOG	A	703	-	20,20,20	1.07	2 (10%)	25,25,25	0.64	0
4	LEU	A	601	3	6,8,8	0.83	0	5,10,10	0.34	0
5	IXX	A	802	-	23,23,23	1.81	5 (21%)	31,31,31	0.93	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BOG	A	701	-	-	2/11/31/31	0/1/1/1
5	IXX	A	801	-	-	1/6/19/19	0/3/3/3
2	BOG	A	705	-	-	3/11/31/31	0/1/1/1
2	BOG	A	704	-	-	4/11/31/31	0/1/1/1
2	BOG	A	702	-	-	5/11/31/31	0/1/1/1
2	BOG	A	703	-	-	0/11/31/31	0/1/1/1
4	LEU	A	601	3	-	1/8/8/8	-
5	IXX	A	802	-	-	4/6/19/19	0/3/3/3

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	801	IXX	C11-C12	6.22	1.49	1.40
5	A	802	IXX	C11-C12	6.06	1.49	1.40
2	A	702	BOG	O5-C1	3.19	1.50	1.41
2	A	703	BOG	O5-C1	3.01	1.49	1.41
2	A	705	BOG	O5-C1	2.95	1.49	1.41
2	A	701	BOG	O5-C1	2.89	1.49	1.41
2	A	704	BOG	O5-C1	2.83	1.49	1.41
5	A	801	IXX	C1-C14	2.68	1.43	1.39
2	A	705	BOG	O1-C1	2.64	1.44	1.40
2	A	702	BOG	O1-C1	2.58	1.44	1.40
5	A	801	IXX	C10-C11	2.40	1.43	1.39
5	A	802	IXX	C2-C1	2.38	1.43	1.38
5	A	802	IXX	C6-C12	2.38	1.56	1.51
5	A	801	IXX	C6-C12	2.30	1.56	1.51
5	A	802	IXX	C1-C14	2.30	1.43	1.39
5	A	801	IXX	C2-C1	2.25	1.42	1.38
2	A	701	BOG	C4-C5	2.17	1.57	1.53
5	A	802	IXX	C10-C11	2.17	1.43	1.39
2	A	705	BOG	C4-C5	2.06	1.57	1.53
2	A	703	BOG	C4-C5	2.02	1.57	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	702	BOG	C1'-O1-C1	3.44	119.56	113.68
5	A	801	IXX	C15-N1-C14	2.59	121.61	117.10
5	A	802	IXX	C10-C11-N1	-2.22	117.52	120.84
2	A	705	BOG	C3'-C2'-C1'	-2.18	103.98	113.47
5	A	802	IXX	C12-C11-N1	2.05	124.22	120.06

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	702	BOG	C2'-C1'-O1-C1
2	A	702	BOG	O5-C5-C6-O6
5	A	802	IXX	C16-C17-N2-C18
5	A	802	IXX	C16-C17-N2-C19
2	A	702	BOG	C4-C5-C6-O6
2	A	705	BOG	O5-C5-C6-O6
2	A	705	BOG	C4-C5-C6-O6
2	A	701	BOG	O1-C1'-C2'-C3'
2	A	704	BOG	O5-C5-C6-O6
2	A	704	BOG	C4-C5-C6-O6
2	A	702	BOG	O1-C1'-C2'-C3'
5	A	801	IXX	C16-C15-N1-C14
2	A	702	BOG	C3'-C4'-C5'-C6'
2	A	701	BOG	C3'-C4'-C5'-C6'
2	A	705	BOG	C2'-C1'-O1-C1
5	A	802	IXX	C15-C16-C17-N2
2	A	704	BOG	O1-C1'-C2'-C3'
2	A	704	BOG	C3'-C4'-C5'-C6'
4	A	601	LEU	O-C-CA-N
5	A	802	IXX	N1-C15-C16-C17


There are no ring outliers.

5 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	801	IXX	2	0
2	A	705	BOG	2	0
2	A	704	BOG	5	0
2	A	702	BOG	1	0
2	A	703	BOG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and


any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand IXX A 801




Bond lengths



Bond angles




Torsions




Rings


Ligand BOG A 705




Bond lengths






Bond angles




Torsions



Rings





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	511/519 (98%)	0.83	75 (14%) 6 5	14, 32, 63, 76	4 (0%)

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	136	ASP	6.2
1	A	510	HIS	5.4
1	A	475	ILE	4.4
1	A	238	LEU	4.4
1	A	513	ALA	4.3
1	A	89	PHE	4.0
1	A	496	PHE	3.9
1	A	516	LEU	3.9
1	A	39	GLY	3.8
1	A	272	ASP	3.8
1	A	132	PRO	3.7
1	A	515	THR	3.7
1	A	466	VAL	3.5
1	A	135	THR	3.5
1	A	507	ARG	3.5
1	A	82	TYR	3.3
1	A	471	TYR	3.3
1	A	495[A]	LEU	3.3
1	A	140	ILE	3.3
1	A	476	MET	3.2
1	A	467	TRP	3.2
1	A	203	PHE	3.1
1	A	514	GLY	3.1
1	A	479	THR	3.1
1	A	425	TRP	3.0
1	A	131	PRO	3.0
1	A	473	PRO	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	462	VAL	3.0
1	A	511	GLU	2.9
1	A	459	PHE	2.8
1	A	129	GLU	2.8
1	A	137	PRO	2.8
1	A	449	TYR	2.8
1	A	214	PHE	2.7
1	A	235	PHE	2.6
1	A	420	LEU	2.6
1	A	159	GLU	2.6
1	A	156	LYS	2.5
1	A	472	ILE	2.5
1	A	512	SER	2.5
1	A	234	ASP	2.5
1	A	314	ILE	2.5
1	A	4	LYS	2.4
1	A	508	ARG	2.4
1	A	477	GLU	2.4
1	A	463	LEU	2.4
1	A	318	GLY	2.4
1	A	233	PRO	2.4
1	A	197	ILE	2.4
1	A	240	ASP	2.3
1	A	414	PHE	2.3
1	A	215	LEU	2.3
1	A	127	VAL	2.3
1	A	303	VAL	2.3
1	A	223	ALA	2.3
1	A	474	LYS	2.3
1	A	130	PRO	2.2
1	A	458	ALA	2.2
1	A	468	ALA	2.2
1	A	446	ARG	2.2
1	A	421	ILE	2.2
1	A	139	SER	2.2
1	A	36	ALA	2.1
1	A	71	ALA	2.1
1	A	319	ALA	2.1
1	A	158	ASP	2.1
1	A	441	ILE	2.1
1	A	42	ALA	2.1
1	A	509	ASN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	88	ARG	2.1
1	A	45	ILE	2.1
1	A	304	ALA	2.0
1	A	271	LYS	2.0
1	A	157	GLY	2.0
1	A	465	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BOG	A	702	20/20	0.57	0.20	62,73,75,75	0
2	BOG	A	704	20/20	0.66	0.22	59,73,76,76	0
2	BOG	A	705	20/20	0.67	0.22	63,71,72,73	0
2	BOG	A	703	20/20	0.70	0.17	41,49,62,62	0
5	IXX	A	802	21/21	0.73	0.23	78,79,83,83	0
2	BOG	A	701	20/20	0.79	0.17	53,57,60,60	0
5	IXX	A	801	21/21	0.94	0.07	25,27,34,35	0
4	LEU	A	601	9/9	0.97	0.06	20,23,24,24	0
3	NA	A	752	1/1	1.00	0.04	22,22,22,22	0
3	NA	A	751	1/1	1.00	0.03	23,23,23,23	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around BOG A 702:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BOG A 704:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BOG A 705:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BOG A 703:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around IXX A 802:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BOG A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around IXX A 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.