



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 11:59 AM UTC

PDB ID : 3Q7L / pdb_00003q7l
Title : Crystal structure of the E2 domain of amyloid precursor-like protein 1
Authors : Xue, Y.; Ha, Y.
Deposited on : 2011-01-05
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	9% 71% 17% •• 9%
1	B	214	2% 72% 19% 8%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3248 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Amyloid-like protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	195	Total	C	N	O	S	0	0	0
			1465	913	278	268	6			
1	B	196	Total	C	N	O	S	0	0	0
			1490	931	283	271	5			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	281	GLY	-	expression tag	UNP P51693
A	282	SER	-	expression tag	UNP P51693
A	283	HIS	-	expression tag	UNP P51693
A	284	MET	-	expression tag	UNP P51693
B	281	GLY	-	expression tag	UNP P51693
B	282	SER	-	expression tag	UNP P51693
B	283	HIS	-	expression tag	UNP P51693
B	284	MET	-	expression tag	UNP P51693

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	121	Total	O	0	0
			121	121		
3	B	162	Total	O	0	0
			162	162		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Amyloid-like protein 1



• Molecule 1: Amyloid-like protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.95Å 83.05Å 89.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.31 – 2.20 39.31 – 2.20	Depositor EDS
% Data completeness (in resolution range)	97.4 (39.31-2.20) 97.3 (39.31-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.14 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.216 , 0.256 0.229 , 0.268	Depositor DCC
R_{free} test set	2712 reflections (9.63%)	wwPDB-VP
Wilson B-factor (Å ²)	39.2	Xtriage
Anisotropy	0.086	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 50.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3248	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/1489	0.92	6/2023 (0.3%)
1	B	0.46	0/1514	0.87	0/2055
All	All	0.45	0/3003	0.90	6/4078 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	402	ASP	CA-C-N	-7.69	112.46	120.38
1	A	402	ASP	C-N-CA	-7.69	112.46	120.38
1	A	438	ASP	CA-C-N	7.51	127.05	119.24
1	A	438	ASP	C-N-CA	7.51	127.05	119.24
1	A	403	PRO	N-CA-C	-5.30	104.23	110.70
1	A	342	LEU	N-CA-C	5.03	115.91	110.13

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1465	0	1363	36	0
1	B	1490	0	1411	36	0
2	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	5	0	0	0	0
3	A	121	0	0	9	0
3	B	162	0	0	6	0
All	All	3248	0	2774	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:335:ALA:HB1	1:A:350:LEU:HD11	1.57	0.84
1:A:402:ASP:CB	1:A:403:PRO:HD3	2.09	0.82
1:A:379:ARG:HD2	3:A:75:HOH:O	1.81	0.80
1:A:396:LEU:HD12	1:A:466:LEU:HD13	1.68	0.76
1:B:402:ASP:CB	1:B:403:PRO:HD3	2.15	0.75
1:B:443:GLN:HG3	1:B:446:ARG:NH2	2.02	0.74
1:A:359:GLN:HE22	1:B:443:GLN:NE2	1.94	0.65
1:A:376:HIS:HA	3:A:75:HOH:O	1.96	0.64
1:A:402:ASP:O	1:A:404:PRO:HD3	1.98	0.64
1:B:317:LEU:HD13	1:B:368:GLU:HG2	1.83	0.60
1:B:414:ARG:HH12	1:B:418:ARG:HD3	1.64	0.60
1:B:418:ARG:HD3	3:B:78:HOH:O	2.01	0.60
1:B:388:ARG:HD3	3:B:25:HOH:O	2.02	0.58
1:A:402:ASP:CB	1:A:403:PRO:CD	2.83	0.56
1:B:381:ILE:HD11	3:B:276:HOH:O	2.06	0.56
1:A:315:MET:HE1	1:A:318:GLU:OE2	2.05	0.56
1:A:399:LEU:HD22	3:A:236:HOH:O	2.06	0.55
1:B:464:LEU:HD11	1:B:485:LEU:HD11	1.87	0.55
1:A:342:LEU:HD23	1:A:343:PRO:HD2	1.88	0.55
1:B:377:ALA:O	1:B:381:ILE:HG13	2.06	0.55
1:A:478:ARG:N	1:A:479:PRO:HD2	2.23	0.54
1:A:406:ALA:HB1	1:A:477:LEU:HD21	1.90	0.54
1:A:468:ASP:C	1:A:470:ASN:H	2.15	0.54
1:A:464:LEU:O	1:A:467:LEU:HB2	2.08	0.54
1:B:402:ASP:CB	1:B:403:PRO:CD	2.86	0.54
1:B:401:ALA:O	1:B:402:ASP:C	2.50	0.52
1:A:383:LEU:O	1:A:387:GLN:HG3	2.09	0.52
1:A:381:ILE:HG13	3:A:14:HOH:O	2.10	0.52
1:B:396:LEU:O	1:B:400:GLN:HG3	2.10	0.52
1:B:317:LEU:HD13	1:B:368:GLU:CG	2.40	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:294:VAL:HB	3:B:228:HOH:O	2.11	0.50
1:A:335:ALA:HB1	1:A:350:LEU:CD1	2.35	0.50
1:A:467:LEU:HA	3:A:236:HOH:O	2.12	0.50
1:A:375:THR:HG22	3:A:75:HOH:O	2.11	0.49
1:A:356:SER:HB3	3:A:231:HOH:O	2.13	0.48
1:B:441:LYS:HA	1:B:444:GLN:OE1	2.12	0.48
1:B:414:ARG:NH1	1:B:418:ARG:HD3	2.27	0.48
1:A:401:ALA:O	1:A:402:ASP:C	2.56	0.48
1:A:396:LEU:HD23	1:A:400:GLN:NE2	2.28	0.48
1:A:339:SER:HA	1:A:342:LEU:HD12	1.96	0.48
1:B:369:ARG:O	1:B:373:VAL:HG23	2.15	0.47
1:B:401:ALA:O	1:B:404:PRO:HD3	2.15	0.46
1:A:351:ASN:HD21	1:A:355:GLN:HE21	1.63	0.46
1:B:318:GLU:O	1:B:322:MET:HG3	2.14	0.46
1:B:414:ARG:HG2	1:B:414:ARG:HH11	1.81	0.46
1:B:383:LEU:O	1:B:387:GLN:HG3	2.16	0.46
1:A:359:GLN:HE22	1:B:443:GLN:HE21	1.65	0.45
1:A:351:ASN:HB2	3:A:62:HOH:O	2.17	0.45
1:B:414:ARG:NH1	1:B:418:ARG:CZ	2.80	0.44
1:B:320:ARG:HG2	1:B:324:GLN:HE21	1.82	0.44
1:A:484:LEU:O	1:A:486:HIS:N	2.51	0.44
1:B:478:ARG:N	1:B:479:PRO:HD2	2.32	0.43
1:A:350:LEU:HA	3:A:156:HOH:O	2.18	0.43
1:B:396:LEU:HD12	1:B:396:LEU:HA	1.90	0.43
1:A:319:GLU:HG2	3:B:229:HOH:O	2.18	0.43
1:A:484:LEU:O	1:A:485:LEU:C	2.62	0.42
1:B:396:LEU:CD1	1:B:466:LEU:HD13	2.48	0.42
1:A:359:GLN:NE2	1:B:443:GLN:HE21	2.18	0.42
1:A:431:TYR:CD2	1:B:325:ILE:HD13	2.55	0.42
1:A:315:MET:HE2	1:A:319:GLU:HG3	2.02	0.41
1:B:414:ARG:NH1	1:B:418:ARG:NE	2.68	0.41
1:B:406:ALA:HB1	1:B:477:LEU:HD21	2.02	0.41
1:B:462:GLN:HG3	3:B:89:HOH:O	2.20	0.41
1:A:315:MET:HE3	1:A:318:GLU:HB3	2.03	0.41
1:B:343:PRO:HG2	1:B:346:ASP:CG	2.45	0.41
1:B:414:ARG:NH1	1:B:418:ARG:CD	2.84	0.41
1:A:406:ALA:CB	1:A:477:LEU:HD21	2.51	0.40
1:B:310:PHE:CD1	1:B:376:HIS:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	193/214 (90%)	181 (94%)	8 (4%)	4 (2%)	5	3
1	B	194/214 (91%)	189 (97%)	4 (2%)	1 (0%)	24	27
All	All	387/428 (90%)	370 (96%)	12 (3%)	5 (1%)	9	8

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	403	PRO
1	A	485	LEU
1	A	469	GLN
1	A	402	ASP
1	A	403	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	136/186 (73%)	128 (94%)	8 (6%)	18	22
1	B	141/186 (76%)	139 (99%)	2 (1%)	59	75
All	All	277/372 (74%)	267 (96%)	10 (4%)	31	42

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	342	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	350	LEU
1	A	352	GLU
1	A	396	LEU
1	A	432	GLN
1	A	443	GLN
1	A	458	GLU
1	A	484	LEU
1	B	361	LEU
1	B	432	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	307	HIS
1	A	326	ASN
1	A	351	ASN
1	A	359	GLN
1	A	400	GLN
1	A	424	GLN
1	A	432	GLN
1	B	326	ASN
1	B	337	ASN
1	B	351	ASN
1	B	353	HIS
1	B	405	GLN
1	B	432	GLN
1	B	443	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	2	-	4,4,4	0.34	0	6,6,6	0.08	0
2	SO4	A	1	-	4,4,4	0.34	0	6,6,6	0.07	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	195/214 (91%)	0.31	19 (9%) 13 10	20, 34, 81, 112	0
1	B	196/214 (91%)	0.05	4 (2%) 65 62	15, 31, 61, 85	0
All	All	391/428 (91%)	0.18	23 (5%) 28 25	15, 32, 75, 112	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	402	ASP	4.2
1	A	470	ASN	4.0
1	A	486	HIS	4.0
1	A	469	GLN	3.3
1	A	471	PRO	3.1
1	A	339	SER	3.0
1	A	477	LEU	2.9
1	B	403	PRO	2.6
1	B	341	ASN	2.6
1	A	485	LEU	2.6
1	A	480	GLN	2.5
1	A	483	GLU	2.4
1	A	468	ASP	2.4
1	A	404	PRO	2.4
1	A	481	ILE	2.3
1	A	437	VAL	2.3
1	A	439	PRO	2.2
1	A	474	ALA	2.2
1	A	484	LEU	2.2
1	A	467	LEU	2.1
1	B	381	ILE	2.1
1	A	473	LEU	2.1
1	B	402	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	A	1	5/5	0.94	0.10	56,65,73,77	0
2	SO4	B	2	5/5	0.95	0.09	51,51,55,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.