



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:17 PM UTC

PDB ID : 4Q7D / pdb_00004q7d
Title : Wild type Fc (wtFc)
Authors : Ahmed, A.A.; Giddens, J.; Pincetic, A.; Lomino, J.V.; Ravetch, J.V.; Wang, L.X.; Bjorkman, P.J.
Deposited on : 2014-04-24
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	22% 71% 20% 6%
1	B	221	2% 81% 12% 6%
2	C	8	38% 62%
2	D	8	25% 75%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BMA	C	3	-	-	X	-
2	MAN	C	6	-	-	X	-

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3685 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ig gamma-1 chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	S	11	0	0
			1660	1057	279	318	6			
1	B	208	Total	C	N	O	S	11	0	0
			1662	1058	280	318	6			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	8	Total	C	N	O	3	0	0
			99	56	4	39			
2	D	8	Total	C	N	O	0	0	0
			99	56	4	39			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	65	Total	O	0	0
			65	65		
3	B	100	Total	O	0	0
			100	100		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ig gamma-1 chain C region



- Molecule 1: Ig gamma-1 chain C region



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1	MAG2	BMA3	MAN4	NAG5	MAN6	MAG7	FUC8
------	------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.52Å 79.48Å 137.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.09 – 2.35 52.09 – 2.35	Depositor EDS
% Data completeness (in resolution range)	(Not available) (52.09-2.35) 99.7 (52.09-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 2.34Å)	Xtriage
Refinement program	PHENIX refine	Depositor
R, R_{free}	0.211 , 0.264 0.221 , 0.271	Depositor DCC
R_{free} test set	1991 reflections (8.49%)	wwPDB-VP
Wilson B-factor (Å ²)	44.6	Xtriage
Anisotropy	0.303	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 44.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3685	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, FUC, MAN, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.45	4/1706 (0.2%)	1.24	10/2324 (0.4%)
1	B	1.01	0/1708	1.04	2/2327 (0.1%)
All	All	1.25	4/3414 (0.1%)	1.15	12/4651 (0.3%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	297	ASN	C-N	-27.39	0.95	1.33
1	A	298	SER	C-N	-25.75	0.96	1.33
1	A	296	TYR	C-N	19.43	1.60	1.33
1	A	348	VAL	C-O	-5.77	1.17	1.24

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	296	TYR	O-C-N	14.64	139.35	122.22
1	A	297	ASN	O-C-N	-12.02	106.99	122.37
1	A	296	TYR	CA-C-N	-11.71	103.22	122.65
1	A	296	TYR	C-N-CA	-11.71	103.22	122.65
1	A	297	ASN	CA-C-N	10.60	141.78	121.54
1	A	297	ASN	C-N-CA	10.60	141.78	121.54
1	A	298	SER	O-C-N	6.43	131.14	122.59
1	A	260	THR	N-CA-C	6.27	118.91	108.13
1	A	342	GLN	CA-C-N	-5.76	113.84	119.78
1	A	342	GLN	C-N-CA	-5.76	113.84	119.78
1	B	290	LYS	CA-C-N	-5.04	114.80	120.14
1	B	290	LYS	C-N-CA	-5.04	114.80	120.14

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1660	0	1629	55	0
1	B	1662	0	1627	22	0
2	C	99	0	86	24	0
2	D	99	0	85	0	0
3	A	65	0	0	8	0
3	B	100	0	0	6	1
All	All	3685	0	3427	88	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:642:HOH:O	2:C:7:NAG:O3	1.61	1.17
1:A:353:PRO:HB2	1:A:358:LEU:CD2	1.77	1.15
1:A:353:PRO:HB2	1:A:358:LEU:HD21	1.13	1.09
1:A:260:THR:HG21	2:C:7:NAG:H61	1.38	1.05
1:A:297:ASN:HD22	2:C:1:NAG:C1	1.54	1.03
2:C:3:BMA:C6	2:C:6:MAN:C1	2.38	1.02
1:A:322:LYS:NZ	3:A:629:HOH:O	2.00	0.95
1:A:241:PHE:CE2	2:C:2:NAG:H4	2.05	0.91
1:A:297:ASN:HD21	2:C:1:NAG:C1	1.75	0.91
1:A:353:PRO:CB	1:A:358:LEU:HD21	2.02	0.90
1:A:301:ARG:NH1	2:C:2:NAG:O7	2.06	0.89
1:B:352:PRO:O	3:B:614:HOH:O	1.88	0.88
1:B:355:ARG:NH1	3:B:681:HOH:O	2.06	0.88
3:A:642:HOH:O	2:C:7:NAG:C4	2.22	0.86
1:A:241:PHE:HE2	2:C:2:NAG:H4	1.44	0.83
2:C:3:BMA:O6	2:C:6:MAN:C5	2.26	0.83
1:A:353:PRO:CB	1:A:358:LEU:CD2	2.55	0.82
2:C:3:BMA:O6	2:C:6:MAN:H5	1.82	0.79
1:B:353:PRO:HB2	1:B:358:LEU:CD2	2.14	0.77
2:C:3:BMA:C6	2:C:6:MAN:O5	2.34	0.76
1:B:347:GLN:OE1	3:B:633:HOH:O	2.05	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:3:BMA:O6	2:C:6:MAN:O5	2.06	0.71
1:A:260:THR:CG2	2:C:7:NAG:H61	2.19	0.71
2:C:3:BMA:H62	2:C:6:MAN:C1	2.24	0.68
1:A:370:LYS:NZ	3:A:640:HOH:O	2.28	0.66
1:A:377:ILE:O	3:A:661:HOH:O	2.13	0.65
1:A:353:PRO:CB	1:A:358:LEU:HD23	2.27	0.65
1:A:241:PHE:CE2	2:C:2:NAG:C4	2.80	0.64
1:B:353:PRO:HB2	1:B:358:LEU:HD23	1.80	0.63
1:A:391:TYR:OH	3:A:655:HOH:O	2.14	0.62
1:A:243:PHE:CE1	2:C:6:MAN:C1	2.83	0.62
1:A:260:THR:HG21	2:C:7:NAG:C6	2.25	0.60
1:A:325:ASN:O	1:A:328:LEU:HG	2.02	0.60
1:A:422:VAL:HG22	1:A:442:SER:OG	2.05	0.57
1:A:332:ILE:CG2	1:A:333:GLU:N	2.68	0.57
1:B:353:PRO:HB2	1:B:358:LEU:HD21	1.86	0.56
1:A:330:ALA:HB1	1:A:331:PRO:HD2	1.87	0.56
2:C:3:BMA:C6	2:C:6:MAN:C5	2.85	0.55
1:A:243:PHE:CZ	2:C:6:MAN:C1	2.91	0.54
1:B:399:ASP:O	1:B:401:ASP:N	2.41	0.54
1:A:266:VAL:HB	1:A:300:TYR:HB2	1.90	0.53
1:A:260:THR:O	1:A:260:THR:HG23	2.10	0.51
1:A:282:VAL:HG23	1:A:283:GLU:O	2.11	0.51
1:A:262:VAL:CG1	1:A:301:ARG:NH1	2.74	0.51
1:A:347:GLN:HG3	1:A:349:TYR:CZ	2.47	0.50
1:B:443:LEU:O	3:B:664:HOH:O	2.18	0.50
1:A:330:ALA:HB1	1:A:331:PRO:CD	2.41	0.50
1:B:297:ASN:OD1	1:B:297:ASN:C	2.54	0.50
1:A:406:LEU:C	1:A:406:LEU:HD12	2.38	0.49
3:A:642:HOH:O	2:C:7:NAG:H4	2.01	0.48
1:A:332:ILE:HG22	1:A:333:GLU:N	2.29	0.48
1:B:353:PRO:HB3	1:B:363:VAL:CG1	2.44	0.48
1:B:411:THR:HG22	3:B:660:HOH:O	2.14	0.48
1:A:246:LYS:NZ	3:A:642:HOH:O	2.46	0.47
1:A:365:LEU:HB3	1:A:441:LEU:HD23	1.96	0.47
1:A:264:VAL:HG11	2:C:1:NAG:O4	2.15	0.47
1:A:263:VAL:HB	1:A:302:VAL:HG13	1.97	0.46
1:A:262:VAL:HG11	1:A:301:ARG:HH12	1.80	0.46
1:A:264:VAL:HG11	2:C:2:NAG:O5	2.16	0.46
1:A:383:SER:OG	1:A:422:VAL:O	2.24	0.45
1:B:266:VAL:HB	1:B:300:TYR:HB2	1.98	0.45
1:B:412:VAL:HG11	1:B:423:PHE:CE1	2.51	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:421:ASN:N	1:A:421:ASN:OD1	2.50	0.45
1:A:266:VAL:O	1:A:300:TYR:N	2.40	0.44
1:B:265:ASP:HA	1:B:299:THR:HB	1.99	0.44
1:B:411:THR:CG2	3:B:660:HOH:O	2.66	0.44
1:A:328:LEU:HD13	1:A:330:ALA:HA	2.00	0.43
1:B:353:PRO:CB	1:B:358:LEU:HD23	2.46	0.43
1:A:320:LYS:HD2	1:A:333:GLU:CD	2.44	0.43
1:B:248:LYS:HD2	1:B:428:MET:HE1	2.00	0.43
1:B:363:VAL:HG12	1:B:364:SER:N	2.34	0.42
2:C:3:BMA:C6	2:C:6:MAN:H5	2.48	0.42
1:A:295:GLN:HB3	1:A:297:ASN:OD1	2.20	0.42
1:A:325:ASN:H	1:A:328:LEU:HD23	1.84	0.42
1:B:278:TYR:CE2	1:B:283:GLU:HB2	2.54	0.42
1:A:365:LEU:HB3	1:A:441:LEU:CD2	2.49	0.42
1:B:433:HIS:O	1:B:434:ASN:HB2	2.19	0.41
1:A:273:VAL:HG21	1:A:302:VAL:HG11	2.02	0.41
1:B:414:LYS:HG2	1:B:418:GLN:NE2	2.35	0.41
1:A:262:VAL:CG1	1:A:301:ARG:HH12	2.33	0.41
1:A:262:VAL:HG11	1:A:301:ARG:NH1	2.35	0.41
1:A:415:SER:O	1:A:418:GLN:N	2.53	0.41
1:B:353:PRO:HG3	1:B:363:VAL:HG12	2.03	0.41
1:A:240:VAL:HG21	1:A:332:ILE:HG21	2.02	0.41
1:A:296:TYR:O	1:A:296:TYR:CG	2.72	0.41
1:A:297:ASN:O	1:A:298:SER:CB	2.67	0.40
1:A:270:ASP:N	1:A:271:PRO:HD3	2.36	0.40
1:A:268:HIS:ND1	1:A:300:TYR:HE1	2.19	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:669:HOH:O	3:B:680:HOH:O[4_555]	2.19	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/221 (93%)	190 (93%)	13 (6%)	2 (1%)	12	12
1	B	206/221 (93%)	200 (97%)	6 (3%)	0	100	100
All	All	411/442 (93%)	390 (95%)	19 (5%)	2 (0%)	24	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	298	SER
1	A	419	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/204 (95%)	191 (98%)	3 (2%)	57	72
1	B	193/204 (95%)	190 (98%)	3 (2%)	55	70
All	All	387/408 (95%)	381 (98%)	6 (2%)	55	70

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	282	VAL
1	A	360	LYS
1	A	442	SER
1	B	248	LYS
1	B	386	GLN
1	B	418	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	361	ASN
1	A	438	GLN
1	B	347	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	2,1	14,14,15	0.76	0	17,19,21	1.20	1 (5%)
2	NAG	C	2	2	14,14,15	0.76	0	17,19,21	1.00	1 (5%)
2	BMA	C	3	2	11,11,12	0.59	0	15,15,17	1.54	3 (20%)
2	MAN	C	4	2	11,11,12	0.59	0	15,15,17	1.35	1 (6%)
2	NAG	C	5	2	14,14,15	0.48	0	17,19,21	1.13	2 (11%)
2	MAN	C	6	2	11,11,12	0.74	0	15,15,17	1.39	1 (6%)
2	NAG	C	7	2	14,14,15	0.97	1 (7%)	17,19,21	1.23	2 (11%)
2	FUC	C	8	2	10,10,11	0.56	0	14,14,16	1.09	1 (7%)
2	NAG	D	1	2,1	14,14,15	0.56	0	17,19,21	1.73	4 (23%)
2	NAG	D	2	2	14,14,15	0.66	0	17,19,21	0.99	0
2	BMA	D	3	2	11,11,12	0.37	0	15,15,17	1.23	1 (6%)
2	MAN	D	4	2	11,11,12	0.63	0	15,15,17	1.10	2 (13%)
2	NAG	D	5	2	14,14,15	0.71	0	17,19,21	1.48	5 (29%)
2	MAN	D	6	2	11,11,12	0.41	0	15,15,17	1.31	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	D	7	2	14,14,15	0.69	0	17,19,21	0.98	1 (5%)
2	FUC	D	8	2	10,10,11	0.63	0	14,14,16	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	NAG	C	5	2	-	4/6/23/26	0/1/1/1
2	MAN	C	6	2	-	0/2/19/22	0/1/1/1
2	NAG	C	7	2	-	0/6/23/26	0/1/1/1
2	FUC	C	8	2	-	-	0/1/1/1
2	NAG	D	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1
2	BMA	D	3	2	-	0/2/19/22	0/1/1/1
2	MAN	D	4	2	-	2/2/19/22	0/1/1/1
2	NAG	D	5	2	-	4/6/23/26	0/1/1/1
2	MAN	D	6	2	-	0/2/19/22	0/1/1/1
2	NAG	D	7	2	-	2/6/23/26	0/1/1/1
2	FUC	D	8	2	-	-	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	7	NAG	O5-C5	-2.21	1.39	1.43

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	6	MAN	C1-O5-C5	4.66	118.44	112.19
2	D	1	NAG	C1-O5-C5	4.16	117.77	112.19
2	D	3	BMA	C1-O5-C5	3.87	117.37	112.19
2	C	4	MAN	C1-O5-C5	3.77	117.24	112.19
2	D	6	MAN	C1-O5-C5	3.68	117.12	112.19
2	C	3	BMA	O3-C3-C4	-3.27	102.66	110.38
2	C	5	NAG	C1-O5-C5	3.23	116.51	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	7	NAG	O4-C4-C3	3.17	117.84	110.38
2	D	1	NAG	C2-N2-C7	3.04	126.98	122.90
2	D	5	NAG	C4-C3-C2	3.00	115.41	111.02
2	C	3	BMA	C1-O5-C5	2.97	116.16	112.19
2	C	3	BMA	O6-C6-C5	-2.71	102.09	111.33
2	D	1	NAG	C8-C7-N2	2.67	120.55	116.12
2	D	4	MAN	C1-O5-C5	2.52	115.56	112.19
2	D	4	MAN	O4-C4-C5	2.45	115.36	109.32
2	D	1	NAG	C1-C2-N2	2.36	114.15	110.43
2	D	5	NAG	C2-N2-C7	2.35	126.05	122.90
2	D	5	NAG	C3-C4-C5	2.29	114.39	110.23
2	C	2	NAG	C8-C7-N2	-2.23	112.42	116.12
2	C	1	NAG	C1-O5-C5	2.15	115.07	112.19
2	D	5	NAG	O5-C1-C2	2.12	114.58	111.29
2	D	5	NAG	C8-C7-N2	2.10	119.61	116.12
2	C	8	FUC	O2-C2-C1	2.08	113.99	109.22
2	C	5	NAG	O5-C1-C2	2.06	114.48	111.29
2	C	7	NAG	O5-C5-C4	-2.06	105.82	110.83
2	D	7	NAG	C8-C7-N2	2.01	119.45	116.12

There are no chirality outliers.

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	5	NAG	C8-C7-N2-C2
2	C	5	NAG	O7-C7-N2-C2
2	D	1	NAG	C8-C7-N2-C2
2	D	1	NAG	O7-C7-N2-C2
2	D	5	NAG	C8-C7-N2-C2
2	D	5	NAG	O7-C7-N2-C2
2	D	7	NAG	C8-C7-N2-C2
2	D	7	NAG	O7-C7-N2-C2
2	D	5	NAG	O5-C5-C6-O6
2	D	5	NAG	C4-C5-C6-O6
2	D	4	MAN	C4-C5-C6-O6
2	D	4	MAN	O5-C5-C6-O6
2	C	5	NAG	C4-C5-C6-O6
2	C	5	NAG	O5-C5-C6-O6


There are no ring outliers.

5 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	7	NAG	6	0
2	C	3	BMA	8	0
2	C	2	NAG	5	0
2	C	6	MAN	10	0
2	C	1	NAG	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	298:SER	C	299:THR	N	0.96
1	A	297:ASN	C	298:SER	N	0.95

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	207/221 (93%)	1.03	49 (23%) 2 1	26, 51, 96, 122	4 (1%)
1	B	208/221 (94%)	0.22	5 (2%) 59 66	20, 43, 67, 81	4 (1%)
All	All	415/442 (93%)	0.62	54 (13%) 7 8	20, 45, 88, 122	8 (1%)

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	273	VAL	5.2
1	A	300	TYR	4.8
1	A	358	LEU	4.7
1	A	264	VAL	4.6
1	B	356	ASP	4.4
1	A	356	ASP	4.2
1	A	238	PRO	4.2
1	A	296	TYR	4.0
1	A	303	VAL	3.7
1	A	323	VAL	3.6
1	A	301	ARG	3.6
1	A	261	CYS	3.6
1	B	358	LEU	3.5
1	A	263	VAL	3.5
1	A	302	VAL	3.5
1	A	291	PRO	3.4
1	A	260	THR	3.4
1	A	324	SER	3.3
1	A	330	ALA	3.2
1	A	297	ASN	3.0
1	A	287	ALA	3.0
1	A	332	ILE	3.0
1	A	239	SER	3.0
1	A	298	SER	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	275	PHE	2.8
1	A	355	ARG	2.8
1	B	421	ASN	2.8
1	B	389	ASN	2.7
1	A	241	PHE	2.7
1	A	277	TRP	2.5
1	A	329	PRO	2.5
1	A	420	GLY	2.5
1	A	299	THR	2.5
1	A	292	ARG	2.5
1	A	266	VAL	2.4
1	A	359	THR	2.4
1	A	284	VAL	2.4
1	A	331	PRO	2.3
1	A	282	VAL	2.3
1	A	265	ASP	2.3
1	A	272	GLU	2.3
1	A	321	CYS	2.2
1	A	444	SER	2.2
1	A	240	VAL	2.2
1	B	236	GLY	2.2
1	A	262	VAL	2.2
1	A	333	GLU	2.2
1	A	281	GLY	2.2
1	A	295	GLN	2.2
1	A	267	SER	2.2
1	A	269	GLU	2.1
1	A	328	LEU	2.1
1	A	268	HIS	2.1
1	A	259	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FUC	C	8	10/11	0.36	0.22	47,49,56,56	3
2	NAG	C	2	14/15	0.44	0.29	37,40,44,46	14
2	FUC	D	8	10/11	0.48	0.36	99,103,108,109	0
2	NAG	C	1	14/15	0.53	0.27	41,42,45,47	14
2	NAG	C	5	14/15	0.60	0.17	76,81,85,86	0
2	BMA	C	3	11/12	0.61	0.17	40,42,43,45	0
2	MAN	C	6	11/12	0.63	0.16	44,46,51,55	0
2	NAG	C	7	14/15	0.64	0.20	37,43,50,52	0
2	NAG	D	5	14/15	0.67	0.17	82,90,92,96	0
2	NAG	D	1	14/15	0.79	0.24	74,88,93,96	0
2	MAN	D	4	11/12	0.80	0.14	52,59,63,74	0
2	NAG	D	7	14/15	0.81	0.13	43,47,56,56	0
2	MAN	C	4	11/12	0.86	0.10	50,53,59,68	0
2	MAN	D	6	11/12	0.89	0.12	48,54,57,58	0
2	NAG	D	2	14/15	0.90	0.17	56,63,67,67	0
2	BMA	D	3	11/12	0.93	0.10	51,52,53,57	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.