



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 09:21 AM UTC

PDB ID : 4Q7N / pdb_00004q7n
Title : Crystal structure of the complex of Buffalo Signalling protein SPB-40 with 4-N-trimethylaminobutyraldehyde at 1.79 Angstrom Resolution
Authors : Chaudhary, A.; Tyagi, T.K.; Singh, A.; Sinha, M.; Bhushan, A.; Kaur, P.; Sharma, S.; Singh, T.P.
Deposited on : 2014-04-25
Resolution : 1.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	361	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	2ZO	A	402	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3236 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chitinase-3-like protein 1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	361	Total	C	N	O	S	0	0	0
			2894	1851	507	527	9			

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is N,N,N-trimethyl-4-oxobutan-1-aminium (CCD ID: 2ZO) (formula: $C_7H_{16}NO$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			9	7	1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	319	Total	O	0	0
			319	319		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chitinase-3-like protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.07Å 66.73Å 106.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	56.49 – 1.79 56.49 – 1.79	Depositor EDS
% Data completeness (in resolution range)	98.7 (56.49-1.79) 98.6 (56.49-1.79)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.185 , 0.229 0.185 , 0.228	Depositor DCC
R_{free} test set	2058 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	23.6	Xtriage
Anisotropy	0.252	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3236	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, 2ZO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.45	9/2974 (0.3%)	1.45	27/4037 (0.7%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	353	SER	CB-OG	9.93	1.62	1.42
1	A	337	ASP	C-O	7.11	1.31	1.23
1	A	219	ALA	CA-CB	6.96	1.64	1.53
1	A	102	VAL	CA-C	6.54	1.59	1.52
1	A	184	THR	CA-CB	6.29	1.60	1.52
1	A	78	TRP	N-CA	5.95	1.53	1.46
1	A	241	GLY	N-CA	5.68	1.50	1.44
1	A	239	THR	CB-CG2	-5.41	1.34	1.52
1	A	138	ALA	C-O	5.30	1.30	1.24

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	107	ARG	NE-CZ-NH1	11.52	133.02	121.50
1	A	212	ARG	N-CA-C	-10.90	96.47	110.53
1	A	107	ARG	CB-CG-CD	10.78	136.10	111.30
1	A	289	ARG	NE-CZ-NH2	-10.31	109.92	119.20
1	A	107	ARG	CD-NE-CZ	10.13	138.58	124.40
1	A	75	VAL	CB-CA-C	-9.83	95.87	110.81
1	A	107	ARG	NE-CZ-NH2	-9.31	110.82	119.20
1	A	289	ARG	CG-CD-NE	-8.28	93.79	112.00
1	A	80	TYR	CA-C-N	-7.53	110.77	121.54
1	A	80	TYR	C-N-CA	-7.53	110.77	121.54
1	A	75	VAL	CA-CB-CG2	7.44	123.05	110.40
1	A	265	ARG	CG-CD-NE	-7.27	96.01	112.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	9	SER	CA-CB-OG	-6.86	97.38	111.10
1	A	81	GLY	N-CA-C	6.84	123.79	112.30
1	A	182	LEU	CB-CA-C	-6.72	98.78	109.80
1	A	107	ARG	CA-CB-CG	6.66	127.41	114.10
1	A	102	VAL	CA-C-N	-6.13	114.07	120.38
1	A	102	VAL	C-N-CA	-6.13	114.07	120.38
1	A	195	VAL	CG1-CB-CG2	5.93	123.85	110.80
1	A	52	ASP	CA-C-N	-5.68	112.68	120.46
1	A	52	ASP	C-N-CA	-5.68	112.68	120.46
1	A	127	ARG	CG-CD-NE	5.53	124.16	112.00
1	A	289	ARG	CD-NE-CZ	5.44	132.02	124.40
1	A	112	ASP	N-CA-C	5.11	119.50	113.16
1	A	182	LEU	CB-CG-CD2	5.08	125.93	110.70
1	A	289	ARG	CA-CB-CG	5.07	124.25	114.10
1	A	239	THR	N-CA-C	-5.07	107.26	113.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2894	0	2818	78	1
2	A	14	0	13	0	0
3	A	9	0	16	8	0
4	A	319	0	0	19	1
All	All	3236	0	2847	80	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:206:GLU:C	1:A:207:ASP:OD2	1.82	1.22

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:GLY:HA3	4:A:813:HOH:O	1.08	1.21
1:A:107:ARG:HD3	1:A:143:GLU:OE2	1.51	1.08
1:A:195:VAL:HG23	1:A:257:ILE:HD11	1.38	1.05
1:A:206:GLU:O	1:A:207:ASP:OD2	1.74	1.04
1:A:60:ASN:HB3	4:A:767:HOH:O	1.55	1.03
1:A:195:VAL:HG23	1:A:257:ILE:CD1	1.95	0.97
1:A:142:ARG:HD2	4:A:773:HOH:O	1.64	0.95
1:A:204:GLY:HA2	1:A:292:ASP:HB2	1.46	0.94
3:A:402:2ZO:H8	4:A:756:HOH:O	1.72	0.90
1:A:311:GLU:HG2	4:A:782:HOH:O	1.72	0.89
1:A:93:GLN:CD	1:A:96:ARG:HH22	1.81	0.88
1:A:250:LYS:HZ3	1:A:252:ASP:HB3	1.37	0.88
1:A:212:ARG:O	1:A:213:PHE:HB2	1.75	0.87
1:A:212:ARG:O	1:A:213:PHE:CB	2.30	0.79
1:A:207:ASP:HB2	4:A:618:HOH:O	1.83	0.77
1:A:204:GLY:CA	1:A:292:ASP:HB2	2.15	0.77
1:A:14:ARG:HH21	3:A:402:2ZO:H10	1.51	0.74
1:A:198:HIS:H	1:A:198:HIS:CD2	2.05	0.74
1:A:251:THR:HG23	4:A:700:HOH:O	1.86	0.73
1:A:93:GLN:NE2	1:A:96:ARG:HH22	1.91	0.68
1:A:225:ARG:O	1:A:225:ARG:NE	2.26	0.68
1:A:14:ARG:NH2	3:A:402:2ZO:H10	2.09	0.68
1:A:148:THR:HG23	4:A:819:HOH:O	1.93	0.67
1:A:10:TRP:HE1	3:A:402:2ZO:C7	2.08	0.66
1:A:57:ASP:O	1:A:61:THR:HG23	1.95	0.65
1:A:120:TRP:CZ3	1:A:158:THR:HB	2.31	0.65
1:A:338:PHE:O	1:A:348:THR:OG1	2.13	0.64
1:A:221:SER:OG	1:A:323:ARG:NH2	2.29	0.64
1:A:251:THR:CG2	4:A:700:HOH:O	2.44	0.63
1:A:206:GLU:C	1:A:207:ASP:CG	2.61	0.62
1:A:225:ARG:HA	1:A:225:ARG:HE	1.65	0.61
1:A:207:ASP:OD2	1:A:207:ASP:N	2.30	0.60
1:A:61:THR:HG22	4:A:727:HOH:O	2.03	0.59
1:A:250:LYS:NZ	1:A:252:ASP:HB3	2.16	0.58
1:A:225:ARG:HE	1:A:225:ARG:CA	2.17	0.58
1:A:323:ARG:HD2	4:A:816:HOH:O	2.04	0.57
1:A:188:HIS:HE1	1:A:194:THR:O	1.88	0.57
1:A:195:VAL:HG23	1:A:257:ILE:HD13	1.83	0.56
1:A:239:THR:HG22	1:A:332:ALA:H	1.71	0.56
1:A:239:THR:HG21	1:A:332:ALA:O	2.05	0.55
1:A:269:TRP:CE3	1:A:272:ILE:HG13	2.41	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:190:ALA:O	1:A:192:ARG:O	2.25	0.55
1:A:53:VAL:CG2	4:A:811:HOH:O	2.55	0.54
1:A:289:ARG:HD3	4:A:815:HOH:O	2.07	0.54
1:A:53:VAL:HG22	4:A:811:HOH:O	2.07	0.54
1:A:10:TRP:HE1	3:A:402:2ZO:H9	1.73	0.54
1:A:119:LEU:HD22	1:A:120:TRP:CZ3	2.43	0.54
1:A:195:VAL:CG2	1:A:257:ILE:HD11	2.26	0.54
1:A:348:THR:HG22	4:A:562:HOH:O	2.08	0.53
1:A:209:SER:N	4:A:618:HOH:O	2.34	0.53
1:A:269:TRP:HE3	1:A:272:ILE:HG13	1.73	0.53
1:A:172:GLN:OE1	1:A:175:ARG:NH1	2.41	0.52
1:A:262:ILE:H	1:A:303:GLN:HE22	1.59	0.51
1:A:320:LEU:HD12	1:A:320:LEU:C	2.37	0.50
1:A:205:ASN:O	1:A:206:GLU:C	2.55	0.49
1:A:239:THR:HG23	1:A:335:LEU:HB2	1.94	0.49
1:A:10:TRP:HE1	3:A:402:2ZO:H10	1.74	0.49
1:A:206:GLU:HG2	1:A:292:ASP:OD1	2.13	0.48
1:A:57:ASP:O	1:A:61:THR:CG2	2.62	0.47
1:A:204:GLY:HA2	1:A:292:ASP:CB	2.33	0.47
1:A:74:SER:HA	1:A:115:ASP:HB3	1.97	0.46
1:A:170:ILE:HD12	1:A:223:MET:HE1	1.97	0.45
1:A:212:ARG:O	1:A:213:PHE:CG	2.70	0.45
1:A:239:THR:HG22	1:A:332:ALA:N	2.31	0.45
1:A:78:TRP:CD1	1:A:119:LEU:HD12	2.52	0.44
1:A:204:GLY:HA3	1:A:292:ASP:CG	2.42	0.44
3:A:402:2ZO:H1	3:A:402:2ZO:H6	1.05	0.43
1:A:142:ARG:CD	4:A:773:HOH:O	2.44	0.42
1:A:321:LYS:CE	4:A:809:HOH:O	2.68	0.42
1:A:240:PHE:HB3	1:A:335:LEU:HD13	2.02	0.41
1:A:225:ARG:NE	1:A:225:ARG:C	2.79	0.41
1:A:83:GLN:HE21	1:A:83:GLN:H	1.68	0.41
1:A:9:SER:O	1:A:12:GLN:HG2	2.20	0.41
1:A:192:ARG:HD3	4:A:762:HOH:O	2.20	0.41
1:A:239:THR:CG2	1:A:332:ALA:H	2.32	0.40
1:A:10:TRP:NE1	3:A:402:2ZO:H9	2.36	0.40
1:A:78:TRP:CD1	1:A:119:LEU:CD1	3.05	0.40
1:A:244:TYR:HB3	1:A:257:ILE:CG1	2.52	0.40
1:A:204:GLY:CA	1:A:292:ASP:CB	2.94	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:357:ASP:OD2	4:A:615:HOH:O[4_445]	1.91	0.29

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	359/361 (99%)	346 (96%)	9 (2%)	4 (1%)	11 3

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	205	ASN
1	A	204	GLY
1	A	213	PHE
1	A	206	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	304/304 (100%)	274 (90%)	30 (10%)	7 2

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	SER
1	A	55	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	61	THR
1	A	75	VAL
1	A	78	TRP
1	A	83	GLN
1	A	85	PHE
1	A	92	THR
1	A	96	ARG
1	A	107	ARG
1	A	127	ARG
1	A	182	LEU
1	A	184	THR
1	A	192	ARG
1	A	193	GLN
1	A	195	VAL
1	A	207	ASP
1	A	225	ARG
1	A	226	LEU
1	A	233	LEU
1	A	251	THR
1	A	253	VAL
1	A	257	ILE
1	A	282	LEU
1	A	289	ARG
1	A	292	ASP
1	A	347	LEU
1	A	353	SER
1	A	359	LEU
1	A	361	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	ASN
1	A	83	GLN
1	A	188	HIS
1	A	193	GLN
1	A	198	HIS
1	A	288	HIS
1	A	294	GLN
1	A	303	GLN
1	A	315	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	401	1	14,14,15	1.07	2 (14%)	17,19,21	1.50	2 (11%)
3	2ZO	A	402	-	7,8,8	0.94	0	9,10,10	2.52	4 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	401	1	-	0/6/23/26	0/1/1/1
3	2ZO	A	402	-	-	3/6/6/6	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	NAG	C3-C2	2.28	1.57	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	NAG	O5-C1	-2.26	1.39	1.43

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	2ZO	C1-C2-C3	-5.08	103.23	109.94
3	A	402	2ZO	C2-C1-N	-4.59	99.14	115.97
2	A	401	NAG	C1-C2-N2	3.32	115.67	110.43
2	A	401	NAG	O7-C7-C8	-2.64	117.35	122.05
3	A	402	2ZO	C5-N-C6	2.14	114.61	108.98
3	A	402	2ZO	C7-N-C1	-2.14	101.41	109.91

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	2ZO	C2-C1-N-C5
3	A	402	2ZO	C2-C1-N-C7
3	A	402	2ZO	C2-C1-N-C6

There are no ring outliers.

1 monomer is involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	2ZO	8	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	361/361 (100%)	0.12	22 (6%)	27 25	14, 23, 44, 91	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	362	VAL	6.0
1	A	269	TRP	5.9
1	A	208	ALA	5.6
1	A	204	GLY	5.5
1	A	210	SER	5.0
1	A	78	TRP	4.4
1	A	209	SER	3.8
1	A	81	GLY	3.0
1	A	158	THR	2.9
1	A	361	ARG	2.9
1	A	206	GLU	2.9
1	A	207	ASP	2.8
1	A	120	TRP	2.7
1	A	191	TRP	2.6
1	A	205	ASN	2.4
1	A	80	TYR	2.3
1	A	253	VAL	2.1
1	A	213	PHE	2.1
1	A	348	THR	2.1
1	A	148	THR	2.1
1	A	251	THR	2.0
1	A	257	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	A	401	14/15	0.86	0.10	24,32,39,42	0
3	2ZO	A	402	9/9	0.88	0.10	18,26,28,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.