



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:17 AM UTC

PDB ID : 1Q88 / pdb_00001q88
Title : Crystal structure of the C-domain of the T.vaginalis Inr binding protein, IBP39 (monoclinic form)
Authors : Schumacher, M.A.; Johnson, P.J.
Deposited on : 2003-08-20
Resolution : 2.42 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6062 (2.44-2.40)
Clashscore	190562	6562 (2.44-2.40)
Ramachandran outliers	187476	6481 (2.44-2.40)
Sidechain outliers	187428	6482 (2.44-2.40)
RSRZ outliers	180081	6066 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	5% 55% 23% 6% 14%
1	B	221	4% 60% 25% • • 10%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3173 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 39 kDa initiator binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	189	Total	C	N	O	S	0	0	0
			1540	988	251	292	9			
1	B	198	Total	C	N	O	S	0	0	0
			1603	1028	260	305	10			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	342	HIS	-	expression tag	UNP Q95VR4
A	343	HIS	-	expression tag	UNP Q95VR4
A	344	HIS	-	expression tag	UNP Q95VR4
A	345	HIS	-	expression tag	UNP Q95VR4
A	346	HIS	-	expression tag	UNP Q95VR4
A	347	HIS	-	expression tag	UNP Q95VR4
B	342	HIS	-	expression tag	UNP Q95VR4
B	343	HIS	-	expression tag	UNP Q95VR4
B	344	HIS	-	expression tag	UNP Q95VR4
B	345	HIS	-	expression tag	UNP Q95VR4
B	346	HIS	-	expression tag	UNP Q95VR4
B	347	HIS	-	expression tag	UNP Q95VR4


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	17	Total	O	0	0
			17	17		
2	B	13	Total	O	0	0
			13	13		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 39 kDa initiator binding protein



• Molecule 1: 39 kDa initiator binding protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	99.60Å 76.40Å 80.30Å 90.00° 93.00° 90.00°	Depositor
Resolution (Å)	49.73 – 2.42 49.73 – 2.42	Depositor EDS
% Data completeness (in resolution range)	84.9 (49.73-2.42) 84.9 (49.73-2.42)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.19 (at 2.34Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.190 , 0.232 0.190 , 0.230	Depositor DCC
R_{free} test set	1939 reflections (9.54%)	wwPDB-VP
Wilson B-factor (Å ²)	44.8	Xtriage
Anisotropy	0.179	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 57.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3173	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	1/1577 (0.1%)	1.24	15/2132 (0.7%)
1	B	0.68	5/1643 (0.3%)	1.82	18/2225 (0.8%)
All	All	0.62	6/3220 (0.2%)	1.56	33/4357 (0.8%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	212	ALA	N-CA	-13.62	1.36	1.46
1	B	212	ALA	CA-C	-8.99	1.43	1.52
1	A	212	ALA	CA-C	-6.51	1.44	1.52
1	B	213	PRO	CA-C	-6.50	1.43	1.52
1	B	211	MET	CA-C	-5.86	1.46	1.53
1	B	214	GLN	N-CA	-5.47	1.38	1.46

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	212	ALA	CA-C-N	47.90	179.71	119.84
1	B	212	ALA	C-N-CA	47.90	179.71	119.84
1	A	212	ALA	CA-C-N	20.95	177.28	127.00
1	A	212	ALA	C-N-CA	20.95	177.28	127.00
1	A	212	ALA	C-N-CD	-15.89	85.63	120.60
1	B	213	PRO	CA-N-CD	-10.71	97.00	112.00
1	B	214	GLN	N-CA-C	-10.50	100.22	113.23
1	A	213	PRO	CA-N-CD	-10.31	97.06	111.50
1	B	212	ALA	C-N-CD	-10.31	82.71	125.00
1	A	213	PRO	N-CD-CG	9.83	115.59	103.80
1	B	211	MET	CA-C-N	-8.38	111.89	123.96
1	B	211	MET	C-N-CA	-8.38	111.89	123.96
1	B	213	PRO	N-CA-CB	8.26	111.92	103.25
1	B	213	PRO	N-CD-CG	7.99	115.18	103.20
1	B	213	PRO	CB-CA-C	7.35	123.69	111.56
1	A	213	PRO	N-CA-CB	7.16	110.48	102.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	232	GLY	CA-C-N	7.02	126.94	119.85
1	B	232	GLY	C-N-CA	7.02	126.94	119.85
1	A	324	GLU	N-CA-C	-6.92	104.99	113.50
1	B	212	ALA	CA-C-O	6.46	127.46	120.94
1	A	232	GLY	CA-C-N	6.25	126.16	119.85
1	A	232	GLY	C-N-CA	6.25	126.16	119.85
1	A	231	PHE	N-CA-C	6.25	119.58	107.71
1	B	258	ASP	CA-C-N	6.17	126.35	119.32
1	B	258	ASP	C-N-CA	6.17	126.35	119.32
1	B	231	PHE	N-CA-C	6.10	119.30	107.71
1	A	212	ALA	CA-C-O	5.89	126.70	120.63
1	A	298	LYS	CA-C-N	5.30	126.47	119.84
1	A	298	LYS	C-N-CA	5.30	126.47	119.84
1	B	217	GLN	OE1-CD-NE2	-5.28	117.32	122.60
1	B	214	GLN	OE1-CD-NE2	-5.21	117.39	122.60
1	A	214	GLN	N-CA-C	-5.01	106.02	112.68
1	A	217	GLN	OE1-CD-NE2	-5.00	117.60	122.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1540	0	1511	47	0
1	B	1603	0	1558	44	0
2	A	17	0	0	1	0
2	B	13	0	0	2	0
All	All	3173	0	3069	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:198:GLN:HE22	1:B:238:MET:HG3	1.28	0.98
1:B:191:GLU:HG2	1:B:200:LYS:HD2	1.47	0.97
1:A:159:GLU:HG3	1:B:175:LYS:HA	1.52	0.90
1:A:186:ILE:HD13	1:A:211:MET:HE2	1.66	0.76
1:B:269:ARG:HD3	1:B:269:ARG:H	1.53	0.73
1:A:232:GLY:H	1:A:297:ASN:HD21	1.38	0.72
1:A:297:ASN:O	1:A:299:PRO:HD3	1.91	0.69
1:A:182:ARG:HD3	2:A:361:HOH:O	1.95	0.67
1:B:195:GLN:H	1:B:198:GLN:HE21	1.43	0.65
1:B:260:GLN:HG3	1:B:263:LYS:HB2	1.80	0.63
1:B:272:ILE:HD11	1:B:331:TYR:HB2	1.82	0.61
1:A:155:PRO:HA	1:A:158:GLN:HE21	1.66	0.61
1:A:159:GLU:CG	1:B:175:LYS:HA	2.30	0.60
1:A:264:ASN:HA	1:A:267:ASN:HD22	1.65	0.60
1:B:229:ALA:HB3	1:B:281:PRO:HB2	1.82	0.59
1:B:244:LEU:HD11	1:B:255:LEU:HG	1.84	0.59
1:B:188:LYS:HA	1:B:188:LYS:HE2	1.84	0.59
1:B:297:ASN:O	1:B:299:PRO:HD3	2.04	0.57
1:B:232:GLY:HA2	1:B:299:PRO:HA	1.85	0.57
1:A:298:LYS:HB3	1:A:301:ILE:HD12	1.88	0.56
1:B:200:LYS:O	1:B:204:ILE:HG13	2.04	0.56
1:A:240:LYS:HZ3	1:A:307:TYR:HB2	1.70	0.56
1:A:298:LYS:CB	1:A:301:ILE:HD12	2.36	0.56
1:B:232:GLY:HA3	1:B:240:LYS:NZ	2.22	0.55
1:A:264:ASN:HA	1:A:267:ASN:ND2	2.21	0.55
1:A:181:MET:HE2	1:A:218:THR:HG22	1.87	0.55
1:B:195:GLN:H	1:B:198:GLN:NE2	2.04	0.54
1:A:317:ASP:HB2	1:A:321:LYS:HD3	1.89	0.53
1:B:302:GLU:C	1:B:304:THR:H	2.15	0.53
1:B:182:ARG:O	1:B:186:ILE:HG12	2.09	0.52
1:A:229:ALA:HB3	1:A:281:PRO:HB2	1.90	0.52
1:A:293:ARG:HH11	1:A:293:ARG:HB3	1.74	0.52
1:B:229:ALA:CB	1:B:281:PRO:HB2	2.39	0.52
1:A:232:GLY:HA2	1:A:299:PRO:HA	1.92	0.51
1:A:325:MET:C	1:A:327:PRO:HD3	2.34	0.51
1:A:316:TYR:HA	1:A:321:LYS:HZ1	1.75	0.51
1:A:298:LYS:HD3	1:A:309:MET:HE2	1.94	0.50
1:A:258:ASP:O	1:A:261:ALA:HB2	2.12	0.50
1:B:241:ILE:O	1:B:245:LEU:HB2	2.11	0.50
1:A:317:ASP:H	1:A:321:LYS:HZ3	1.60	0.49
1:A:216:GLU:HG2	1:A:218:THR:O	2.12	0.49
1:A:247:ILE:HD13	1:A:323:PHE:HD2	1.77	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:185:LEU:HD13	1:A:211:MET:HE3	1.95	0.48
1:A:197:GLU:H	1:A:197:GLU:CD	2.21	0.48
1:A:198:GLN:OE1	1:A:238:MET:HG2	2.13	0.48
1:B:324:GLU:HB3	2:B:359:HOH:O	2.14	0.48
1:A:244:LEU:HD13	1:A:255:LEU:HD11	1.96	0.48
1:B:294:LYS:HB2	1:B:311:GLU:HG2	1.95	0.48
1:A:181:MET:CE	1:A:218:THR:HG22	2.44	0.47
1:B:240:LYS:NZ	1:B:297:ASN:HD21	2.12	0.47
1:B:306:GLN:HA	1:B:316:TYR:O	2.14	0.47
1:B:201:GLN:O	1:B:205:GLU:HG3	2.15	0.47
1:B:193:PHE:CE1	1:B:234:TYR:HB2	2.50	0.47
1:A:247:ILE:HD11	1:A:320:ASP:HA	1.97	0.47
1:B:294:LYS:HB2	1:B:311:GLU:CG	2.45	0.47
1:A:227:PHE:HA	1:A:230:MET:HE3	1.97	0.46
1:B:226:LYS:O	1:B:230:MET:HG3	2.14	0.46
1:A:182:ARG:O	1:A:186:ILE:HG12	2.15	0.46
1:B:232:GLY:HA3	1:B:240:LYS:HZ2	1.80	0.46
1:A:293:ARG:HH11	1:A:293:ARG:CB	2.29	0.46
1:B:231:PHE:O	1:B:232:GLY:O	2.33	0.46
1:B:232:GLY:CA	1:B:240:LYS:HZ2	2.28	0.45
1:B:302:GLU:C	1:B:304:THR:N	2.74	0.45
1:A:240:LYS:NZ	1:A:307:TYR:HB2	2.33	0.44
1:A:317:ASP:H	1:A:321:LYS:NZ	2.15	0.44
1:B:321:LYS:O	1:B:325:MET:HG2	2.17	0.44
1:A:240:LYS:HZ3	1:A:307:TYR:CB	2.31	0.44
1:A:224:LEU:HD23	1:A:224:LEU:HA	1.88	0.43
1:A:231:PHE:O	1:A:232:GLY:O	2.37	0.43
1:A:193:PHE:CD1	1:A:234:TYR:HB2	2.54	0.43
1:A:287:LYS:HG2	1:A:292:ILE:CD1	2.48	0.43
1:B:168:ILE:O	1:B:172:LEU:HG	2.19	0.43
1:A:232:GLY:H	1:A:297:ASN:ND2	2.11	0.42
1:A:232:GLY:N	1:A:297:ASN:HD21	2.12	0.42
1:B:292:ILE:O	1:B:292:ILE:HG13	2.18	0.42
1:A:298:LYS:NZ	1:A:309:MET:HE1	2.34	0.42
1:B:198:GLN:NE2	1:B:238:MET:HG3	2.12	0.42
1:B:258:ASP:HA	1:B:259:PRO:HD3	1.87	0.42
1:B:190:ALA:O	1:B:194:LYS:HG2	2.20	0.41
1:B:246:LEU:HD23	2:B:360:HOH:O	2.20	0.41
1:A:295:ILE:HD13	1:A:322:TYR:CE2	2.56	0.41
1:B:290:THR:O	1:B:290:THR:HG22	2.20	0.41
1:A:182:ARG:HD2	1:A:216:GLU:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:THR:OG1	1:A:290:THR:HG23	2.20	0.41
1:B:240:LYS:HZ2	1:B:297:ASN:HD21	1.68	0.41
1:B:247:ILE:HD13	1:B:323:PHE:HD2	1.86	0.41
1:A:258:ASP:HA	1:A:259:PRO:HD3	1.93	0.40
1:B:193:PHE:CD1	1:B:234:TYR:HB2	2.57	0.40
1:B:286:LEU:HA	1:B:286:LEU:HD23	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	185/221 (84%)	168 (91%)	13 (7%)	4 (2%)	5	5
1	B	196/221 (89%)	179 (91%)	13 (7%)	4 (2%)	6	7
All	All	381/442 (86%)	347 (91%)	26 (7%)	8 (2%)	5	6

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	232	GLY
1	B	213	PRO
1	B	232	GLY
1	A	269	ARG
1	B	303	ALA
1	A	177	ASP
1	B	305	GLY
1	A	213	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/198 (84%)	145 (87%)	22 (13%)	4	5
1	B	172/198 (87%)	157 (91%)	15 (9%)	9	15
All	All	339/396 (86%)	302 (89%)	37 (11%)	6	9

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	152	ASN
1	A	159	GLU
1	A	164	LYS
1	A	174	GLN
1	A	180	VAL
1	A	185	LEU
1	A	197	GLU
1	A	201	GLN
1	A	211	MET
1	A	213	PRO
1	A	215	GLU
1	A	219	LYS
1	A	224	LEU
1	A	238	MET
1	A	244	LEU
1	A	255	LEU
1	A	258	ASP
1	A	260	GLN
1	A	287	LYS
1	A	290	THR
1	A	293	ARG
1	A	304	THR
1	B	152	ASN
1	B	156	ASN
1	B	171	ARG
1	B	174	GLN
1	B	201	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	213	PRO
1	B	217	GLN
1	B	244	LEU
1	B	245	LEU
1	B	255	LEU
1	B	258	ASP
1	B	260	GLN
1	B	270	ASP
1	B	286	LEU
1	B	329	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	152	ASN
1	A	174	GLN
1	A	214	GLN
1	A	260	GLN
1	A	267	ASN
1	A	297	ASN
1	B	152	ASN
1	B	198	GLN
1	B	202	ASN
1	B	214	GLN
1	B	217	GLN
1	B	253	HIS
1	B	267	ASN
1	B	278	GLN
1	B	297	ASN
1	B	306	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	189/221 (85%)	0.05	11 (5%) 29 25	31, 45, 112, 123	0
1	B	198/221 (89%)	0.14	8 (4%) 42 38	31, 56, 100, 135	0
All	All	387/442 (87%)	0.10	19 (4%) 35 31	31, 50, 108, 135	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	332	LEU	6.0
1	A	339	ALA	5.5
1	B	345	HIS	4.7
1	B	270	ASP	4.2
1	A	334	ALA	3.6
1	A	333	THR	3.6
1	A	337	THR	3.4
1	A	338	PHE	3.3
1	A	335	TYR	2.7
1	B	271	SER	2.7
1	B	330	THR	2.7
1	B	305	GLY	2.6
1	B	232	GLY	2.5
1	B	265	ALA	2.5
1	A	270	ASP	2.4
1	A	328	ILE	2.3
1	A	232	GLY	2.2
1	A	271	SER	2.2
1	B	269	ARG	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.