



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:25 PM UTC

PDB ID : 6Q8C / pdb_00006q8c
Title : Neisseria gonorrhoeae Leucyl-tRNA Synthetase in Complex with 5'-O-(N-(L-Leucyl)-Sulfamoyl)Uridine
Authors : Pang, L.; De Graef, S.; Strelkov, S.V.; Weeks, S.D.
Deposited on : 2018-12-14
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7754 (2.34-2.30)
Clashscore	190562	8383 (2.34-2.30)
Ramachandran outliers	187476	8303 (2.34-2.30)
Sidechain outliers	187428	8303 (2.34-2.30)
RSRZ outliers	180081	7760 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	877	8% 87% 8% 5%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6540 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Leucine-tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	832	6311	4028	1063	1185	35	0	1	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP A0A1D3HNX6
A	454	ASN	ASP	engineered mutation	UNP A0A1D3HNX6
A	508	ILE	MET	conflict	UNP A0A1D3HNX6

- Molecule 2 is 5'-O-(N-(L-Leucyl)-Sulfamoyl)Uridine (CCD ID: LSU) (formula: C₁₅H₂₄N₄O₉S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	29	15	4	9	1	0	0

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	190	Total	O	0	0
			190	190		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Leucine-tRNA ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.32Å 82.00Å 227.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	113.86 – 2.31 113.86 – 2.31	Depositor EDS
% Data completeness (in resolution range)	100.0 (113.86-2.31) 100.0 (113.86-2.31)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.16Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.200 , 0.251 0.205 , 0.256	Depositor DCC
R_{free} test set	2550 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	46.9	Xtriage
Anisotropy	0.229	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 62.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6540	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, EDO, MG, LSU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.21	0/6474	0.40	0/8822

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6311	0	5841	47	0
2	A	29	0	0	1	0
3	A	1	0	0	0	0
4	A	8	0	12	0	0
5	A	1	0	0	0	0
6	A	190	0	0	2	0
All	All	6540	0	5853	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:314:THR:HG23	1:A:316:ARG:H	1.31	0.95
1:A:314:THR:HG21	1:A:329:VAL:HB	1.62	0.80
1:A:361:TYR:HB2	1:A:363:LEU:HD13	1.64	0.78
1:A:262:LEU:HD23	1:A:322:LEU:HD22	1.73	0.70
1:A:439:TYR:HB2	1:A:478:PRO:HD2	1.79	0.64
1:A:314:THR:HG23	1:A:316:ARG:N	2.10	0.64
1:A:219:MET:HE1	1:A:545:ILE:HD11	1.78	0.64
1:A:53:VAL:HG12	1:A:661:MET:HE1	1.80	0.62
1:A:230:MET:HE1	1:A:412:ALA:HB2	1.85	0.57
1:A:155:THR:HG22	1:A:187:GLU:OE2	2.05	0.57
1:A:134:ARG:HB3	1:A:463:PRO:HB3	1.88	0.55
1:A:52:HIS:HD2	2:A:901:LSU:C6	2.20	0.55
1:A:267:TYR:CE1	1:A:363:LEU:HD23	2.43	0.54
1:A:361:TYR:CB	1:A:363:LEU:HD13	2.35	0.54
1:A:32:LYS:HB3	1:A:72:ASN:HB2	1.90	0.52
1:A:424:LYS:NZ	1:A:426:GLN:HE21	2.08	0.52
1:A:57:THR:O	1:A:61:VAL:HG13	2.10	0.52
1:A:661:MET:HA	1:A:671:LEU:HD11	1.92	0.51
1:A:262:LEU:HD13	1:A:345:MET:HE1	1.93	0.51
1:A:215:GLN:NE2	6:A:1008:HOH:O	2.43	0.50
1:A:394:VAL:HG12	1:A:395:ASN:ND2	2.27	0.50
1:A:58:ILE:O	1:A:61:VAL:HG22	2.11	0.49
1:A:314:THR:HG22	1:A:329:VAL:O	2.13	0.49
1:A:55:ASN:C	1:A:55:ASN:HD22	2.22	0.48
1:A:222:ASN:O	1:A:428[B]:ARG:NH1	2.47	0.48
1:A:11:GLU:HG2	1:A:785:HIS:CG	2.48	0.48
1:A:309:LYS:HD2	1:A:334:TYR:HB3	1.94	0.48
1:A:49:HIS:O	1:A:52:HIS:HB2	2.15	0.46
1:A:230:MET:HE3	1:A:411:VAL:HG23	1.97	0.46
1:A:528:SER:OG	1:A:531:SER:OG	2.24	0.46
1:A:42:PRO:HD2	1:A:78:GLY:O	2.16	0.46
1:A:259:PRO:HB2	1:A:411:VAL:HG21	1.98	0.46
1:A:602:PRO:HD3	1:A:641:ASN:HA	1.98	0.45
1:A:635:MET:HE1	1:A:643:VAL:HG22	1.98	0.45
1:A:545:ILE:HB	1:A:579:THR:HG21	1.99	0.45
1:A:557:PHE:CZ	1:A:561:LEU:HD11	2.53	0.44
1:A:35:TYR:CE2	1:A:66:LYS:HG3	2.54	0.43
1:A:11:GLU:HG2	1:A:785:HIS:CD2	2.53	0.43
1:A:40:MET:SD	1:A:543:GLY:HA3	2.60	0.42
1:A:230:MET:HE2	1:A:230:MET:HB2	1.99	0.41
1:A:244:LEU:HB3	1:A:249:ALA:HA	2.01	0.41
1:A:374:ASN:HB3	1:A:384:TRP:CE2	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:86:ALA:HB1	1:A:97:PRO:HB3	2.02	0.41
1:A:276:LEU:HD21	1:A:329:VAL:HG21	2.03	0.41
1:A:137:GLN:O	1:A:141:THR:HG23	2.22	0.40
1:A:82:PHE:CG	1:A:441:GLY:HA2	2.57	0.40
1:A:722:LYS:NZ	6:A:1025:HOH:O	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	821/877 (94%)	794 (97%)	27 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	603/717 (84%)	601 (100%)	2 (0%)	86	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	238	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	ASN
1	A	52	HIS
1	A	362	ASN
1	A	382	GLN
1	A	390	ASN
1	A	395	ASN
1	A	426	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	LSU	A	901	-	29,30,30	0.67	0	38,44,44	0.71	2 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	904	-	3,3,3	0.45	0	2,2,2	0.27	0
4	EDO	A	903	-	3,3,3	0.46	0	2,2,2	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LSU	A	901	-	-	3/22/39/39	0/2/2/2
4	EDO	A	904	-	-	0/1/1/1	-
4	EDO	A	903	-	-	0/1/1/1	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	901	LSU	O-C-N3S	-2.74	118.01	122.98
2	A	901	LSU	C-N3S-S	2.12	131.84	124.07

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	901	LSU	O-C-CA-N
2	A	901	LSU	N3S-C-CA-N
2	A	901	LSU	C5'-O5'-S-N3S


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	LSU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	832/877 (94%)	0.47	68 (8%) 17 20	29, 61, 103, 141	1 (0%)

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	608	THR	6.0
1	A	817	ILE	5.4
1	A	607	LEU	4.4
1	A	626	PRO	4.1
1	A	301	GLU	4.0
1	A	598	ASP	4.0
1	A	872	VAL	3.8
1	A	606	GLU	3.7
1	A	871	LEU	3.7
1	A	623	ASP	3.7
1	A	379	ASN	3.5
1	A	172	ILE	3.4
1	A	864	ILE	3.4
1	A	302	ALA	3.4
1	A	339	TYR	3.3
1	A	875	VAL	3.2
1	A	475	MET	3.1
1	A	850	GLU	3.1
1	A	824	ASN	3.0
1	A	827	LEU	3.0
1	A	874	ILE	3.0
1	A	10	ILE	3.0
1	A	834	ALA	2.9
1	A	319	VAL	2.9
1	A	815	SER	2.8
1	A	173	ASP	2.7
1	A	177	TRP	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	599	TRP	2.7
1	A	589	TYR	2.6
1	A	826	LYS	2.6
1	A	628	VAL	2.6
1	A	825	GLY	2.6
1	A	683	ARG	2.6
1	A	297	GLY	2.5
1	A	867	VAL	2.5
1	A	378	ALA	2.5
1	A	306	THR	2.4
1	A	800	TRP	2.4
1	A	366	LYS	2.4
1	A	698	GLN	2.4
1	A	813	VAL	2.4
1	A	847	LEU	2.4
1	A	641	ASN	2.3
1	A	869	GLY	2.3
1	A	603	ALA	2.3
1	A	375	ALA	2.3
1	A	600	ILE	2.3
1	A	868	PRO	2.3
1	A	477	SER	2.3
1	A	823	VAL	2.2
1	A	866	VAL	2.2
1	A	304	MET	2.2
1	A	327	LEU	2.2
1	A	363	LEU	2.2
1	A	629	ILE	2.2
1	A	180	GLY	2.2
1	A	848	ALA	2.2
1	A	627	VAL	2.1
1	A	155	THR	2.1
1	A	48	LEU	2.1
1	A	174	GLY	2.1
1	A	340	GLY	2.1
1	A	314	THR	2.1
1	A	338	GLY	2.1
1	A	840	ALA	2.0
1	A	399	LEU	2.0
1	A	171	VAL	2.0
1	A	870	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EDO	A	903	4/4	0.92	0.15	45,60,62,64	0
4	EDO	A	904	4/4	0.93	0.13	63,63,73,73	0
2	LSU	A	901	29/29	0.95	0.08	30,41,69,92	0
3	ZN	A	902	1/1	0.99	0.02	50,50,50,50	0
5	MG	A	905	1/1	0.99	0.03	71,71,71,71	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.