



Full wwPDB EM Validation Report ⓘ

Mar 8, 2026 – 01:37 PM UTC

PDB ID : 8Q8V / pdb_00008q8v
EMDB ID : EMD-18262
Title : Tau - CTE-MIA3
Authors : Lovestam, S.; Li, D.; Scheres, S.H.W.; Goedert, M.
Deposited on : 2023-08-18
Resolution : 3.80 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	10198 (3.30 - 4.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	382	
1	B	382	
1	C	382	
1	D	382	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	382	 18% 80%
1	F	382	 18% 80%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 7032 atoms, of which 3588 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isoform Tau-D of Microtubule-associated protein tau.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	75	Total	C	H	N	O	S	0	0
			1172	360	598	106	107	1		
1	C	75	Total	C	H	N	O	S	0	0
			1172	360	598	106	107	1		
1	E	75	Total	C	H	N	O	S	0	0
			1172	360	598	106	107	1		
1	B	75	Total	C	H	N	O	S	0	0
			1172	360	598	106	107	1		
1	D	75	Total	C	H	N	O	S	0	0
			1172	360	598	106	107	1		
1	F	75	Total	C	H	N	O	S	0	0
			1172	360	598	106	107	1		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	ALA	deletion	UNP P10636
C	?	-	ALA	deletion	UNP P10636
E	?	-	ALA	deletion	UNP P10636
B	?	-	ALA	deletion	UNP P10636
D	?	-	ALA	deletion	UNP P10636
F	?	-	ALA	deletion	UNP P10636

VAL SER SER THR GLY SER ILE ASP MET VAL ASP SER SER PRO GLN LEU ALA THR LEU LEU ALA ASP GLU VAL SER ALA LEU LEU

• Molecule 1: Isoform Tau-D of Microtubule-associated protein tau

Chain E:  18% 80%

MET ALA GLU PRO THR LEU ARG GLN SER GLU PHE MET VAL MET ASP SER SER PRO GLN LEU ALA THR LEU LEU

GLY HIS VAL THR PRO THR LEU SER GLN ARG ALA MET VAL MET VAL SER LYS SER LYS ASP HIS ALA THR LEU LEU

LYS THR PRO THR PRO THR SER GLY GLU VAL PRO LYS SER LYS ASP GLY ASP ARG THR LEU LEU LEU LEU

LYS SER ARG THR GLN ALA PRO VAL PRO MET LYS SER LYS ASP GLY ASP ARG THR LEU LEU LEU LEU

VAL PRO GLY GLY S395 Y310 D314 G335 Q336 V337 E338 S341 E342 K343 L344 D345 F346 K347 D348 R349 D358 R379

VAL SER SER THR GLY SER ILE ASP MET VAL ASP SER SER PRO GLN LEU ALA THR LEU LEU ALA ASP GLU VAL SER ALA LEU LEU

• Molecule 1: Isoform Tau-D of Microtubule-associated protein tau

Chain B:  19% 80%

MET ALA GLU PRO THR LEU ARG GLN SER GLU PHE MET VAL MET ASP SER SER PRO GLN LEU ALA THR LEU LEU

GLY HIS VAL THR PRO THR LEU SER GLN ARG ALA MET VAL MET VAL SER LYS SER LYS ASP HIS ALA THR LEU LEU

LYS THR PRO THR PRO THR SER GLY GLU VAL PRO LYS SER LYS ASP GLY ASP ARG THR LEU LEU LEU LEU

LYS SER ARG THR GLN ALA PRO VAL VAL PRO MET LYS SER LYS ASP GLY ASP ARG THR LEU LEU LEU LEU

VAL PRO GLY GLY S395 D314 V339 R340 S341 L344 D345 F346 K347 D348 R349 D358 R379

ILE ASP MET VAL ASP PRO GLN LEU VAL ALA THR LEU LEU LEU LEU LEU LEU LEU LEU LEU LEU


• Molecule 1: Isoform Tau-D of Microtubule-associated protein tau

Chain D:  19% 80%

MET ALA GLU PRO THR LEU ARG GLN SER GLU PHE MET VAL MET ASP SER SER PRO GLN LEU ALA THR LEU LEU

GLY HIS VAL THR PRO THR LEU SER GLN ARG ALA MET VAL MET VAL SER LYS SER LYS ASP HIS ALA THR LEU LEU

- Molecule 1: Isoform Tau-D of Microtubule-associated protein tau



4 Experimental information

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=-1.13°, rise=4.77 Å, axial sym=C1	Depositor
Number of segments used	4243	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{Å}^2$)	40	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	0.011	Depositor
Minimum map value	-0.006	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.00261	Depositor
Map size (Å)	316.41602, 316.41602, 316.41602	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.82400006, 0.82400006, 0.82400006	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/583	1.30	0/780
1	B	0.76	0/583	1.27	0/780
1	C	0.79	0/583	1.33	2/780 (0.3%)
1	D	0.79	0/583	1.33	2/780 (0.3%)
1	E	0.79	0/583	1.34	4/780 (0.5%)
1	F	0.79	0/583	1.34	4/780 (0.5%)
All	All	0.78	0/3498	1.32	12/4680 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1
1	F	0	1
All	All	0	2

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	348	ASP	CA-C-N	5.63	129.69	121.31
1	E	348	ASP	C-N-CA	5.63	129.69	121.31
1	F	348	ASP	CA-C-N	5.59	129.65	121.31
1	F	348	ASP	C-N-CA	5.59	129.65	121.31
1	E	343	LYS	CA-C-N	5.30	128.24	120.71
1	E	343	LYS	C-N-CA	5.30	128.24	120.71
1	F	343	LYS	CA-C-N	5.29	128.22	120.71
1	F	343	LYS	C-N-CA	5.29	128.22	120.71
1	D	348	ASP	CA-C-N	5.11	128.92	121.31
1	D	348	ASP	C-N-CA	5.11	128.92	121.31

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	348	ASP	CA-C-N	5.10	128.91	121.31
1	C	348	ASP	C-N-CA	5.10	128.91	121.31

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	310	TYR	Sidechain
1	F	310	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	574	598	597	2	0
1	B	574	598	597	2	0
1	C	574	598	597	2	0
1	D	574	598	597	2	0
1	E	574	598	597	2	0
1	F	574	598	597	2	0
All	All	3444	3588	3582	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:341:SER:HB3	1:D:344:LEU:HD11	1.89	0.55
1:C:341:SER:HB3	1:C:344:LEU:HD11	1.89	0.54
1:D:341:SER:CB	1:D:344:LEU:HD11	2.41	0.51
1:C:341:SER:CB	1:C:344:LEU:HD11	2.41	0.49
1:F:341:SER:CB	1:F:344:LEU:HD11	2.47	0.44
1:E:341:SER:CB	1:E:344:LEU:HD11	2.47	0.44
1:A:341:SER:HB3	1:A:344:LEU:HD11	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:341:SER:CB	1:B:344:LEU:HD11	2.49	0.42
1:B:341:SER:HB3	1:B:344:LEU:HD11	2.01	0.42
1:E:341:SER:HB3	1:E:344:LEU:HD11	2.02	0.41
1:A:341:SER:CB	1:A:344:LEU:HD11	2.49	0.41
1:F:341:SER:HB3	1:F:344:LEU:HD11	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	73/382 (19%)	70 (96%)	3 (4%)	0	100	100
1	B	73/382 (19%)	70 (96%)	3 (4%)	0	100	100
1	C	73/382 (19%)	67 (92%)	6 (8%)	0	100	100
1	D	73/382 (19%)	67 (92%)	6 (8%)	0	100	100
1	E	73/382 (19%)	68 (93%)	5 (7%)	0	100	100
1	F	73/382 (19%)	68 (93%)	5 (7%)	0	100	100
All	All	438/2292 (19%)	410 (94%)	28 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	66/312 (21%)	66 (100%)	0	100	100
1	B	66/312 (21%)	66 (100%)	0	100	100
1	C	66/312 (21%)	66 (100%)	0	100	100
1	D	66/312 (21%)	66 (100%)	0	100	100
1	E	66/312 (21%)	66 (100%)	0	100	100
1	F	66/312 (21%)	66 (100%)	0	100	100
All	All	396/1872 (21%)	396 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	307	GLN
1	A	351	GLN
1	A	362	HIS
1	C	307	GLN
1	C	351	GLN
1	E	307	GLN
1	E	351	GLN
1	B	307	GLN
1	B	351	GLN
1	B	362	HIS
1	D	307	GLN
1	D	330	HIS
1	D	351	GLN
1	F	307	GLN
1	F	351	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-18262. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 192




Y Index: 192




Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 219




Y Index: 209




Z Index: 198


6.3.2 Raw map



X Index: 191



Y Index: 187




Z Index: 182


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

This section was not generated.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 317 nm³; this corresponds to an approximate mass of 286 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.263 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.263 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.80	-	-
Author-provided FSC curve	3.77	8.02	4.06
Unmasked-calculated*	6.94	15.87	7.03

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.94 differs from the reported value 3.8 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMD map EMD-18262 and PDB model 8Q8V. Per-residue inclusion information can be found in section 3 on page 5.

9.1 Map-model overlay [i](#)


This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.00261).

9.4 Atom inclusion [i](#)



At the recommended contour level, 61% of all backbone atoms, 60% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.00261) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6020	0.4370
A	0.6130	0.4510
B	0.6220	0.4290
C	0.6080	0.4490
D	0.6120	0.4160
E	0.6170	0.4530
F	0.6100	0.4240

