



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:54 PM UTC

PDB ID : 6Q9N / pdb_00006q9n
Title : Crystal structure of PBP2a from MRSA in complex with piperacillin and quinazolinone
Authors : Martinez-Caballero, S.; Batuecas, M.T.; Hermoso, J.A.
Deposited on : 2018-12-18
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	642	
1	B	642	

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 10419 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Penicillin binding protein 2 prime.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	638	Total	C	N	O	S	0	1	0
			5127	3231	865	1016	15			
1	B	637	Total	C	N	O	S	0	0	0
			5112	3223	863	1011	15			

- Molecule 2 is 3-[2-[(E)-2-(4-ethynylphenyl)ethenyl]-4-oxidanylidene-quinazolin-3-yl]benzoic acid (CCD ID: QLN) (formula: C₂₅H₁₆N₂O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			30	25	2	3		

- Molecule 3 is Piperacillin (Open Form) (CCD ID: JPP) (formula: C₂₃H₂₉N₅O₇S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			36	23	5	7	1		


- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Cl	0	0
			2	2		
4	B	2	Total	Cl	0	0
			2	2		

- Molecule 5 is CADMIUM ION (CCD ID: CD) (formula: Cd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Cd	0	0
			2	2		
5	B	2	Total	Cd	0	0
			2	2		

- Molecule 6 is beta-muramic acid (CCD ID: MUR) (formula: C₉H₁₇NO₇).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			17	9	1	7		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	59	Total	O	0	0
			59	59		
7	B	30	Total	O	0	0
			30	30		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Penicillin binding protein 2 prime



• Molecule 1: Penicillin binding protein 2 prime



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.17Å 102.88Å 187.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.55 – 2.50 49.55 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.55-2.50) 99.9 (49.55-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.221 , 0.285 0.222 , 0.281	Depositor DCC
R_{free} test set	2763 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	58.5	Xtriage
Anisotropy	0.774	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 39.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10419	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, QLN, CD, MUR, JPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.97	1/5212 (0.0%)	1.39	3/7008 (0.0%)
1	B	0.98	0/5197	1.39	0/6988
All	All	0.97	1/10409 (0.0%)	1.39	3/13996 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	133	ILE	N-CA	5.05	1.49	1.45

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	403	SER	CA-CB-OG	8.98	129.06	111.10
1	A	403	SER	CB-CA-C	-6.04	98.55	110.27
1	A	58	GLY	CA-C-O	-5.19	118.09	122.29

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	306	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5127	0	5119	42	0
1	B	5112	0	5110	44	0
2	A	30	0	0	1	0
3	A	36	0	27	7	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	17	0	16	2	0
7	A	59	0	0	1	0
7	B	30	0	0	0	0
All	All	10419	0	10272	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:462:SER:HB3	3:A:702:JPP:C03	1.99	0.93
1:B:403:SER:OG	1:B:599:GLY:HA3	1.79	0.82
1:B:425:TYR:OH	1:B:473:GLU:HG3	1.81	0.80
1:A:112:VAL:HG13	1:A:134:PRO:HB3	1.70	0.72
1:B:403:SER:OG	1:B:599:GLY:CA	2.39	0.70
1:A:462:SER:HB3	3:A:702:JPP:H13	1.74	0.69
1:B:583:HIS:O	1:B:587:ILE:HG22	1.97	0.64
1:B:112:VAL:HG13	1:B:134:PRO:HB3	1.79	0.63
1:A:408:LEU:HD22	1:A:534:ILE:HG21	1.82	0.62
1:B:146:ASN:HD21	1:B:295:ASP:HB3	1.65	0.62
1:B:38:GLU:OE2	1:B:83:ARG:NH2	2.33	0.61
1:B:173:ILE:HG22	1:B:178:VAL:HG21	1.82	0.61
1:B:264:LEU:HD21	1:B:273:LYS:O	2.01	0.60
1:A:241:ARG:HB2	6:A:707:MUR:H1	1.84	0.59
1:A:462:SER:HB3	3:A:702:JPP:H14	1.81	0.57
1:A:256:VAL:O	1:A:372:MET:HG2	2.05	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:464:ASN:ND2	3:A:702:JPP:O06	2.34	0.56
1:A:612:ARG:NH2	1:A:635:ASP:OD2	2.39	0.54
1:A:471:ALA:HB1	1:A:514:LEU:HD22	1.90	0.53
1:B:351:HIS:CD2	1:B:666:ILE:HD13	2.43	0.53
1:B:408:LEU:HD22	1:B:534:ILE:HG21	1.91	0.53
1:A:516:ASP:HA	1:A:519:TYR:CE1	2.44	0.52
1:A:631:ILE:HD13	1:A:648:ILE:HG22	1.91	0.52
1:A:327:THR:HB	1:A:356:GLU:HB3	1.92	0.51
1:B:306:SER:O	1:B:308:THR:N	2.45	0.50
1:B:269:TYR:HA	1:B:272:TYR:CD1	2.47	0.49
1:A:99:TYR:HD2	1:A:112:VAL:HG11	1.78	0.48
1:A:220:MET:HA	1:A:224:LEU:CD2	2.43	0.48
1:A:462:SER:CB	3:A:702:JPP:C03	2.83	0.48
1:A:65:ARG:HB3	1:A:66:PRO:HD3	1.96	0.47
1:A:305:ASN:HD22	1:B:68:LYS:HB3	1.80	0.47
1:A:462:SER:O	3:A:702:JPP:H14	2.15	0.47
1:A:506:LYS:HE2	1:A:506:LYS:O	2.15	0.47
1:A:143:HIS:HB3	1:A:145:GLU:OE1	2.15	0.47
1:B:535:TYR:O	1:B:628:MET:HE1	2.15	0.47
3:A:702:JPP:H6	3:A:702:JPP:H15	1.56	0.46
1:B:309:ILE:HD12	1:B:309:ILE:N	2.30	0.46
1:B:327:THR:OG1	1:B:549:LEU:HA	2.16	0.46
1:B:89:SER:HB3	1:B:92:LYS:HB2	1.98	0.45
1:B:156:ASP:OD1	1:B:156:ASP:C	2.58	0.45
6:A:707:MUR:H7	6:A:707:MUR:H2	1.64	0.45
1:B:89:SER:OG	1:B:90:LYS:N	2.50	0.45
1:A:68:LYS:HB3	1:B:305:ASN:HD22	1.80	0.45
1:A:595:ILE:HD11	1:A:620:TYR:CZ	2.52	0.45
1:A:157:ARG:HD3	1:A:666:ILE:O	2.18	0.44
1:B:99:TYR:HD2	1:B:112:VAL:HG11	1.82	0.44
1:A:421:ASP:OD1	1:A:421:ASP:N	2.51	0.44
1:A:351:HIS:HD2	1:A:626:ASN:O	2.00	0.44
1:A:595:ILE:HD11	1:A:620:TYR:OH	2.17	0.43
1:A:62:MET:HE3	1:A:127:TRP:CG	2.53	0.43
1:B:173:ILE:CG2	1:B:178:VAL:HG21	2.47	0.43
1:A:171:ILE:HG12	1:A:224:LEU:HD11	2.01	0.43
1:A:432:TRP:CZ2	1:A:469:ARG:HD3	2.53	0.43
1:B:33:THR:O	1:B:36:ALA:HB3	2.19	0.43
1:A:112:VAL:CG1	1:A:134:PRO:HB3	2.45	0.43
1:B:603:LEU:HD21	1:B:612:ARG:NE	2.33	0.43
1:B:392:LEU:HD21	1:B:397:ILE:HD13	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:537:ALA:CB	1:B:562:ILE:HD11	2.48	0.42
1:A:146:ASN:HD22	2:A:701:QLN:C16	2.32	0.42
1:A:665:ASP:OD2	1:B:554:LYS:NZ	2.53	0.42
1:B:173:ILE:HG13	1:B:201:MET:CE	2.49	0.42
1:B:252:LEU:HD23	1:B:252:LEU:HA	1.90	0.42
1:B:516:ASP:HA	1:B:519:TYR:CE1	2.54	0.42
1:B:615:GLY:C	1:B:645:ASN:ND2	2.77	0.42
1:B:173:ILE:HG13	1:B:201:MET:SD	2.59	0.42
1:A:287:TYR:CZ	1:A:550:LEU:HD11	2.55	0.42
1:B:603:LEU:C	1:B:603:LEU:HD12	2.45	0.42
1:A:612:ARG:HH21	1:A:635:ASP:CG	2.27	0.41
1:A:187:ALA:HB1	1:A:192:ILE:O	2.20	0.41
1:A:209:ASP:OD2	1:B:135:GLY:HA2	2.20	0.41
1:B:172:GLY:HA3	1:B:212:VAL:O	2.21	0.41
1:A:408:LEU:HD23	1:A:408:LEU:N	2.36	0.41
1:B:263:GLU:O	1:B:266:GLN:HB2	2.20	0.41
1:B:403:SER:CB	1:B:599:GLY:HA2	2.50	0.41
1:B:177:ASN:C	1:B:177:ASN:HD22	2.29	0.41
1:B:587:ILE:HD13	1:B:617:PHE:CE1	2.55	0.41
1:A:220:MET:HA	1:A:224:LEU:HD22	2.03	0.41
1:A:174:VAL:HA	1:A:210:THR:O	2.22	0.40
1:B:603:LEU:O	1:B:604:LYS:C	2.64	0.40
1:A:394:LYS:NZ	7:A:805:HOH:O	2.54	0.40
1:B:42:PHE:HB3	1:B:63:THR:HA	2.02	0.40
1:A:288:ASP:O	1:A:292:GLN:HG2	2.21	0.40
1:B:256:VAL:O	1:B:372:MET:HG2	2.22	0.40
1:B:345:GLY:HA3	1:B:632:ASN:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	635/642 (99%)	598 (94%)	33 (5%)	4 (1%)	21	38
1	B	633/642 (99%)	599 (95%)	29 (5%)	5 (1%)	16	31
All	All	1268/1284 (99%)	1197 (94%)	62 (5%)	9 (1%)	18	34

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	176	LYS
1	A	306	SER
1	B	307	ASN
1	B	555	ASN
1	A	307	ASN
1	A	504	SER
1	B	89	SER
1	B	221	ASP
1	B	635	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	570/572 (100%)	535 (94%)	35 (6%)	17	35
1	B	568/572 (99%)	526 (93%)	42 (7%)	13	27
All	All	1138/1144 (100%)	1061 (93%)	77 (7%)	14	31

All (77) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	LYS
1	A	75	VAL
1	A	82	ASP
1	A	87	LYS
1	A	89	SER
1	A	92	LYS
1	A	93	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	94	ARG
1	A	102	LYS
1	A	112	VAL
1	A	125	LEU
1	A	156	ASP
1	A	171	ILE
1	A	177	ASN
1	A	195	ASP
1	A	219	LYS
1	A	220	MET
1	A	229	LYS
1	A	252	LEU
1	A	365	SER
1	A	367	ASP
1	A	391	LEU
1	A	421	ASP
1	A	433	GLN
1	A	444	THR
1	A	447	GLU
1	A	506	LYS
1	A	508	LEU
1	A	514	LEU
1	A	563	ILE
1	A	604	LYS
1	A	609	GLU
1	A	610	THR
1	A	622	LYS
1	A	662	LYS
1	B	30	ILE
1	B	34	ILE
1	B	40	LYS
1	B	44	GLN
1	B	75	VAL
1	B	86	LYS
1	B	87	LYS
1	B	89	SER
1	B	93	LYS
1	B	96	ASP
1	B	101	ILE
1	B	102	LYS
1	B	112	VAL
1	B	120	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	156	ASP
1	B	173	ILE
1	B	177	ASN
1	B	181	LYS
1	B	219	LYS
1	B	222	GLU
1	B	245	LEU
1	B	252	LEU
1	B	308	THR
1	B	309	ILE
1	B	391	LEU
1	B	403	SER
1	B	419	LEU
1	B	424	SER
1	B	428	ASP
1	B	433	GLN
1	B	444	THR
1	B	503	ILE
1	B	506	LYS
1	B	514	LEU
1	B	523	GLU
1	B	532	LEU
1	B	536	SER
1	B	554	LYS
1	B	557	VAL
1	B	595	ILE
1	B	641	MET
1	B	662	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (30) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	ASN
1	A	79	ASN
1	A	91	ASN
1	A	137	GLN
1	A	146	ASN
1	A	177	ASN
1	A	200	GLN
1	A	293	HIS
1	A	305	ASN
1	A	342	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	442	ASN
1	A	452	ASN
1	A	583	HIS
1	A	624	ASN
1	A	661	ASN
1	B	81	GLN
1	B	91	ASN
1	B	107	ASN
1	B	146	ASN
1	B	177	ASN
1	B	203	GLN
1	B	305	ASN
1	B	339	ASN
1	B	351	HIS
1	B	452	ASN
1	B	510	ASN
1	B	569	ASN
1	B	624	ASN
1	B	632	ASN
1	B	661	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 8 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	MUR	A	707	-	17,17,17	0.77	0	21,24,24	1.84	5 (23%)
2	QLN	A	701	-	32,33,33	1.38	4 (12%)	42,46,46	1.55	8 (19%)
3	JPP	A	702	1	33,38,38	4.37	11 (33%)	41,55,55	2.97	18 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MUR	A	707	-	-	5/10/30/30	0/1/1/1
2	QLN	A	701	-	-	10/15/15/15	0/4/4/4
3	JPP	A	702	1	-	5/26/64/64	0/3/3/3

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	702	JPP	O04-C06	11.47	1.43	1.23
3	A	702	JPP	O-C14	10.35	1.44	1.23
3	A	702	JPP	O05-C11	10.16	1.43	1.23
3	A	702	JPP	O06-C21	10.14	1.42	1.23
3	A	702	JPP	C14-C11	-7.38	1.41	1.53
3	A	702	JPP	C10-N	-6.97	1.34	1.47
3	A	702	JPP	C09-N01	-5.96	1.38	1.47
3	A	702	JPP	C10-C09	-3.14	1.39	1.51
3	A	702	JPP	O02-C04	-2.41	1.23	1.30
2	A	701	QLN	C19-C13	2.28	1.43	1.39
3	A	702	JPP	C06-N01	-2.25	1.37	1.41
3	A	702	JPP	C17-S	-2.23	1.79	1.84
2	A	701	QLN	C15-C13	2.19	1.43	1.39
2	A	701	QLN	C19-C18	2.08	1.42	1.39
2	A	701	QLN	C25-C24	2.07	1.42	1.38

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	702	JPP	C20-N03-C21	-7.52	114.47	123.08

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	702	JPP	C19-S-C17	-6.89	75.93	93.41
6	A	707	MUR	O5-C1-C2	6.43	117.01	109.51
3	A	702	JPP	C18-C19-S	5.91	115.09	103.86
3	A	702	JPP	C03-C19-S	-5.38	100.52	109.18
3	A	702	JPP	C02-C19-S	5.15	117.47	109.18
3	A	702	JPP	C10-N-C14	4.59	129.99	120.82
3	A	702	JPP	O-C14-N	4.51	128.66	123.55
2	A	701	QLN	C13-N02-C14	4.37	125.61	119.86
3	A	702	JPP	C02-C19-C18	-4.25	98.23	112.33
3	A	702	JPP	CA-N04-C06	-4.25	113.08	121.22
2	A	701	QLN	C11-N05-C14	3.66	122.66	118.23
2	A	701	QLN	O03-C12-N02	3.56	125.12	120.38
3	A	702	JPP	C05-CA-C21	-3.38	100.47	108.42
2	A	701	QLN	C01-C12-N02	3.14	118.98	114.69
3	A	702	JPP	C15-N-C14	-3.10	115.88	119.59
3	A	702	JPP	C10-C09-N01	3.01	113.72	109.06
2	A	701	QLN	O03-C12-C01	-2.81	115.90	123.11
3	A	702	JPP	O04-C06-N01	2.78	122.81	119.59
3	A	702	JPP	O06-C21-CA	2.67	125.04	120.58
6	A	707	MUR	C1-C2-C3	2.63	114.80	110.10
6	A	707	MUR	C3-C4-C5	2.61	115.13	109.57
3	A	702	JPP	C17-C20-N03	-2.59	104.44	109.93
3	A	702	JPP	O04-C06-N04	-2.49	116.07	122.96
6	A	707	MUR	O1-C1-O5	-2.33	103.48	110.41
6	A	707	MUR	C4-C3-C2	2.22	115.75	111.37
2	A	701	QLN	C01-C11-N05	-2.19	120.21	122.41
2	A	701	QLN	C19-C13-N02	2.15	121.96	119.22
3	A	702	JPP	C04-C18-N02	-2.15	107.30	110.62
2	A	701	QLN	C18-C19-C13	2.04	121.91	119.61
3	A	702	JPP	C05-CA-N04	-2.00	107.69	112.75

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	702	JPP	C-C15-N-C10
6	A	707	MUR	C2-C3-O3-C7
6	A	707	MUR	O3-C7-C8-O8
6	A	707	MUR	O3-C7-C8-O9
6	A	707	MUR	C9-C7-C8-O8
6	A	707	MUR	C9-C7-C8-O9
2	A	701	QLN	C19-C18-C20-O08

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	701	QLN	C17-C18-C20-O08
2	A	701	QLN	C19-C18-C20-O06
2	A	701	QLN	C17-C18-C20-O06
2	A	701	QLN	C21-C22-C23-C24
2	A	701	QLN	C21-C22-C23-C28
2	A	701	QLN	C15-C13-N02-C14
2	A	701	QLN	C19-C13-N02-C14
2	A	701	QLN	C15-C13-N02-C12
2	A	701	QLN	C19-C13-N02-C12
3	A	702	JPP	C01-C20-N03-C21
3	A	702	JPP	C21-CA-N04-C06
3	A	702	JPP	O02-C04-C18-N02
3	A	702	JPP	C-C15-N-C14


There are no ring outliers.


3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	707	MUR	2	0
2	A	701	QLN	1	0
3	A	702	JPP	7	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	638/642 (99%)	0.18	15 (2%) 59 55	32, 71, 108, 150	1 (0%)
1	B	637/642 (99%)	0.24	11 (1%) 69 65	47, 74, 117, 145	0
All	All	1275/1284 (99%)	0.21	26 (2%) 65 61	32, 73, 111, 150	1 (0%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	587	ILE	3.8
1	A	435	ASP	3.1
1	A	434	LYS	3.0
1	A	458	ALA	2.9
1	A	571	LEU	2.7
1	A	459	ILE	2.7
1	A	223	TYR	2.7
1	A	508	LEU	2.6
1	B	610	THR	2.6
1	B	579	VAL	2.5
1	A	455	LEU	2.4
1	A	27	ASP	2.4
1	A	503	ILE	2.4
1	A	504	SER	2.3
1	A	28	LYS	2.3
1	B	223	TYR	2.3
1	B	580	ASN	2.2
1	A	214	LEU	2.2
1	B	88	VAL	2.1
1	A	467	PHE	2.1
1	B	617	PHE	2.1
1	B	645	ASN	2.1
1	B	90	LYS	2.0
1	B	446	TYR	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	85	ILE	2.0
1	B	586	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	MUR	A	707	17/17	0.67	0.19	76,83,99,101	0
2	QLN	A	701	30/30	0.72	0.21	106,135,141,142	0
3	JPP	A	702	36/36	0.85	0.11	68,77,84,86	0
4	CL	A	703	1/1	0.97	0.06	74,74,74,74	0
4	CL	B	702	1/1	0.98	0.09	50,50,50,50	0
4	CL	A	704	1/1	0.99	0.03	50,50,50,50	0
4	CL	B	701	1/1	0.99	0.08	54,54,54,54	0
5	CD	A	706	1/1	1.00	0.01	58,58,58,58	0
5	CD	B	703	1/1	1.00	0.02	58,58,58,58	0
5	CD	B	704	1/1	1.00	0.01	60,60,60,60	0
5	CD	A	705	1/1	1.00	0.02	73,73,73,73	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MUR A 707:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around QLN A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around JPP A 702:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.