



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:59 AM UTC

PDB ID : 6Q9T / pdb_00006q9t
Title : Protein-aromatic foldamer complex crystal structure
Authors : Post, S.; Langlois d'Estaintot, B.; Fischer, L.; Granier, T.; Huc, I.
Deposited on : 2018-12-18
Resolution : 2.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5070 (2.70-2.66)
Clashscore	190562	5409 (2.70-2.66)
Ramachandran outliers	187476	5324 (2.70-2.66)
Sidechain outliers	187428	5324 (2.70-2.66)
RSRZ outliers	180081	5070 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	259	
2	B	16	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2009 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbonic anhydrase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	237	Total	C	N	O	S	0	1	0
			1778	1142	306	328	2			

- Molecule 2 is a protein called Aromatic foldamer.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	16	Total	C	N	O	S	0	0	0
			224	157	33	33	1			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	O	0	0
			4	4		
4	B	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Carbonic anhydrase 2



- Molecule 2: Aromatic foldamer



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	79.30Å 80.96Å 45.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.62 – 2.68 45.62 – 2.68	Depositor EDS
% Data completeness (in resolution range)	98.5 (45.62-2.68) 98.9 (45.62-2.68)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.215 , 0.260 0.219 , 0.263	Depositor DCC
R_{free} test set	431 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	73.2	Xtriage
Anisotropy	0.682	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 68.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.010 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2009	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZY9, 4SO, QUK, QDD, ZN, QUJ, QVE, QVS, A1IJ4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.17	2/1830 (0.1%)	1.49	7/2503 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	6

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	195	GLY	C-O	6.99	1.29	1.24
1	A	149	VAL	C-O	5.73	1.31	1.24

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	132	LYS	N-CA-C	-5.82	105.28	112.90
1	A	209	ILE	CB-CA-C	5.76	117.05	111.23
1	A	111	LYS	CB-CA-C	-5.43	100.80	111.06
1	A	209	ILE	N-CA-CB	-5.19	107.46	112.65
1	A	130	PHE	CA-C-N	5.15	125.81	119.99
1	A	130	PHE	C-N-CA	5.15	125.81	119.99
1	A	51	TYR	CB-CA-C	-5.09	102.77	111.02

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	303	QUJ	Peptide
2	B	307	ZY9	Peptide
2	B	308	QVE	Peptide
2	B	310	ZY9	Peptide
2	B	311	QVS	Peptide
2	B	314	ZY9	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1778	0	1631	17	0
2	B	224	0	5	1	0
3	A	1	0	0	0	0
4	A	4	0	0	0	0
4	B	2	0	0	0	0
All	All	2009	0	1636	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:ASN:ND2	1:A:168:THR:C	2.66	0.54
1:A:95:PHE:CE2	1:A:118:LEU:HD13	2.42	0.54
1:A:88:TYR:CE2	1:A:124:ASN:HB2	2.44	0.53
1:A:66:PHE:CZ	1:A:95:PHE:CE1	2.97	0.52
1:A:82:GLY:HA2	1:A:190:TYR:OH	2.09	0.52
1:A:66:PHE:CZ	1:A:95:PHE:CD1	2.99	0.51
1:A:89:ARG:O	1:A:122:HIS:HA	2.13	0.48
1:A:114:TYR:CE2	1:A:214:PRO:HG3	2.49	0.47
1:A:29:SER:O	1:A:245:ARG:NH1	2.46	0.47
1:A:109:VAL:O	1:A:112:LYS:HB3	2.15	0.47
1:A:59:ILE:HD11	1:A:66:PHE:CD1	2.49	0.46
1:A:61:ASN:HD22	1:A:168:THR:C	2.24	0.46
1:A:124:ASN:OD1	1:A:124:ASN:C	2.59	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:133:ALA:O	1:A:139:GLY:HA3	2.16	0.45
1:A:149:VAL:HA	1:A:217:VAL:O	2.18	0.43
2:B:316:QUK:OB	2:B:316:QUK:N1	2.52	0.43
1:A:126:LYS:HD2	1:A:127:TYR:CZ	2.54	0.42
1:A:127:TYR:CZ	1:A:136:GLN:HG3	2.56	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	233/259 (90%)	204 (88%)	29 (12%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/224 (78%)	173 (99%)	1 (1%)	78	90

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

15 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	QVE	B	315	2	19,19,20	2.70	5 (26%)	25,26,28	2.71	8 (32%)
2	QUJ	B	303	2	19,19,20	2.43	3 (15%)	25,26,28	2.17	7 (28%)
2	A1IJ4	B	302	2	16,16,17	1.53	2 (12%)	17,18,20	2.09	5 (29%)
2	QDD	B	313	2	18,18,19	1.82	3 (16%)	24,25,27	2.69	9 (37%)
2	QVE	B	308	2	19,19,20	1.99	2 (10%)	25,26,28	2.09	7 (28%)
2	QVS	B	305	2	15,15,16	2.14	2 (13%)	20,21,23	2.47	7 (35%)
2	QUK	B	316	2	20,20,20	1.68	2 (10%)	27,27,27	1.69	6 (22%)
2	ZY9	B	307	2	10,10,11	3.95	4 (40%)	11,12,14	2.37	5 (45%)
2	QVS	B	311	2	15,15,16	1.67	2 (13%)	20,21,23	3.32	8 (40%)
2	ZY9	B	312	2	10,10,11	3.35	2 (20%)	11,12,14	3.31	5 (45%)
2	QUK	B	306	2	19,19,20	1.79	5 (26%)	24,25,27	2.78	7 (29%)
2	ZY9	B	314	2	10,10,11	3.09	2 (20%)	11,12,14	2.64	5 (45%)
2	ZY9	B	309	2	10,10,11	3.33	2 (20%)	11,12,14	2.57	7 (63%)
2	ZY9	B	310	2	10,10,11	5.32	5 (50%)	11,12,14	2.60	4 (36%)
2	ZY9	B	304	2	10,10,11	3.36	2 (20%)	11,12,14	2.72	5 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	QVE	B	315	2	-	2/7/7/9	0/2/2/2
2	QUJ	B	303	2	-	2/7/7/9	0/2/2/2
2	A1IJ4	B	302	2	-	2/10/10/11	0/1/1/1
2	QDD	B	313	2	-	0/6/6/8	0/2/2/2
2	QVE	B	308	2	-	3/7/7/9	0/2/2/2
2	QVS	B	305	2	-	0/2/2/4	0/2/2/2
2	QUK	B	316	2	-	5/9/9/9	0/2/2/2
2	ZY9	B	307	2	-	0/4/4/6	0/1/1/1
2	QVS	B	311	2	-	0/2/2/4	0/2/2/2
2	ZY9	B	312	2	-	0/4/4/6	0/1/1/1
2	QUK	B	306	2	-	2/7/7/9	0/2/2/2
2	ZY9	B	314	2	-	2/4/4/6	0/1/1/1
2	ZY9	B	309	2	-	0/4/4/6	0/1/1/1
2	ZY9	B	310	2	-	0/4/4/6	0/1/1/1
2	ZY9	B	304	2	-	0/4/4/6	0/1/1/1

All (43) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	310	ZY9	CA-C	-12.53	1.35	1.48
2	B	307	ZY9	C2-C7	-11.07	1.37	1.51
2	B	312	ZY9	C2-C7	-9.88	1.39	1.51
2	B	315	QVE	CA-C	-9.51	1.38	1.48
2	B	310	ZY9	C2-C7	-9.12	1.40	1.51
2	B	303	QUJ	CA-C	-8.71	1.39	1.48
2	B	309	ZY9	C2-C7	-8.71	1.40	1.51
2	B	314	ZY9	C2-C7	-8.68	1.40	1.51
2	B	304	ZY9	C2-C7	-7.65	1.41	1.51
2	B	308	QVE	CA-C	-7.06	1.41	1.48
2	B	304	ZY9	CA-C	-7.00	1.41	1.48
2	B	305	QVS	CA-C	-6.01	1.42	1.48
2	B	313	QDD	C10-C	-5.48	1.42	1.48
2	B	316	QUK	C10-C	-5.21	1.41	1.50
2	B	309	ZY9	CA-C	-5.15	1.43	1.48
2	B	311	QVS	CA-C	-4.65	1.43	1.48
2	B	310	ZY9	C8-C6	4.38	1.46	1.38
2	B	306	QUK	C10-C	-4.25	1.44	1.48
2	B	314	ZY9	CA-C	-3.89	1.44	1.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	302	A1IJ4	C128-C127	-3.82	1.37	1.51
2	B	315	QVE	CG-CD	3.82	1.62	1.50
2	B	303	QUJ	CA-N11	3.81	1.37	1.33
2	B	305	QVS	CA-N11	3.79	1.37	1.33
2	B	307	ZY9	CA-C	-3.72	1.44	1.48
2	B	316	QUK	OB-CG	3.67	1.56	1.43
2	B	310	ZY9	C7-N11	3.32	1.40	1.34
2	B	313	QDD	CB-C8	-3.05	1.42	1.52
2	B	312	ZY9	CA-C	-2.98	1.45	1.48
2	B	315	QVE	CA-N11	2.98	1.36	1.33
2	B	310	ZY9	C8-C9	2.78	1.43	1.38
2	B	306	QUK	C6-C7	-2.76	1.35	1.42
2	B	306	QUK	C10-N11	2.69	1.35	1.33
2	B	311	QVS	C6-C7	-2.56	1.35	1.42
2	B	308	QVE	CA-N11	2.53	1.35	1.33
2	B	313	QDD	C7-N11	-2.44	1.32	1.37
2	B	306	QUK	CA-C7	-2.41	1.38	1.42
2	B	315	QVE	C4-C5	2.29	1.41	1.36
2	B	307	ZY9	C9-CA	2.25	1.43	1.39
2	B	307	ZY9	C7-N11	2.20	1.38	1.34
2	B	306	QUK	C7-N11	-2.20	1.32	1.37
2	B	302	A1IJ4	C124-C127	-2.18	1.35	1.39
2	B	303	QUJ	C4-C5	2.07	1.41	1.36
2	B	315	QVE	C4-C3	2.06	1.42	1.38

All (95) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	311	QVS	CA-N11-C7	11.78	127.76	118.06
2	B	306	QUK	C10-N11-C7	8.32	124.91	118.06
2	B	313	QDD	C10-N11-C7	7.69	124.39	118.06
2	B	315	QVE	C-CA-N11	7.34	121.85	114.66
2	B	305	QVS	CA-N11-C7	7.13	123.93	118.06
2	B	310	ZY9	O-C-CA	-6.62	118.11	124.22
2	B	303	QUJ	C-CA-N11	6.25	120.78	114.66
2	B	306	QUK	C-C10-N11	6.05	120.59	114.66
2	B	312	ZY9	O-C-CA	-5.68	118.98	124.22
2	B	312	ZY9	CA-N11-C7	5.66	126.10	118.35
2	B	315	QVE	CA-N11-C7	5.66	122.72	118.06
2	B	308	QVE	C-CA-N11	5.46	120.01	114.66
2	B	311	QVS	C9-CA-N11	-5.43	116.93	123.18
2	B	313	QDD	C2-C7-C6	5.41	124.05	119.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	312	ZY9	C9-CA-N11	-5.26	115.21	122.44
2	B	304	ZY9	C2-C7-N11	5.06	124.68	115.85
2	B	302	A1IJ4	C122-C124-C127	4.94	125.16	119.76
2	B	306	QUK	C9-C10-N11	-4.79	117.67	123.18
2	B	314	ZY9	C2-C7-N11	4.72	124.10	115.85
2	B	315	QVE	C9-CA-N11	-4.63	117.85	123.18
2	B	303	QUJ	CA-N11-C7	4.43	121.71	118.06
2	B	303	QUJ	C7-C2-N	-4.29	110.20	118.16
2	B	306	QUK	CA-C7-C6	4.20	122.93	119.02
2	B	308	QVE	CA-N11-C7	4.17	121.49	118.06
2	B	315	QVE	C7-C2-N	-4.13	110.49	118.16
2	B	314	ZY9	C6-C7-N11	-4.10	117.22	122.40
2	B	312	ZY9	C2-C7-N11	4.02	122.87	115.85
2	B	315	QVE	O-C-CA	-3.99	120.54	124.22
2	B	304	ZY9	C6-C7-N11	-3.97	117.38	122.40
2	B	303	QUJ	C3-C2-N	3.96	128.25	120.39
2	B	304	ZY9	O-C-CA	-3.94	120.58	124.22
2	B	316	QUK	OXT-C-O	-3.94	114.89	123.35
2	B	315	QVE	C3-C2-N	3.91	128.14	120.39
2	B	313	QDD	C9-C10-N11	-3.87	118.72	123.18
2	B	307	ZY9	C9-CA-N11	-3.79	117.24	122.44
2	B	305	QVS	C9-CA-N11	-3.74	118.88	123.18
2	B	313	QDD	C3-C2-N	3.65	127.64	120.39
2	B	309	ZY9	C6-C7-N11	-3.55	117.91	122.40
2	B	314	ZY9	C8-C6-C7	3.48	123.29	118.93
2	B	316	QUK	C10-N11-C7	3.46	124.75	117.41
2	B	311	QVS	C2-C7-N11	3.45	122.18	118.55
2	B	309	ZY9	O-C-CA	-3.44	121.05	124.22
2	B	313	QDD	O-C-C10	-3.38	121.10	124.22
2	B	307	ZY9	CA-N11-C7	3.36	122.95	118.35
2	B	309	ZY9	C2-C7-N11	3.36	121.71	115.85
2	B	308	QVE	CG-OB-C8	-3.27	109.73	116.68
2	B	316	QUK	C3-CA-N	3.25	126.85	120.39
2	B	310	ZY9	CA-N11-C7	3.24	122.78	118.35
2	B	309	ZY9	C8-C6-C7	3.22	122.96	118.93
2	B	307	ZY9	C2-C7-N11	3.21	121.45	115.85
2	B	310	ZY9	C6-C7-N11	-3.21	118.35	122.40
2	B	314	ZY9	CA-N11-C7	3.20	122.72	118.35
2	B	308	QVE	C3-C2-N	3.14	126.62	120.39
2	B	308	QVE	OB-CG-CD	-3.13	100.08	110.40
2	B	306	QUK	O-C-C10	-3.08	121.38	124.22
2	B	311	QVS	C6-C7-N11	-3.06	116.42	122.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	316	QUK	C7-CA-N	-3.05	112.49	118.16
2	B	305	QVS	C4-C5-C6	-3.05	116.79	120.91
2	B	307	ZY9	O-C-CA	-3.04	121.42	124.22
2	B	308	QVE	C9-CA-N11	-3.01	119.71	123.18
2	B	302	A1IJ4	C121-O71-C122	2.98	125.68	117.93
2	B	309	ZY9	CA-N11-C7	2.96	122.40	118.35
2	B	304	ZY9	CA-N11-C7	2.93	122.35	118.35
2	B	311	QVS	C9-CA-C	2.74	126.03	121.11
2	B	309	ZY9	C9-CA-N11	-2.73	118.69	122.44
2	B	308	QVE	C7-C2-N	-2.72	113.12	118.16
2	B	305	QVS	C-CA-N11	2.71	117.31	114.66
2	B	302	A1IJ4	C125-C126-C127	-2.68	116.85	120.61
2	B	316	QUK	C9-C10-N11	-2.67	117.69	124.31
2	B	303	QUJ	C9-CA-N11	-2.65	120.12	123.18
2	B	311	QVS	C2-C7-C6	2.55	121.39	119.02
2	B	305	QVS	C3-C2-N	2.55	125.44	120.39
2	B	302	A1IJ4	C126-C125-C123	2.54	123.51	120.24
2	B	314	ZY9	O-C-CA	-2.48	121.93	124.22
2	B	312	ZY9	C6-C7-N11	-2.47	119.28	122.40
2	B	311	QVS	C-CA-N11	2.43	117.04	114.66
2	B	313	QDD	C2-C7-N11	-2.42	116.01	118.55
2	B	309	ZY9	C6-C8-C9	-2.41	117.15	120.24
2	B	302	A1IJ4	O-C-N31	-2.32	122.21	125.03
2	B	303	QUJ	C4-C3-C2	-2.30	116.92	121.19
2	B	315	QVE	OE2-CD-CG	-2.29	112.45	121.99
2	B	313	QDD	C9-C8-C6	-2.27	116.46	118.87
2	B	313	QDD	OD1-CG-CB	-2.23	116.23	122.94
2	B	304	ZY9	C8-C6-C7	2.22	121.71	118.93
2	B	305	QVS	C2-C7-C6	2.22	121.08	119.02
2	B	306	QUK	C3-CA-N	2.20	124.75	120.39
2	B	310	ZY9	C2-C7-N11	2.17	119.64	115.85
2	B	316	QUK	OXT-C-C10	2.15	119.83	114.71
2	B	313	QDD	C-C10-N11	2.14	116.76	114.66
2	B	303	QUJ	O-C-CA	-2.14	122.25	124.22
2	B	306	QUK	C6-C7-N11	-2.14	118.29	122.66
2	B	315	QVE	C2-C7-N11	2.08	120.74	118.55
2	B	311	QVS	C3-C2-N	2.06	124.47	120.39
2	B	305	QVS	C6-C7-N11	-2.04	118.49	122.66
2	B	307	ZY9	C6-C7-N11	-2.03	119.83	122.40

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	314	ZY9	O-C-CA-N11
2	B	314	ZY9	O-C-CA-C9
2	B	316	QUK	O-C-C10-N11
2	B	316	QUK	O-C-C10-C9
2	B	316	QUK	OXT-C-C10-N11
2	B	316	QUK	OXT-C-C10-C9
2	B	308	QVE	C9-C8-OB-CG
2	B	308	QVE	C6-C8-OB-CG
2	B	316	QUK	CE-CD-CG-OB
2	B	303	QUJ	C6-C8-OB-CG
2	B	306	QUK	CE-CD-CG-OB
2	B	303	QUJ	C9-C8-OB-CG
2	B	315	QVE	C9-C8-OB-CG
2	B	315	QVE	C6-C8-OB-CG
2	B	302	A1IJ4	C119-C120-C121-O71
2	B	306	QUK	CG-CD-CE-N1
2	B	302	A1IJ4	C119-C118-N31-C
2	B	308	QVE	CD-CG-OB-C8

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	316	QUK	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/259 (91%)	0.24	4 (1%) 69 66	45, 78, 124, 142	0
2	B	0/16	-	-	-	-
All	All	237/275 (86%)	0.24	4 (1%) 69 66	45, 78, 124, 142	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	95	PHE	2.8
1	A	183	LEU	2.8
1	A	149	VAL	2.4
1	A	36	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	QUK	B	316	19/19	0.89	0.11	66,70,86,86	0
2	QDD	B	313	17/18	0.91	0.10	59,61,69,72	0
2	QVE	B	315	18/19	0.92	0.09	60,62,73,75	0
2	QVE	B	308	18/19	0.92	0.10	57,60,81,86	0
2	QUK	B	306	18/19	0.93	0.10	57,60,70,71	0
2	QUJ	B	303	18/19	0.94	0.08	55,59,77,78	0
2	ZY9	B	314	10/11	0.94	0.08	62,64,65,65	0
2	ZY9	B	304	10/11	0.94	0.09	52,53,55,55	0
2	ZY9	B	310	10/11	0.94	0.09	62,64,65,65	0
2	QVS	B	311	14/15	0.95	0.08	57,61,64,64	0
2	ZY9	B	312	10/11	0.95	0.06	55,57,60,60	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	QVS	B	305	14/15	0.95	0.06	54,55,57,58	0
2	ZY9	B	309	10/11	0.96	0.07	57,58,60,60	0
2	ZY9	B	307	10/11	0.96	0.08	53,56,58,58	0
2	A1IJ4	B	302	16/17	0.97	0.07	52,55,61,63	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ZN	A	301	1/1	0.97	0.06	87,87,87,87	0

6.5 Other polymers [i](#)

There are no such residues in this entry.