



wwPDB EM Validation Summary Report ⓘ

Mar 15, 2026 – 12:51 PM UTC

PDB ID : 9Q90 / pdb_00009q90
EMDB ID : EMD-14171
Title : CryoEM structure of bacterial transcription intermediate complex mediated by activator PspF
Authors : Ye, F.; Gao, F.; Zhang, X.
Deposited on : 2025-02-26
Resolution : 3.50 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	13950 (3.00 - 4.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1	275	
1	2	275	
1	3	275	
1	4	275	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	5	275	
1	6	275	
2	A	329	
2	B	329	
3	C	1342	
4	D	1407	
5	E	91	
6	M	497	
7	N	36	
8	T	36	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	AF3	3	602	-	-	X	-
10	AF3	4	602	-	-	X	-

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 41495 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Psp operon transcriptional activator.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	1	258	Total	C	N	O	S	0	0
			2052	1302	360	380	10		
1	2	259	Total	C	N	O	S	0	0
			2052	1301	361	380	10		
1	3	259	Total	C	N	O	S	0	0
			2053	1304	358	381	10		
1	4	259	Total	C	N	O	S	0	0
			2062	1308	364	381	9		
1	5	256	Total	C	N	O	S	0	0
			2041	1295	361	376	9		
1	6	256	Total	C	N	O	S	0	0
			2000	1270	349	372	9		

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	A	309	Total	C	N	O	S	0	0
			2322	1453	407	455	7		
2	B	223	Total	C	N	O	S	0	0
			1676	1045	294	332	5		

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	1341	Total	C	N	O	S	0	0
			10125	6360	1761	1964	40		

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	1334	Total	C	N	O	S	0	0
			9634	6052	1730	1814	38		

- Molecule 5 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	74	Total	C	N	O	S	0	0
			546	335	109	101	1		

- Molecule 6 is a protein called RNA polymerase sigma-54 factor.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	M	417	Total	C	N	O	S	0	0
			3301	2071	572	646	12		

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	-19	MET	-	initiating methionine	UNP A6TEM1
M	-18	GLY	-	expression tag	UNP A6TEM1
M	-17	SER	-	expression tag	UNP A6TEM1
M	-16	SER	-	expression tag	UNP A6TEM1
M	-15	HIS	-	expression tag	UNP A6TEM1
M	-14	HIS	-	expression tag	UNP A6TEM1
M	-13	HIS	-	expression tag	UNP A6TEM1
M	-12	HIS	-	expression tag	UNP A6TEM1
M	-11	HIS	-	expression tag	UNP A6TEM1
M	-10	HIS	-	expression tag	UNP A6TEM1
M	-9	SER	-	expression tag	UNP A6TEM1
M	-8	SER	-	expression tag	UNP A6TEM1
M	-7	GLY	-	expression tag	UNP A6TEM1
M	-6	LEU	-	expression tag	UNP A6TEM1
M	-5	VAL	-	expression tag	UNP A6TEM1
M	-4	PRO	-	expression tag	UNP A6TEM1
M	-3	ARG	-	expression tag	UNP A6TEM1
M	-2	GLY	-	expression tag	UNP A6TEM1
M	-1	SER	-	expression tag	UNP A6TEM1
M	0	HIS	-	expression tag	UNP A6TEM1
M	1	MET	-	expression tag	UNP A6TEM1
M	2	LYS	-	expression tag	UNP A6TEM1
M	3	GLN	-	expression tag	UNP A6TEM1
M	4	GLY	-	expression tag	UNP A6TEM1
M	5	LEU	-	expression tag	UNP A6TEM1
M	6	GLN	-	expression tag	UNP A6TEM1
M	7	LEU	-	expression tag	UNP A6TEM1
M	8	ARG	-	expression tag	UNP A6TEM1
M	9	LEU	-	expression tag	UNP A6TEM1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
M	10	SER	-	expression tag	UNP A6TEM1
M	11	GLN	-	expression tag	UNP A6TEM1
M	12	GLN	-	expression tag	UNP A6TEM1
M	13	LEU	-	expression tag	UNP A6TEM1
M	14	ALA	-	expression tag	UNP A6TEM1


- Molecule 7 is a DNA chain called DNA (36-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
7	N	36	Total	C	N	O	P	0	0
			738	349	137	216	36		

- Molecule 8 is a DNA chain called DNA (36-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
8	T	36	Total	C	N	O	P	0	0
			738	349	137	216	36		


- Molecule 9 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
9	3	1	Total	C	N	O	P	0
			27	10	5	10	2	
9	4	1	Total	C	N	O	P	0
			27	10	5	10	2	
9	6	1	Total	C	N	O	P	0
			27	10	5	10	2	

- Molecule 10 is ALUMINUM FLUORIDE (CCD ID: AF3) (formula: AlF_3).



Mol	Chain	Residues	Atoms			AltConf
10	1	1	Total	Al	F	0
			4	1	3	
10	2	1	Total	Al	F	0
			4	1	3	
10	3	1	Total	Al	F	0
			4	1	3	
10	4	1	Total	Al	F	0
			4	1	3	

- Molecule 11 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
11	1	1	Total	Mg	0
			1	1	
11	2	1	Total	Mg	0
			1	1	

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
11	3	1	Total 1	Mg 1	0
11	4	1	Total 1	Mg 1	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Psp operon transcriptional activator




- Molecule 1: Psp operon transcriptional activator




- Molecule 1: Psp operon transcriptional activator









• Molecule 1: Psp operon transcriptional activator




• Molecule 1: Psp operon transcriptional activator




• Molecule 1: Psp operon transcriptional activator






• Molecule 2: DNA-directed RNA polymerase subunit alpha

Chain A: 76% 18% 6%




• Molecule 2: DNA-directed RNA polymerase subunit alpha


Chain B: 47% 21% 32%



• Molecule 3: DNA-directed RNA polymerase subunit beta


Chain C: 81% 19%






• Molecule 4: DNA-directed RNA polymerase subunit beta'


Chain D: 74% 21% 5%






• Molecule 5: DNA-directed RNA polymerase subunit omega

Chain E: 71% 10% 19%




• Molecule 6: RNA polymerase sigma-54 factor

Chain M: 58% 25% 16%



• Molecule 7: DNA (36-MER)

Chain N: 33% 67%



• Molecule 8: DNA (36-MER)

Chain T: 58% 42%



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	23354	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	1400	Depositor
Maximum defocus (nm)	2600	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.302	Depositor
Minimum map value	-0.152	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.006	Depositor
Recommended contour level	0.012	Depositor
Map size (Å)	308.0, 308.0, 308.0	wwPDB
Map dimensions	280, 280, 280	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.1, 1.1, 1.1	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AF3, ADP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	1	0.37	0/2093	0.63	0/2832
1	2	0.19	0/2093	0.47	0/2833
1	3	0.21	0/2095	0.49	0/2837
1	4	0.19	0/2104	0.45	0/2848
1	5	0.19	0/2082	0.51	1/2817 (0.0%)
1	6	0.25	0/2038	0.55	0/2762
2	A	0.15	0/2351	0.39	0/3202
2	B	0.16	0/1693	0.41	0/2300
3	C	0.15	0/10283	0.39	2/13940 (0.0%)
4	D	0.15	0/9768	0.38	2/13270 (0.0%)
5	E	0.17	0/547	0.46	0/740
6	M	0.28	2/3348 (0.1%)	0.55	6/4533 (0.1%)
7	N	0.16	0/827	0.38	0/1274
8	T	0.17	0/827	0.35	0/1274
All	All	0.19	2/42149 (0.0%)	0.44	11/57462 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	1	0	1
6	M	0	1
All	All	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	M	269	PRO	CG-CD	-10.51	1.15	1.50
6	M	269	PRO	N-CD	5.80	1.55	1.47

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	M	269	PRO	CA-N-CD	-12.57	94.41	112.00
6	M	269	PRO	N-CD-CG	-11.47	86.00	103.20
3	C	535	PRO	N-CD-CG	-6.69	93.16	103.20
6	M	269	PRO	CA-CB-CG	-6.51	92.12	104.50
4	D	506	VAL	N-CA-C	-6.49	106.67	112.12

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	1	227	ARG	Sidechain
6	M	1	MET	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	2052	0	2048	100	0
1	2	2052	0	2048	84	0
1	3	2053	0	2044	67	0
1	4	2062	0	2059	90	0
1	5	2041	0	2038	103	0
1	6	2000	0	1967	83	0
2	A	2322	0	2308	45	0
2	B	1676	0	1676	46	0
3	C	10125	0	9829	185	0
4	D	9634	0	9244	208	0
5	E	546	0	548	10	0
6	M	3301	0	3331	108	0
7	N	738	0	404	20	0
8	T	738	0	404	16	0
9	1	27	0	12	2	0
9	2	27	0	12	3	0
9	3	27	0	12	4	0
9	4	27	0	12	3	0
9	6	27	0	12	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	1	4	0	0	0	0
10	2	4	0	0	0	0
10	3	4	0	0	2	0
10	4	4	0	0	3	0
11	1	1	0	0	0	0
11	2	1	0	0	0	0
11	3	1	0	0	0	0
11	4	1	0	0	0	0
All	All	41495	0	40008	1059	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 1059 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:3:42:LYS:NZ	10:3:602:AF3:F3	1.84	1.00
2:A:10:LYS:HD2	2:B:227:GLN:HA	1.52	0.89
1:1:81:GLU:HA	1:1:91:ARG:HE	1.42	0.85
3:C:903:ARG:HD3	3:C:910:ALA:HB2	1.58	0.83
1:2:38:ARG:HH12	1:2:224:GLY:HA2	1.43	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1	256/275 (93%)	242 (94%)	14 (6%)	0	100	100
1	2	257/275 (94%)	248 (96%)	9 (4%)	0	100	100
1	3	257/275 (94%)	246 (96%)	11 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	4	257/275 (94%)	247 (96%)	10 (4%)	0	100	100
1	5	254/275 (92%)	243 (96%)	11 (4%)	0	100	100
1	6	254/275 (92%)	244 (96%)	10 (4%)	0	100	100
2	A	305/329 (93%)	280 (92%)	25 (8%)	0	100	100
2	B	219/329 (67%)	207 (94%)	12 (6%)	0	100	100
3	C	1339/1342 (100%)	1261 (94%)	78 (6%)	0	100	100
4	D	1322/1407 (94%)	1221 (92%)	101 (8%)	0	100	100
5	E	72/91 (79%)	69 (96%)	3 (4%)	0	100	100
6	M	413/497 (83%)	397 (96%)	16 (4%)	0	100	100
All	All	5205/5645 (92%)	4905 (94%)	300 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	219/235 (93%)	213 (97%)	6 (3%)	39	62
1	2	219/235 (93%)	217 (99%)	2 (1%)	70	76
1	3	219/235 (93%)	219 (100%)	0	100	100
1	4	220/235 (94%)	219 (100%)	1 (0%)	81	80
1	5	218/235 (93%)	217 (100%)	1 (0%)	81	80
1	6	208/235 (88%)	208 (100%)	0	100	100
2	A	249/286 (87%)	248 (100%)	1 (0%)	84	81
2	B	180/286 (63%)	179 (99%)	1 (1%)	78	79
3	C	1047/1157 (90%)	1045 (100%)	2 (0%)	87	85
4	D	913/1168 (78%)	911 (100%)	2 (0%)	87	85
5	E	53/75 (71%)	53 (100%)	0	100	100
6	M	367/440 (83%)	365 (100%)	2 (0%)	81	80

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	4112/4822 (85%)	4094 (100%)	18 (0%)	81	81

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	144	TYR
6	M	285	TRP
6	M	42	GLU
1	4	43	GLU
3	C	737	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	965	GLN
6	M	12	GLN
4	D	200	GLN
6	M	20	GLN
4	D	805	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	ADP	6	301	-	28,29,29	1.41	4 (14%)	43,45,45	1.85	9 (20%)
9	ADP	4	601	10,11	28,29,29	1.41	4 (14%)	43,45,45	1.84	11 (25%)
10	AF3	1	602	9	0,3,3	-	-	-		
10	AF3	2	602	9	0,3,3	-	-	-		
9	ADP	2	601	10,11	28,29,29	1.41	4 (14%)	43,45,45	1.86	8 (18%)
10	AF3	4	602	9	0,3,3	-	-	-		
9	ADP	1	601	10,11	28,29,29	1.40	4 (14%)	43,45,45	1.88	8 (18%)
9	ADP	3	601	10,11	28,29,29	1.41	4 (14%)	43,45,45	1.83	8 (18%)
10	AF3	3	602	9	0,3,3	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	ADP	6	301	-	-	2/16/32/32	0/3/3/3
9	ADP	4	601	10,11	-	1/16/32/32	0/3/3/3
9	ADP	2	601	10,11	-	4/16/32/32	0/3/3/3
9	ADP	1	601	10,11	-	0/16/32/32	0/3/3/3
9	ADP	3	601	10,11	-	2/16/32/32	0/3/3/3

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	3	601	ADP	C5-C4	4.74	1.47	1.39
9	2	601	ADP	C5-C4	4.72	1.47	1.39
9	1	601	ADP	C5-C4	4.70	1.47	1.39
9	4	601	ADP	C5-C4	4.69	1.47	1.39
9	6	301	ADP	C5-C4	4.69	1.47	1.39

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	2	601	ADP	C5-C4-N3	-6.00	118.46	126.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	3	601	ADP	C5-C4-N3	-5.98	118.48	126.72
9	1	601	ADP	C5-C4-N3	-5.97	118.50	126.72
9	4	601	ADP	C5-C4-N3	-5.87	118.63	126.72
9	6	301	ADP	C5-C4-N3	-5.87	118.63	126.72

There are no chirality outliers.

5 of 9 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
9	2	601	ADP	PA-O3A-PB-O3B
9	2	601	ADP	C5'-O5'-PA-O1A
9	2	601	ADP	C5'-O5'-PA-O3A
9	3	601	ADP	C5'-O5'-PA-O3A
9	6	301	ADP	C5'-O5'-PA-O1A



There are no ring outliers.


7 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	6	301	ADP	1	0
9	4	601	ADP	3	0
9	2	601	ADP	3	0
10	4	602	AF3	3	0
9	1	601	ADP	2	0
9	3	601	ADP	4	0
10	3	602	AF3	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-14171. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 140



Y Index: 140




Z Index: 140


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 173



Y Index: 111




Z Index: 149


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.012. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 723 nm³; this corresponds to an approximate mass of 653 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.286 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-14171 and PDB model 9Q90. Per-residue inclusion information can be found in section [3](#) on page [9](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.012 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.012).

9.4 Atom inclusion [i](#)



At the recommended contour level, 97% of all backbone atoms, 95% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.012) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9530	0.3560
1	0.9450	0.1570
2	0.9070	0.1390
3	0.9020	0.1270
4	0.9330	0.1320
5	0.9100	0.1300
6	0.8820	0.1290
A	0.9240	0.4430
B	0.9660	0.4290
C	0.9760	0.4740
D	0.9740	0.4590
E	0.9850	0.5050
M	0.9680	0.4220
N	0.9860	0.3140
T	0.9780	0.3020

