



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 03:07 AM UTC

PDB ID : 2QA6 / pdb_00002qa6
Title : Crystal Structure of Estrogen Receptor Alpha mutant 537S Complexed with 4-(6-hydroxy-1H-indazol-3-yl)benzene-1,3-diol
Authors : Nettles, K.W.; Bruning, J.B.; Nowak, J.; Sharma, S.K.; Hahm, J.B.; Shi, Y.; Kulp, K.; Hochberg, R.B.; Zhou, H.; Katzenellenbogen, J.A.; Katzenellenbogen, B.S.; Kim, Y.; Joachmiak, A.; Greene, G.L.
Deposited on : 2007-06-14
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	
1	B	258	
2	C	13	
2	D	13	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4062 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Estrogen receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	243	1906	1222	325	340	19	0	0	0
1	B	242	1912	1226	326	341	19	0	0	0


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	297	SER	-	expression tag	UNP P03372
A	537	SER	TYR	engineered mutation	UNP P03372
B	297	SER	-	expression tag	UNP P03372
B	537	SER	TYR	engineered mutation	UNP P03372

- Molecule 2 is a protein called nuclear receptor coactivator 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	9	79	51	16	12	0	0	0
2	D	11	88	56	18	14	0	0	0

- Molecule 3 is 4-(6-HYDROXY-1H-INDAZOL-3-YL)BENZENE-1,3-DIOL (CCD ID: KN2) (formula: C₁₃H₁₀N₂O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
3	A	1	Total	C	N	O	0	0
			18	13	2	3		
3	B	1	Total	C	N	O	0	0
			18	13	2	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	21	Total	O	0	0
			21	21		
4	B	20	Total	O	0	0
			20	20		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Estrogen receptor



- Molecule 1: Estrogen receptor



- Molecule 2: nuclear receptor coactivator 2



- Molecule 2: nuclear receptor coactivator 2





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.01Å 82.94Å 58.58Å 90.00° 109.25° 90.00°	Depositor
Resolution (Å)	15.00 – 2.60 15.00 – 2.60	Depositor EDS
% Data completeness (in resolution range)	91.7 (15.00-2.60) 91.2 (15.00-2.60)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.06 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.195 , 0.249 0.197 , 0.254	Depositor DCC
R_{free} test set	694 reflections (4.46%)	wwPDB-VP
Wilson B-factor (Å ²)	65.0	Xtrriage
Anisotropy	0.123	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 45.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.029 for l,-k,h	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4062	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: KN2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.93	2/1940 (0.1%)	1.00	1/2622 (0.0%)
1	B	1.01	0/1947	1.05	4/2632 (0.2%)
2	C	0.67	0/79	1.23	2/104 (1.9%)
2	D	0.78	0/88	1.06	1/116 (0.9%)
All	All	0.96	2/4054 (0.0%)	1.03	8/5474 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	435	PHE	CA-C	-5.61	1.45	1.52
1	A	452	ILE	CA-C	5.39	1.59	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	512	SER	N-CA-C	-8.13	102.50	111.36
2	D	690	LEU	N-CA-C	-6.25	104.39	111.14
2	C	689	ILE	CB-CA-C	-6.20	103.66	112.22
2	C	690	LEU	N-CA-C	-5.48	105.22	111.14
1	A	331	TYR	N-CA-C	5.47	118.40	107.62
1	B	377	HIS	N-CA-C	5.38	117.56	111.11
1	B	382	ALA	N-CA-C	5.11	119.67	113.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	458	VAL	N-CA-C	5.08	115.80	110.62

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	330	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1906	0	1940	55	0
1	B	1912	0	1947	35	0
2	C	79	0	88	3	0
2	D	88	0	91	2	0
3	A	18	0	10	2	0
3	B	18	0	8	1	0
4	A	21	0	0	4	0
4	B	20	0	0	4	0
All	All	4062	0	4084	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:465:THR:N	1:B:468:SER:HG	1.82	0.77
1:A:501:HIS:HB2	4:A:569:HOH:O	1.85	0.75
1:B:547:HIS:O	1:B:548:ARG:C	2.34	0.69
1:B:392:VAL:HG11	1:B:431:THR:HG22	1.73	0.69
1:B:366:GLY:HA2	1:B:369:ASP:OD2	1.93	0.69
1:A:472:LYS:HE3	4:A:558:HOH:O	1.94	0.66
1:A:435:PHE:HE1	1:A:510:ILE:HG21	1.61	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:519:ASN:OD1	1:B:519:ASN:ND2	2.31	0.62
1:A:421:MET:HB3	1:A:524:HIS:CE1	2.34	0.62
1:B:472:LYS:HE2	4:B:566:HOH:O	1.99	0.62
1:A:392:VAL:HG13	1:A:432:SER:HA	1.82	0.61
1:B:547:HIS:O	1:B:549:LEU:HD13	2.00	0.61
1:A:331:TYR:HB3	4:A:575:HOH:O	1.98	0.61
1:A:418:VAL:HG12	1:A:528:MET:CE	2.30	0.61
1:B:331:TYR:CD1	1:B:333:PRO:HD3	2.35	0.61
1:A:358:ILE:HG23	2:C:694:LEU:HD23	1.83	0.60
1:B:522:MET:HE3	1:B:547:HIS:CE1	2.39	0.58
1:A:536:LEU:HB2	1:A:541:LEU:HD13	1.85	0.58
1:A:401:LYS:HD3	1:A:409:LEU:HD21	1.86	0.58
1:A:342:MET:HA	1:A:342:MET:HE2	1.86	0.57
1:B:321:ASP:CB	4:B:562:HOH:O	2.51	0.57
1:A:413:ASN:ND2	1:A:413:ASN:H	2.03	0.56
1:A:533:VAL:HG12	1:A:534:VAL:HG22	1.87	0.56
1:B:486:LEU:O	1:B:490:MET:HG3	2.06	0.56
1:B:472:LYS:CE	4:B:566:HOH:O	2.53	0.56
1:A:355:VAL:O	1:A:359:ASN:ND2	2.39	0.55
1:A:490:MET:HB3	1:A:495:LEU:HD22	1.89	0.55
1:A:310:LEU:O	1:A:481:LYS:HE2	2.07	0.54
1:A:427:MET:HE3	1:A:517:MET:HG2	1.89	0.54
1:B:465:THR:N	1:B:468:SER:OG	2.41	0.53
1:A:383:TRP:CD1	1:A:543:MET:HE3	2.44	0.52
1:A:392:VAL:HG13	1:A:432:SER:CA	2.40	0.52
1:B:342:MET:HG2	1:B:418:VAL:CG2	2.40	0.52
1:A:374:ASP:OD2	1:A:471:GLU:HG2	2.10	0.52
1:B:389:ILE:HD12	1:B:452:ILE:HD11	1.92	0.52
1:A:335:ARG:N	4:A:556:HOH:O	2.44	0.51
3:A:1:KN2:HAE	3:A:1:KN2:HAG	1.92	0.51
1:B:461:PHE:HB2	1:B:472:LYS:NZ	2.26	0.50
1:B:536:LEU:HB3	1:B:541:LEU:HD13	1.93	0.50
1:A:329:SER:O	1:A:330:GLU:HG3	2.10	0.50
1:B:392:VAL:HG11	1:B:431:THR:CG2	2.40	0.50
1:A:514:ILE:HA	1:A:517:MET:CE	2.42	0.50
1:A:424:ILE:HA	1:A:427:MET:HE2	1.94	0.50
1:A:427:MET:HA	4:B:572:HOH:O	2.12	0.50
1:A:329:SER:OG	1:A:330:GLU:N	2.46	0.49
2:D:689:ILE:HG22	2:D:693:LEU:HD11	1.94	0.48
1:A:370:LEU:HD21	1:A:475:ILE:HG13	1.96	0.48
1:A:435:PHE:CE1	1:A:510:ILE:HG21	2.45	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:392:VAL:HG11	1:A:431:THR:HG22	1.95	0.48
1:B:312:ALA:O	1:B:316:VAL:HG23	2.13	0.48
1:A:320:LEU:HD23	1:A:446:VAL:HG11	1.95	0.48
1:A:418:VAL:CG1	1:A:528:MET:HE3	2.43	0.48
1:B:549:LEU:HD22	1:B:549:LEU:H	1.79	0.47
1:B:412:ARG:NH2	1:B:422:VAL:HG22	2.29	0.47
1:A:343:MET:HE1	3:A:1:KN2:HAH	1.95	0.46
1:B:440:LEU:HD13	1:B:507:LEU:HD22	1.97	0.46
1:A:371:THR:HG21	1:A:467:LYS:HE2	1.96	0.46
1:A:455:ASN:O	1:A:458:VAL:HG12	2.16	0.46
1:B:331:TYR:CE1	1:B:333:PRO:HB3	2.50	0.46
1:A:337:PHE:CD2	1:A:342:MET:HE3	2.51	0.46
1:B:332:ASP:OD1	1:B:334:THR:OG1	2.33	0.46
1:B:395:SER:O	1:B:396:MET:C	2.59	0.46
1:A:418:VAL:CG1	1:A:528:MET:CE	2.93	0.46
2:D:689:ILE:O	2:D:693:LEU:HG	2.15	0.46
1:B:461:PHE:HB2	1:B:472:LYS:HZ1	1.81	0.45
1:A:329:SER:O	1:A:330:GLU:CG	2.64	0.45
1:A:343:MET:HB3	1:A:534:VAL:HG22	1.98	0.45
1:A:533:VAL:HG12	1:A:534:VAL:CG2	2.46	0.45
3:B:1:KN2:HAG	3:B:1:KN2:HAE	1.98	0.45
1:A:514:ILE:HA	1:A:517:MET:HE2	1.99	0.44
1:B:424:ILE:HA	1:B:427:MET:HE3	1.99	0.44
1:A:370:LEU:O	1:A:375:GLN:NE2	2.41	0.43
1:A:424:ILE:HD13	1:A:521:GLY:HA2	2.00	0.43
1:A:408:LEU:HG	1:A:410:LEU:HD21	2.01	0.43
1:B:536:LEU:CB	1:B:541:LEU:HD13	2.48	0.43
1:A:358:ILE:O	1:A:362:LYS:HG3	2.18	0.43
1:A:438:MET:HE3	1:A:506:GLN:OE1	2.18	0.43
1:A:474:HIS:HD2	1:A:477:ARG:HH21	1.65	0.43
1:A:539:LEU:O	1:A:543:MET:HG2	2.19	0.43
1:A:538:ASP:HB3	2:C:689:ILE:HD11	2.01	0.43
1:A:529:LYS:HD2	1:A:529:LYS:C	2.44	0.42
1:A:448:LEU:HD22	1:A:511:LEU:HD21	2.01	0.42
1:B:364:VAL:O	1:B:365:PRO:C	2.61	0.42
1:B:366:GLY:O	1:B:369:ASP:HB2	2.20	0.42
1:B:357:MET:HE2	1:B:386:ILE:HG22	2.01	0.42
1:A:402:LEU:HD12	1:A:425:PHE:CE2	2.55	0.42
2:C:695:GLN:O	2:C:696:ASP:C	2.62	0.41
1:B:438:MET:O	1:B:439:ASN:C	2.62	0.41
1:A:429:LEU:HA	1:A:429:LEU:HD23	1.79	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:383:TRP:CE2	1:B:543:MET:HB3	2.55	0.41
1:B:412:ARG:HA	1:B:425:PHE:CE2	2.56	0.41
1:A:421:MET:HE3	1:A:421:MET:HB2	1.96	0.41
1:A:402:LEU:HD12	1:A:425:PHE:CZ	2.55	0.40
1:B:424:ILE:HD13	1:B:521:GLY:HA2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/258 (93%)	227 (95%)	10 (4%)	2 (1%)	16	34
1	B	238/258 (92%)	228 (96%)	6 (2%)	4 (2%)	7	15
2	C	7/13 (54%)	7 (100%)	0	0	100	100
2	D	9/13 (69%)	8 (89%)	1 (11%)	0	100	100
All	All	493/542 (91%)	470 (95%)	17 (3%)	6 (1%)	10	23

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	548	ARG
1	A	329	SER
1	A	533	VAL
1	B	330	GLU
1	B	416	LYS
1	B	336	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	211/233 (91%)	194 (92%)	17 (8%)	11	24
1	B	212/233 (91%)	196 (92%)	16 (8%)	12	28
2	C	9/13 (69%)	7 (78%)	2 (22%)	1	2
2	D	9/13 (69%)	9 (100%)	0	100	100
All	All	441/492 (90%)	406 (92%)	35 (8%)	11	26

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	330	GLU
1	A	331	TYR
1	A	372	LEU
1	A	413	ASN
1	A	421	MET
1	A	425	PHE
1	A	437	MET
1	A	462	LEU
1	A	465	THR
1	A	467	LYS
1	A	468	SER
1	A	525	LEU
1	A	529	LYS
1	A	532	ASN
1	A	534	VAL
1	A	541	LEU
1	A	542	GLU
1	B	309	SER
1	B	337	PHE
1	B	397	GLU
1	B	432	SER
1	B	433	SER
1	B	460	THR
1	B	468	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	481	LYS
1	B	492	LYS
1	B	497	LEU
1	B	518	SER
1	B	525	LEU
1	B	536	LEU
1	B	540	LEU
1	B	541	LEU
1	B	549	LEU
2	C	693	LEU
2	C	696	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	413	ASN
1	A	474	HIS
1	A	476	HIS
1	A	501	HIS
1	A	513	HIS
1	A	519	ASN
1	B	455	ASN
1	B	488	HIS
1	B	501	HIS
1	B	519	ASN
1	B	532	ASN
2	D	695	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	KN2	A	1	-	20,20,20	1.87	3 (15%)	27,29,29	1.85	3 (11%)
3	KN2	B	1	-	20,20,20	1.95	4 (20%)	27,29,29	1.94	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KN2	A	1	-	-	2/4/4/4	0/3/3/3
3	KN2	B	1	-	-	2/4/4/4	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1	KN2	CAO-CAP	-5.55	1.39	1.49
3	B	1	KN2	N1K-N1J	-5.46	1.23	1.36
3	A	1	KN2	N1K-N1J	-4.82	1.24	1.36
3	B	1	KN2	CAO-CAP	-4.73	1.40	1.49
3	A	1	KN2	CAR-CAP	-2.86	1.40	1.47
3	B	1	KN2	CAR-CAP	-2.75	1.40	1.47
3	B	1	KN2	CAQ-N1K	-2.20	1.32	1.36

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1	KN2	CAR-CAP-N1J	-8.00	105.79	110.36
3	A	1	KN2	CAR-CAP-N1J	-7.86	105.87	110.36
3	A	1	KN2	CAP-N1J-N1K	2.50	112.32	106.36
3	B	1	KN2	CAP-N1J-N1K	2.47	112.25	106.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	B	1	KN2	CAG-CAR-CAP	2.21	138.48	132.41
3	A	1	KN2	CAQ-N1K-N1J	-2.08	110.36	111.90
3	B	1	KN2	CAE-CAD-CAL	2.01	122.00	119.88

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1	KN2	CAN-CAO-CAP-CAR
3	A	1	KN2	CAE-CAO-CAP-CAR
3	B	1	KN2	CAN-CAO-CAP-CAR
3	B	1	KN2	CAE-CAO-CAP-CAR

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1	KN2	2	0
3	B	1	KN2	1	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	243/258 (94%)	-0.44	2 (0%) 82 80	16, 40, 60, 72	8 (3%)
1	B	242/258 (93%)	-0.49	0 100 100	18, 40, 63, 72	10 (4%)
2	C	9/13 (69%)	-0.00	0 100 100	35, 37, 40, 41	0
2	D	11/13 (84%)	0.33	1 (9%) 15 11	32, 34, 40, 43	0
All	All	505/542 (93%)	-0.44	3 (0%) 85 83	16, 40, 61, 72	18 (3%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	363	ARG	2.4
1	A	432	SER	2.4
2	D	696	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	KN2	A	1	18/18	0.92	0.05	58,59,63,64	0
3	KN2	B	1	18/18	0.95	0.05	66,67,69,69	0

6.5 Other polymers [i](#)

There are no such residues in this entry.