



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 07:51 PM UTC

PDB ID : 2QAR / pdb_00002qar
Title : Structure of the 2TEL crystallization module fused to T4 lysozyme with a helical linker.
Authors : Nauli, S.; Bowie, J.U.
Deposited on : 2007-06-15
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	86	7% 72% 27% .
1	D	86	6% 69% 23% 6% .
2	B	93	2% 90% 10%
2	E	93	94% 6%
3	C	163	31% 82% 11% 5% .
3	F	163	48% 85% 12% .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5591 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E80-TELSAM domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	86	Total	C	N	O	S	0	1	0
			710	452	125	132	1			
1	D	84	Total	C	N	O	S	0	0	0
			684	437	122	124	1			


- Molecule 2 is a protein called TELSAM domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	93	Total	C	N	O	S	0	1	0
			750	480	128	141	1			
2	E	93	Total	C	N	O	S	0	1	0
			750	480	129	140	1			

- Molecule 3 is a protein called Lysozyme.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	163	Total	C	N	O	S	0	1	0
			1296	816	233	241	6			
3	F	163	Total	C	N	O	S	0	1	0
			1290	813	230	241	6			

- Molecule 4 is NITRATE ION (CCD ID: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	N	O	0	0
			4	1	3		
4	B	1	Total	N	O	0	0
			4	1	3		
4	D	1	Total	N	O	0	0
			4	1	3		
4	D	1	Total	N	O	0	0
			4	1	3		
4	D	1	Total	N	O	0	0
			4	1	3		
4	E	1	Total	N	O	0	0
			4	1	3		
4	E	1	Total	N	O	0	0
			4	1	3		

- Molecule 5 is AMMONIUM ION (CCD ID: NH4) (formula: H₄N).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total N 1 1	0	0
5	D	1	Total N 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	14	Total O 14 14	0	0
6	B	19	Total O 19 19	0	0
6	C	5	Total O 5 5	0	0
6	D	17	Total O 17 17	0	0
6	E	18	Total O 18 18	0	0
6	F	4	Total O 4 4	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: E80-TELSAM domain



- Molecule 1: E80-TELSAM domain





- Molecule 2: TELSAM domain




- Molecule 2: TELSAM domain



- Molecule 3: Lysozyme



● Molecule 3: Lysozyme



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	119.51Å 119.51Å 58.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	103.70 – 2.40 103.70 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.9 (103.70-2.40) 87.7 (103.70-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 2.39Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.220 , 0.255 (Not available) , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	48.3	Xtriage
Anisotropy	0.002	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 63.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.487 for -h,-k,l 0.041 for h,-h-k,-l 0.040 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5591	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 31.99 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0103e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NH4, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/729	0.88	2/987 (0.2%)
1	D	0.36	0/703	0.88	2/952 (0.2%)
2	B	0.33	0/771	0.78	0/1048
2	E	0.33	0/768	0.78	0/1044
3	C	0.43	0/1317	1.01	9/1776 (0.5%)
3	F	0.42	0/1311	0.97	3/1769 (0.2%)
All	All	0.38	0/5599	0.91	16/7576 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	6
3	F	0	6
All	All	0	12

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	16	LEU	CA-C-N	6.29	133.03	121.70
3	F	16	LEU	C-N-CA	6.29	133.03	121.70
3	C	34	LEU	CA-C-N	5.89	132.78	121.54
3	C	34	LEU	C-N-CA	5.89	132.78	121.54
3	C	56	ASN	CA-C-N	5.86	132.24	121.70
3	C	56	ASN	C-N-CA	5.86	132.24	121.70
3	C	57	GLY	CA-C-N	5.80	132.13	121.70
3	C	57	GLY	C-N-CA	5.80	132.13	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	15	SER	CA-C-N	5.24	131.13	121.70
1	D	15	SER	C-N-CA	5.24	131.13	121.70
3	C	35	THR	N-CA-CB	5.19	119.27	110.49
3	C	127	TRP	N-CA-C	5.12	116.67	111.14
1	A	15	SER	CA-C-N	5.11	130.90	121.70
1	A	15	SER	C-N-CA	5.11	130.90	121.70
3	F	127	TRP	N-CA-C	5.08	116.63	111.14
3	C	34	LEU	N-CA-CB	5.07	119.06	110.49

There are no chirality outliers.

All (12) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	15	ARG	Peptide
3	C	16	LEU	Peptide
3	C	33	LEU	Peptide
3	C	34	LEU	Peptide
3	C	55	THR	Peptide
3	C	57	GLY	Peptide
3	F	15	ARG	Peptide
3	F	16	LEU	Peptide
3	F	33	LEU	Peptide
3	F	34	LEU	Peptide
3	F	55	THR	Peptide
3	F	57	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	710	0	689	16	0
1	D	684	0	660	17	0
2	B	750	0	723	6	0
2	E	750	0	720	4	0
3	C	1296	0	1308	16	0
3	F	1290	0	1297	14	0
4	A	4	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	4	0	0	0	0
4	D	16	0	0	1	0
4	E	8	0	0	0	0
5	A	1	0	0	0	0
5	D	1	0	0	0	0
6	A	14	0	0	0	0
6	B	19	0	0	1	0
6	C	5	0	0	0	0
6	D	17	0	0	0	0
6	E	18	0	0	1	0
6	F	4	0	0	0	0
All	All	5591	0	5397	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:14:ALA:N	1:D:15:SER:HA	1.96	0.78
1:A:16:ILE:O	1:A:16:ILE:HG22	1.84	0.77
3:C:16:LEU:H	3:C:17:LYS:HE2	1.56	0.70
3:F:55:THR:HG23	3:F:56:ASN:H	1.60	0.67
3:F:16:LEU:HB2	3:F:17:LYS:HB3	1.76	0.66
1:A:61:ALA:HB2	2:B:45:PHE:HB3	1.79	0.65
1:D:61:ALA:HB2	2:E:45:PHE:HB3	1.79	0.64
3:F:55:THR:CG2	3:F:56:ASN:H	2.11	0.64
1:D:17:ARG:HB3	1:D:18:LEU:O	1.99	0.62
1:A:16:ILE:O	1:A:16:ILE:CG2	2.51	0.58
2:B:79:ASP:O	2:B:83:GLU:HG2	2.06	0.56
1:A:16:ILE:HA	1:A:18:LEU:N	2.20	0.56
2:E:79:ASP:O	2:E:83:GLU:HG2	2.06	0.56
1:D:16:ILE:H	1:D:18:LEU:HB2	1.72	0.55
3:F:55:THR:CG2	3:F:56:ASN:N	2.70	0.55
3:F:37:SER:HB3	3:F:42:ALA:HB3	1.93	0.51
3:C:37:SER:HB3	3:C:42:ALA:HB3	1.92	0.51
1:D:14:ALA:N	1:D:15:SER:CA	2.72	0.51
3:C:34:LEU:HB3	3:C:35:THR:HB	1.93	0.50
3:F:35:THR:O	3:F:35:THR:HG23	2.12	0.50
3:C:35:THR:HG23	3:C:35:THR:O	2.14	0.48
3:C:127:TRP:HB3	3:C:155:ARG:HA	1.94	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:68:GLU:HG3	4:D:108:NO3:N	2.28	0.48
1:A:99:THR:HA	2:B:15:SER:HA	1.96	0.48
1:D:15:SER:O	1:D:87:HIS:NE2	2.47	0.47
3:F:127:TRP:HB3	3:F:155:ARG:HA	1.95	0.47
1:A:25:GLN:HB3	4:A:102:NO3:O2	2.15	0.47
3:C:23:GLU:HB3	3:C:25:TYR:HE1	1.80	0.47
3:C:23:GLU:CB	3:C:25:TYR:HE1	2.28	0.47
3:F:23:GLU:CB	3:F:25:TYR:HE1	2.28	0.47
3:F:25:TYR:CD1	3:F:25:TYR:N	2.81	0.47
1:A:97:GLY:HA2	1:A:98:SER:HA	1.62	0.47
3:C:88:VAL:HG21	3:C:119:LEU:HB3	1.97	0.47
1:D:18:LEU:HA	1:D:19:PRO:HD2	1.81	0.47
3:F:23:GLU:HB3	3:F:25:TYR:HE1	1.80	0.46
3:F:88:VAL:HG21	3:F:119:LEU:HB3	1.97	0.46
1:D:95:GLY:HA3	1:D:96:GLY:HA2	1.64	0.46
1:A:14:ALA:HA	1:A:15:SER:HA	1.65	0.45
3:C:128[B]:ASP:OD2	3:C:128[B]:ASP:N	2.45	0.45
1:A:26:PRO:O	1:A:59:GLY:HA3	2.17	0.45
1:A:15:SER:N	1:A:16:ILE:HB	2.31	0.45
2:B:24:LEU:HB2	6:B:112:HOH:O	2.15	0.44
3:C:17:LYS:HD2	3:C:17:LYS:C	2.42	0.44
1:D:15:SER:H	1:D:16:ILE:CB	2.30	0.44
3:C:44:LYS:HG3	3:C:56:ASN:O	2.18	0.43
1:D:82:TYR:O	1:D:86:GLN:HG2	2.18	0.43
1:A:29:TRP:O	1:A:58:ASN:HB2	2.18	0.43
1:A:79:ASP:O	1:A:83:GLU:HG2	2.18	0.43
1:D:79:ASP:O	1:D:83:GLU:HG2	2.17	0.43
2:E:24:LEU:HB2	6:E:115:HOH:O	2.18	0.43
1:D:26:PRO:O	1:D:59:GLY:HA3	2.18	0.43
1:D:87:HIS:O	1:D:91:GLN:HG2	2.18	0.42
1:A:14:ALA:HA	1:A:17:ARG:HD2	2.00	0.42
1:A:82:TYR:O	1:A:86:GLN:HG2	2.20	0.42
1:D:29:TRP:O	1:D:58:ASN:HB2	2.18	0.42
3:C:28:ILE:HB	3:C:34:LEU:HD11	2.02	0.42
3:C:23:GLU:HB3	3:C:25:TYR:CE1	2.55	0.41
3:C:156:THR:HB	3:C:158:THR:HG22	2.03	0.41
3:F:23:GLU:HB3	3:F:25:TYR:CE1	2.55	0.41
3:F:156:THR:HB	3:F:158:THR:HG22	2.03	0.41
1:A:93:PRO:HA	1:A:94:GLY:HA2	1.77	0.41
1:D:42:GLU:CD	1:D:49:PRO:HA	2.45	0.41
2:E:20:ALA:HA	2:E:23:ARG:HG3	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:GLU:CD	1:A:49:PRO:HA	2.45	0.40
3:C:55:THR:HG23	3:C:56:ASN:OD1	2.21	0.40
2:B:51:ASP:HB3	2:B:54:THR:HG23	2.03	0.40
1:D:57:MET:HB2	1:D:61:ALA:HB3	2.03	0.40
3:F:86:LYS:HB3	3:F:87:PRO:HD3	2.03	0.40
2:B:47:LEU:HD13	2:B:77:SER:HB3	2.03	0.40
3:C:86:LYS:HB3	3:C:87:PRO:HD3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	85/86 (99%)	75 (88%)	10 (12%)	0	100	100
1	D	82/86 (95%)	76 (93%)	4 (5%)	2 (2%)	4	5
2	B	92/93 (99%)	90 (98%)	2 (2%)	0	100	100
2	E	92/93 (99%)	90 (98%)	2 (2%)	0	100	100
3	C	162/163 (99%)	140 (86%)	18 (11%)	4 (2%)	4	5
3	F	162/163 (99%)	139 (86%)	22 (14%)	1 (1%)	21	32
All	All	675/684 (99%)	610 (90%)	58 (9%)	7 (1%)	12	20

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	35	THR
3	C	58	VAL
3	F	17	LYS
3	C	17	LYS
3	C	34	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	18	LEU
1	D	95	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	76/75 (101%)	74 (97%)	2 (3%)	40	63
1	D	71/75 (95%)	67 (94%)	4 (6%)	19	33
2	B	76/78 (97%)	76 (100%)	0	100	100
2	E	75/78 (96%)	75 (100%)	0	100	100
3	C	136/137 (99%)	127 (93%)	9 (7%)	15	26
3	F	135/137 (98%)	129 (96%)	6 (4%)	25	43
All	All	569/580 (98%)	548 (96%)	21 (4%)	30	51

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	ARG
1	A	80	GLU
3	C	3	ASN
3	C	4	ILE
3	C	8	LEU
3	C	17	LYS
3	C	25	TYR
3	C	33	LEU
3	C	35	THR
3	C	58	VAL
3	C	67	LEU
1	D	17	ARG
1	D	31	ARG
1	D	68	GLU
1	D	80	GLU
3	F	3	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	F	4	ILE
3	F	8	LEU
3	F	25	TYR
3	F	33	LEU
3	F	67	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	GLN
1	A	36	GLN
1	A	43	ASN
3	C	41	ASN
3	C	70	GLN
3	C	124	GLN
3	C	141	ASN
3	C	142	GLN
1	D	25	GLN
1	D	36	GLN
1	D	43	ASN
3	F	41	ASN
3	F	70	GLN
3	F	124	GLN
3	F	141	ASN
3	F	142	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 10 ligands modelled in this entry, 2 are modelled with single atom - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NO3	D	106	-	1,3,3	3.25	1 (100%)	0,3,3	-	-
4	NO3	D	104	-	1,3,3	3.27	1 (100%)	0,3,3	-	-
4	NO3	B	109	-	1,3,3	3.34	1 (100%)	0,3,3	-	-
4	NO3	D	101	-	1,3,3	3.32	1 (100%)	0,3,3	-	-
4	NO3	A	102	-	1,3,3	3.26	1 (100%)	0,3,3	-	-
4	NO3	D	108	-	1,3,3	3.29	1 (100%)	0,3,3	-	-
4	NO3	E	108	-	1,3,3	3.22	1 (100%)	0,3,3	-	-
4	NO3	E	109	-	1,3,3	3.31	1 (100%)	0,3,3	-	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	109	NO3	O1-N	3.34	1.40	1.24
4	D	101	NO3	O1-N	3.32	1.40	1.24
4	E	109	NO3	O1-N	3.31	1.40	1.24
4	D	108	NO3	O1-N	3.29	1.40	1.24
4	D	104	NO3	O1-N	3.27	1.40	1.24
4	A	102	NO3	O1-N	3.26	1.40	1.24
4	D	106	NO3	O1-N	3.25	1.40	1.24
4	E	108	NO3	O1-N	3.22	1.40	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	102	NO3	1	0
4	D	108	NO3	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	86/86 (100%)	0.61	6 (6%)	22 19	25, 49, 113, 157	1 (1%)
1	D	84/86 (97%)	0.46	5 (5%)	27 24	37, 50, 97, 122	0
2	B	93/93 (100%)	0.47	2 (2%)	62 58	30, 49, 74, 98	1 (1%)
2	E	93/93 (100%)	0.52	0	100 100	25, 49, 74, 98	1 (1%)
3	C	163/163 (100%)	1.57	50 (30%)	1 1	48, 96, 127, 150	1 (0%)
3	F	163/163 (100%)	2.13	79 (48%)	0 0	48, 96, 127, 150	1 (0%)
All	All	682/684 (99%)	1.15	142 (20%)	2 2	25, 71, 123, 157	5 (0%)

All (142) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	16	ILE	5.3
1	A	95	GLY	5.3
1	A	14	ALA	5.1
3	F	98	CYS	5.0
3	F	87	PRO	4.8
3	F	79	ILE	4.6
3	F	89	TYR	4.4
1	D	95	GLY	4.3
3	F	95	VAL	4.1
3	F	11	ASP	4.0
3	F	69	CYS	4.0
3	C	31	GLY	4.0
3	F	107	MET	4.0
3	F	100	LEU	4.0
1	A	96	GLY	3.6
3	F	29	GLY	3.6
3	F	62	ASP	3.6
3	F	104	VAL	3.5
3	F	74	ALA	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	F	80	LEU	3.5
3	F	75	ALA	3.5
3	F	97	ARG	3.5
3	F	51	ILE	3.5
3	F	85	LEU	3.4
3	F	31	GLY	3.4
3	C	62	ASP	3.4
3	C	24	GLY	3.4
3	C	10	ILE	3.4
3	C	107	MET	3.4
3	F	78	GLY	3.3
3	F	4	ILE	3.3
3	F	94	CYS	3.3
3	F	8	LEU	3.3
3	F	14	LEU	3.2
3	F	122	LEU	3.2
3	C	128[A]	ASP	3.1
3	F	24	GLY	3.1
3	F	139	TRP	3.1
3	F	72	VAL	3.1
3	C	98	CYS	3.0
3	F	10	ILE	3.0
3	F	52	GLY	3.0
3	F	113	ALA	3.0
3	C	87	PRO	3.0
3	F	134	LEU	3.0
3	C	156	THR	3.0
3	C	52	GLY	3.0
3	F	77	ARG	2.9
3	F	18	ILE	2.9
3	F	59	ILE	2.9
3	C	95	VAL	2.9
3	F	156	THR	2.9
3	C	14	LEU	2.9
3	F	32	HIS	2.9
3	F	47	LEU	2.9
3	F	140	TYR	2.9
3	F	27	THR	2.9
3	F	43	ALA	2.9
3	F	76	VAL	2.8
3	C	8	LEU	2.8
3	C	139	TRP	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	89	TYR	2.8
3	F	67	LEU	2.8
3	F	86	LYS	2.8
3	F	119	LEU	2.7
3	C	11	ASP	2.7
3	F	102	ASN	2.7
3	F	19	TYR	2.7
3	F	162	TYR	2.7
3	C	43	ALA	2.7
3	F	39	SER	2.7
3	C	162	TYR	2.7
3	F	50	ALA	2.7
1	A	94	GLY	2.6
3	F	154	PHE	2.6
3	C	157	GLY	2.6
3	C	151	ILE	2.6
3	F	117	ASN	2.6
3	F	137	SER	2.6
3	F	157	GLY	2.5
3	C	94	CYS	2.5
1	D	96	GLY	2.5
3	C	51	ILE	2.5
3	F	92	LEU	2.5
3	C	140	TYR	2.5
3	C	29	GLY	2.5
3	C	18	ILE	2.5
3	C	50	ALA	2.5
3	C	143	THR	2.5
3	F	65	GLU	2.5
3	F	71	ASP	2.5
3	F	151	ILE	2.5
3	F	149	ARG	2.5
3	C	78	GLY	2.4
3	F	124	GLN	2.4
3	C	74	ALA	2.4
3	F	128[A]	ASP	2.4
3	C	80	LEU	2.4
2	B	35	ALA	2.4
3	F	42	ALA	2.4
3	C	27	THR	2.4
1	A	36	GLN	2.4
3	F	159	TRP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	14	ALA	2.3
3	F	109	GLU	2.3
3	C	79	ILE	2.3
3	F	73	ASP	2.3
3	F	150	VAL	2.3
3	C	113	ALA	2.3
3	F	28	ILE	2.3
3	C	121	MET	2.3
3	F	90	ASP	2.3
3	C	39	SER	2.2
3	F	82	ASN	2.2
3	F	125	LYS	2.2
3	C	4	ILE	2.2
3	F	115	PHE	2.2
1	D	15	SER	2.2
3	C	37	SER	2.2
3	F	103	MET	2.2
3	C	19	TYR	2.2
3	F	38	PRO	2.2
3	C	42	ALA	2.1
3	C	134	LEU	2.1
1	A	20	ALA	2.1
3	C	122	LEU	2.1
3	F	118	SER	2.1
3	C	124	GLN	2.1
3	C	75	ALA	2.1
2	B	59	GLY	2.1
3	F	34	LEU	2.1
3	C	65	GLU	2.1
3	C	158	THR	2.1
3	F	7	MET	2.1
3	C	76	VAL	2.0
3	C	135	ALA	2.0
3	C	47	LEU	2.0
3	C	25	TYR	2.0
3	F	112	VAL	2.0
3	C	85	LEU	2.0
3	F	108	GLY	2.0
3	F	101	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NO3	D	108	4/4	0.37	0.14	114,117,117,118	0
4	NO3	B	109	4/4	0.73	0.15	87,88,89,90	0
4	NO3	E	109	4/4	0.73	0.16	87,93,95,95	0
4	NO3	D	104	4/4	0.78	0.21	94,96,96,97	0
4	NO3	D	101	4/4	0.79	0.22	66,69,72,76	0
4	NO3	E	108	4/4	0.84	0.23	71,73,74,77	0
4	NO3	A	102	4/4	0.86	0.28	68,70,71,76	0
4	NO3	D	106	4/4	0.94	0.21	79,80,81,82	0
5	NH4	D	202	1/1	0.96	0.07	41,41,41,41	0
5	NH4	A	201	1/1	0.97	0.06	37,37,37,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.