



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:15 AM UTC

PDB ID : 9QA3 / pdb_00009qa3
Title : Drosophila melanogaster angiotensin converting enzyme homologue, AnCE in complex with YW dipeptide
Authors : Zukowska, J.; Gregory, K.S.; Acharya, K.R.
Deposited on : 2025-02-27
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	597	86% 12% .
2	B	2	50% 100%
2	D	2	50% 50%
3	C	6	100%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5335 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Angiotensin-converting enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	597	Total	C	N	O	S	0	0	0
			4874	3120	804	930	20			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	346	ILE	THR	conflict	UNP Q10714

- Molecule 2 is a protein called TYR-TRP.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			27	20	3	4			
2	D	2	Total	C	N	O	0	0	0
			27	20	3	4			

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	301	Total	O	0	0
			301	301		
6	B	3	Total	O	0	0
			3	3		
6	D	2	Total	O	0	0
			2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Angiotensin-converting enzyme




- Molecule 2: TYR-TRP



- Molecule 2: TYR-TRP



- Molecule 3: beta-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	172.43Å 172.43Å 103.50Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	86.22 – 2.20 86.22 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (86.22-2.20) 100.0 (86.22-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0430 (refmacat 0.4.100)	Depositor
R, R_{free}	0.163 , 0.206 0.172 , 0.212	Depositor DCC
R_{free} test set	2789 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	31.2	Xtriage
Anisotropy	0.494	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 36.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.003 for -2/3*h-1/3*k-4/3*l,-1/3*h-2/3*k+4/3*l,-1/3*h+1/3*k+1/3*l 0.000 for -h,1/3*h-1/3*k-4/3*l,-1/3*h-2/3*k+1/3*l 0.000 for -1/3*h+1/3*k+4/3*l,-k,2/3*h+1/3*k+1/3*l 0.000 for -h,2/3*h+1/3*k+4/3*l,1/3*h+2/3*k-1/3*l 0.009 for -1/3*h-2/3*k+4/3*l,-2/3*h-1/3*k-4/3*l,1/3*h-1/3*k-1/3*l 0.000 for 1/3*h+2/3*k-4/3*l,-k,-2/3*h-1/3*k-1/3*l 0.015 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5335	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, ZN, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.94	2/5000 (0.0%)	1.52	42/6772 (0.6%)
2	B	1.04	0/29	1.67	0/38
2	D	0.85	0/29	1.84	1/38 (2.6%)
All	All	0.94	2/5058 (0.0%)	1.52	43/6848 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	157	ARG	CZ-NH1	5.48	1.40	1.32
1	A	481	PRO	CA-CB	-5.43	1.48	1.53

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	437	ASP	CA-CB-CG	9.39	121.99	112.60
1	A	157	ARG	NE-CZ-NH2	-9.16	110.95	119.20
1	A	157	ARG	NE-CZ-NH1	8.32	129.82	121.50
1	A	211	ASP	CB-CA-C	-8.29	95.29	110.63
1	A	198	THR	CA-CB-OG1	-8.11	97.44	109.60
1	A	343	PHE	CA-CB-CG	-7.99	105.81	113.80
1	A	401	LEU	N-CA-CB	7.90	121.53	110.07
1	A	211	ASP	CA-CB-CG	7.00	119.60	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	383	PHE	CA-CB-CG	-6.83	106.97	113.80
1	A	332	ARG	CB-CG-CD	-6.53	96.29	111.30
1	A	27	GLU	CB-CG-CD	6.50	123.65	112.60
1	A	515	PHE	CA-CB-CG	-6.35	107.45	113.80
1	A	566	ASP	CA-CB-CG	6.26	118.86	112.60
2	D	1629	TYR	CE1-CZ-OH	-6.13	101.51	119.90
1	A	481	PRO	N-CA-C	6.12	116.35	110.47
1	A	474	ASP	CA-CB-CG	5.99	118.59	112.60
1	A	401	LEU	CA-C-O	-5.87	114.66	120.70
1	A	529	ASP	CA-CB-CG	5.86	118.46	112.60
1	A	38	LYS	N-CA-CB	-5.83	101.62	110.07
1	A	38	LYS	CB-CA-C	5.79	120.06	110.90
1	A	369	LEU	N-CA-C	-5.79	105.71	112.89
1	A	375	PHE	CB-CA-C	-5.78	101.56	110.81
1	A	574	GLU	CG-CD-OE2	-5.76	105.15	118.40
1	A	81	THR	CA-CB-OG1	-5.67	101.09	109.60
1	A	53	ASN	CA-CB-CG	5.62	118.22	112.60
1	A	316	PRO	CB-CA-C	-5.55	102.83	111.22
1	A	24	GLN	N-CA-CB	5.46	118.75	110.28
1	A	280	LYS	CB-CA-C	5.46	117.33	109.09
1	A	182	GLU	CB-CA-C	-5.45	101.74	110.79
1	A	169	PHE	CA-CB-CG	-5.34	108.46	113.80
1	A	599	LYS	CB-CA-C	5.31	119.71	110.68
1	A	186	GLU	CB-CA-C	-5.30	101.83	110.85
1	A	481	PRO	CA-C-O	5.19	124.64	120.90
1	A	42	VAL	N-CA-CB	-5.17	102.83	110.58
1	A	315	LEU	CA-C-O	5.14	124.58	119.75
1	A	140	LYS	CB-CA-C	5.13	119.34	110.77
1	A	403	LEU	CA-C-N	5.12	127.09	120.44
1	A	403	LEU	C-N-CA	5.12	127.09	120.44
1	A	595	ALA	N-CA-C	-5.07	105.46	111.69
1	A	574	GLU	CG-CD-OE1	5.07	130.05	118.40
1	A	115	GLU	CB-CG-CD	5.04	121.16	112.60
1	A	305	ASP	N-CA-CB	5.03	117.30	110.01
1	A	604	ILE	CA-C-O	-5.02	115.00	120.72

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	183	ARG	Sidechain
1	A	332	ARG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	356	ARG	Sidechain
1	A	473	ARG	Sidechain
1	A	590	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4874	0	4675	25	0
2	B	27	0	21	0	0
2	D	27	0	18	2	0
3	C	72	0	61	0	0
4	A	1	0	0	0	0
5	A	28	0	26	0	0
6	A	301	0	0	4	0
6	B	3	0	0	0	0
6	D	2	0	0	1	0
All	All	5335	0	4801	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:380:HIS:HE1	6:A:1064:HOH:O	1.71	0.73
1:A:121:SER:O	1:A:125:SER:HB3	2.03	0.59
1:A:306:PHE:CD1	1:A:401:LEU:HD13	2.40	0.57
1:A:380:HIS:HD2	6:A:988:HOH:O	1.90	0.55
1:A:132:VAL:HG22	1:A:149:ILE:HD11	1.90	0.53
1:A:276:PRO:HB3	1:A:592:TRP:CH2	2.44	0.53
1:A:380:HIS:H	1:A:380:HIS:CD2	2.29	0.51
2:D:1629:TYR:N	6:D:1701:HOH:O	2.44	0.50
1:A:113:TYR:CE2	1:A:117:LEU:HD11	2.47	0.49
1:A:564:TRP:N	1:A:565:PRO:CD	2.76	0.49
1:A:442:LEU:N	1:A:443:PRO:HD2	2.29	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:LEU:HA	1:A:149:ILE:HB	1.97	0.45
1:A:442:LEU:HB2	1:A:443:PRO:HD3	1.99	0.44
1:A:587:GLU:N	1:A:588:PRO:HD2	2.32	0.44
1:A:146:ASP:HA	1:A:147:PRO:HA	1.71	0.44
1:A:20:LYS:HE3	1:A:24:GLN:NE2	2.33	0.43
1:A:86:ARG:HG3	6:A:1029:HOH:O	2.17	0.43
1:A:157:ARG:NH2	1:A:272:ASP:OD1	2.50	0.43
1:A:266:GLN:NE2	2:D:1630:TRP:HE1	2.16	0.43
1:A:479:ILE:C	1:A:479:ILE:HD12	2.44	0.42
1:A:96:GLN:O	1:A:100:LEU:HG	2.19	0.42
1:A:285:VAL:O	1:A:286:SER:C	2.62	0.41
1:A:285:VAL:HA	6:A:865:HOH:O	2.19	0.41
1:A:315:LEU:HD13	1:A:319:PHE:CD1	2.56	0.41
1:A:410:HIS:ND1	1:A:541:ASP:OD2	2.45	0.41
1:A:532:ASN:OD1	1:A:534:GLU:HG2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	595/597 (100%)	585 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	520/520 (100%)	505 (97%)	15 (3%)	37	51
2	B	2/2 (100%)	2 (100%)	0	100	100
2	D	2/2 (100%)	2 (100%)	0	100	100
All	All	524/524 (100%)	509 (97%)	15 (3%)	37	51

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	GLU
1	A	34	LYS
1	A	60	LYS
1	A	98	LYS
1	A	125	SER
1	A	211	ASP
1	A	273	ILE
1	A	276	PRO
1	A	326	GLU
1	A	360	ASP
1	A	378	TYR
1	A	394	HIS
1	A	462	LYS
1	A	488	LYS
1	A	540	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	266	GLN
1	A	380	HIS
1	A	464	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	3,1	14,14,15	0.56	0	17,19,21	1.35	2 (11%)
3	NAG	C	2	3	14,14,15	0.47	0	17,19,21	1.93	4 (23%)
3	BMA	C	3	3	11,11,12	1.28	1 (9%)	15,15,17	1.63	3 (20%)
3	MAN	C	4	3	11,11,12	1.21	1 (9%)	15,15,17	1.82	5 (33%)
3	BMA	C	5	3	11,11,12	1.13	1 (9%)	15,15,17	2.47	6 (40%)
3	MAN	C	6	3	11,11,12	1.29	1 (9%)	15,15,17	3.00	9 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1
3	BMA	C	3	3	-	2/2/19/22	0/1/1/1
3	MAN	C	4	3	-	2/2/19/22	0/1/1/1
3	BMA	C	5	3	-	1/2/19/22	0/1/1/1
3	MAN	C	6	3	-	0/2/19/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	5	BMA	C2-C3	-3.16	1.47	1.52
3	C	6	MAN	C4-C5	2.48	1.58	1.53
3	C	4	MAN	C4-C5	2.44	1.58	1.53
3	C	3	BMA	C4-C5	2.33	1.58	1.53

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	6	MAN	C1-C2-C3	7.54	120.62	109.64
3	C	5	BMA	C2-C3-C4	-5.14	101.82	110.86
3	C	2	NAG	C1-O5-C5	4.57	118.31	112.19
3	C	6	MAN	O3-C3-C4	4.34	120.60	110.38
3	C	2	NAG	C2-N2-C7	-4.30	117.14	122.90
3	C	5	BMA	O5-C1-C2	4.20	120.82	110.79
3	C	5	BMA	C1-O5-C5	4.08	117.65	112.19
3	C	1	NAG	C2-N2-C7	3.84	128.04	122.90
3	C	3	BMA	C3-C4-C5	3.63	116.81	110.23
3	C	3	BMA	O4-C4-C3	-3.60	101.88	110.38
3	C	4	MAN	C3-C4-C5	3.58	116.73	110.23
3	C	6	MAN	O6-C6-C5	3.40	122.92	111.33
3	C	5	BMA	C3-C4-C5	-3.15	104.52	110.23
3	C	6	MAN	C6-C5-C4	3.12	120.68	113.02
3	C	5	BMA	O3-C3-C4	3.04	117.53	110.38
3	C	6	MAN	O5-C1-C2	2.85	117.58	110.79
3	C	6	MAN	C1-O5-C5	2.74	115.86	112.19
3	C	4	MAN	C1-C2-C3	-2.73	105.66	109.64
3	C	4	MAN	C1-O5-C5	2.70	115.80	112.19
3	C	2	NAG	O5-C1-C2	2.62	115.35	111.29
3	C	6	MAN	O3-C3-C2	-2.46	105.03	110.05
3	C	5	BMA	C1-C2-C3	2.40	113.14	109.64
3	C	4	MAN	O2-C2-C3	2.27	114.86	110.15
3	C	1	NAG	C4-C3-C2	2.20	114.25	111.02
3	C	6	MAN	O5-C5-C6	2.18	111.91	107.66
3	C	3	BMA	C1-O5-C5	2.15	115.07	112.19
3	C	2	NAG	O6-C6-C5	-2.12	104.10	111.33
3	C	4	MAN	O5-C5-C6	-2.07	103.64	107.66
3	C	6	MAN	O5-C5-C4	-2.06	105.81	110.83

There are no chirality outliers.


All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	2	NAG	O5-C5-C6-O6
3	C	4	MAN	O5-C5-C6-O6
3	C	2	NAG	C4-C5-C6-O6
3	C	4	MAN	C4-C5-C6-O6
3	C	3	BMA	C4-C5-C6-O6
3	C	3	BMA	O5-C5-C6-O6
3	C	5	BMA	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	A	702	1	14,14,15	0.93	0	17,19,21	2.69	6 (35%)
5	NAG	A	703	1	14,14,15	0.50	0	17,19,21	1.89	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	702	1	-	2/6/23/26	0/1/1/1
5	NAG	A	703	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	702	NAG	C1-C2-N2	-8.03	97.77	110.43
5	A	703	NAG	C3-C4-C5	-4.13	102.75	110.23
5	A	702	NAG	O4-C4-C5	3.66	118.35	109.32
5	A	703	NAG	C2-N2-C7	3.51	127.60	122.90
5	A	702	NAG	C4-C3-C2	3.29	115.84	111.02
5	A	702	NAG	O7-C7-N2	-3.11	116.48	121.98
5	A	703	NAG	C4-C3-C2	-2.74	107.00	111.02
5	A	702	NAG	O7-C7-C8	2.61	126.70	122.05
5	A	703	NAG	O3-C3-C2	2.44	114.46	109.40
5	A	702	NAG	C3-C4-C5	-2.42	105.84	110.23
5	A	703	NAG	O4-C4-C5	2.18	114.68	109.32

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	702	NAG	C8-C7-N2-C2
5	A	703	NAG	C4-C5-C6-O6
5	A	702	NAG	O7-C7-N2-C2
5	A	703	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	597/597 (100%)	-0.31	5 (0%) 82 80	23, 33, 57, 88	0
2	B	2/2 (100%)	3.28	1 (50%) 0 0	55, 55, 55, 87	0
2	D	2/2 (100%)	1.69	1 (50%) 0 0	49, 49, 49, 49	0
All	All	601/601 (100%)	-0.29	7 (1%) 76 74	23, 33, 57, 88	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1	TYR	4.7
1	A	20	LYS	2.5
1	A	27	GLU	2.3
1	A	102	LYS	2.1
1	A	18	LEU	2.1
2	D	1629	TYR	2.1
1	A	279	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	C	1	14/15	-	-	28,35,38,38	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	C	2	14/15	-	-	37,46,62,63	0
3	BMA	C	3	11/12	-	-	51,67,73,73	0
3	MAN	C	4	11/12	-	-	56,67,76,79	0
3	BMA	C	5	11/12	-	-	55,85,90,134	0
3	MAN	C	6	11/12	-	-	63,81,93,101	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	702	14/15	0.85	0.15	55,70,89,93	0
5	NAG	A	703	14/15	0.85	0.16	43,79,92,96	0
4	ZN	A	701	1/1	1.00	0.02	29,29,29,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.