



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 10:05 AM UTC

PDB ID : 4QBY / pdb_00004qby
Title : yCP in complex with BOC-ALA-ALA-ALA-CHO
Authors : Arciniega, M.; Beck, P.; Lange, O.; Groll, M.; Huber, R.
Deposited on : 2014-05-09
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	
1	O	250	
2	B	258	
2	P	258	
3	C	254	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Q	254	 78%12% • 6%
4	D	260	 71%17% • 10%
4	R	260	 71%17% • 10%
5	E	234	 73%21% • •
5	S	234	 73%23% • •
6	F	288	 72%11% • 16%
6	T	288	 71%13% • 16%
7	G	252	 79%14% • •
7	U	252	 77%16% • •
8	H	232	 77%18% • •
8	V	232	 78%17% • •
9	I	205	 78%20% •
9	W	205	 78%20% •
10	J	198	 %81%15% • •
10	X	198	 %81%15% • •
11	K	212	 83%17%
11	Y	212	 86%13% •
12	L	222	 82%17% •
12	Z	222	 82%16% •
13	M	246	 84%9% • 5%
13	a	246	 83%10% • 5%
14	N	196	 84%15% •
14	b	196	 83%15% •
15	1	4	 75%25%
15	2	4	 75%25%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain	
15	3	4		
15	4	4		

2 Entry composition

There are 17 unique types of molecules in this entry. The entry contains 49645 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit alpha type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			
1	O	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			

- Molecule 2 is a protein called Proteasome subunit alpha type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	244	Total	C	N	O	S	0	0	0
			1904	1201	321	379	3			
2	P	244	Total	C	N	O	S	0	0	0
			1904	1201	321	379	3			

- Molecule 3 is a protein called Proteasome subunit alpha type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	240	Total	C	N	O	S	0	0	0
			1881	1176	329	372	4			
3	Q	240	Total	C	N	O	S	0	0	0
			1881	1176	329	372	4			

- Molecule 4 is a protein called Proteasome subunit alpha type-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	235	Total	C	N	O	S	0	0	0
			1813	1136	304	366	7			
4	R	235	Total	C	N	O	S	0	0	0
			1813	1136	304	366	7			

- Molecule 5 is a protein called Proteasome subunit alpha type-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	231	Total	C	N	O	S	0	0	0
			1773	1114	307	348	4			
5	S	231	Total	C	N	O	S	0	0	0
			1773	1114	307	348	4			

- Molecule 6 is a protein called Probable proteasome subunit alpha type-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	243	Total	C	N	O	S	0	0	0
			1892	1203	329	356	4			
6	T	243	Total	C	N	O	S	0	0	0
			1892	1203	329	356	4			

- Molecule 7 is a protein called Proteasome subunit alpha type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	241	Total	C	N	O	S	0	0	0
			1907	1214	320	365	8			
7	U	241	Total	C	N	O	S	0	0	0
			1907	1214	320	365	8			

- Molecule 8 is a protein called Proteasome subunit beta type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	226	Total	C	N	O	S	0	0	0
			1719	1082	298	332	7			
8	V	226	Total	C	N	O	S	0	0	0
			1719	1082	298	332	7			

- Molecule 9 is a protein called Proteasome subunit beta type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			
9	W	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			

- Molecule 10 is a protein called Proteasome subunit beta type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	195	Total	C	N	O	S	0	0	0
			1561	992	264	299	6			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	X	195	Total	C	N	O	S	0	0	0
			1561	992	264	299	6			

- Molecule 11 is a protein called Proteasome subunit beta type-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			
11	Y	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			

- Molecule 12 is a protein called Proteasome subunit beta type-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			
12	Z	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			

- Molecule 13 is a protein called Proteasome subunit beta type-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			
13	a	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			

- Molecule 14 is a protein called Proteasome subunit beta type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			
14	b	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			

- Molecule 15 is a protein called BOC-ALA-ALA-ALA-CHO.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
15	1	4	Total	C	N	O	0	0	0
			22	14	3	5			
15	2	4	Total	C	N	O	0	0	0
			22	14	3	5			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
15	3	4	Total	C	N	O	0	0	0
			22	14	3	5			
15	4	4	Total	C	N	O	0	0	0
			22	14	3	5			

- Molecule 16 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	G	1	Total	Mg	0	0
			1	1		
16	H	1	Total	Mg	0	0
			1	1		
16	I	2	Total	Mg	0	0
			2	2		
16	J	1	Total	Mg	0	0
			1	1		
16	K	2	Total	Mg	0	0
			2	2		
16	N	1	Total	Mg	0	0
			1	1		
16	V	1	Total	Mg	0	0
			1	1		
16	Y	1	Total	Mg	0	0
			1	1		
16	Z	1	Total	Mg	0	0
			1	1		

- Molecule 17 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	A	7	Total	O	0	0
			7	7		
17	B	6	Total	O	0	0
			6	6		
17	C	5	Total	O	0	0
			5	5		
17	D	5	Total	O	0	0
			5	5		
17	E	5	Total	O	0	0
			5	5		
17	F	6	Total	O	0	0
			6	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	G	10	Total 10	O 10	0	0
17	H	8	Total 8	O 8	0	0
17	I	4	Total 4	O 4	0	0
17	J	8	Total 8	O 8	0	0
17	K	3	Total 3	O 3	0	0
17	L	7	Total 7	O 7	0	0
17	M	10	Total 10	O 10	0	0
17	N	6	Total 6	O 6	0	0
17	O	5	Total 5	O 5	0	0
17	P	8	Total 8	O 8	0	0
17	Q	6	Total 6	O 6	0	0
17	R	3	Total 3	O 3	0	0
17	S	1	Total 1	O 1	0	0
17	T	7	Total 7	O 7	0	0
17	U	5	Total 5	O 5	0	0
17	V	11	Total 11	O 11	0	0
17	W	3	Total 3	O 3	0	0
17	X	7	Total 7	O 7	0	0
17	Y	5	Total 5	O 5	0	0
17	Z	4	Total 4	O 4	0	0
17	2	1	Total 1	O 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	a	16	Total 16	O 16	0	0
17	3	2	Total 2	O 2	0	0
17	b	5	Total 5	O 5	0	0
17	4	1	Total 1	O 1	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome subunit alpha type-2

Chain A: 




- Molecule 1: Proteasome subunit alpha type-2

Chain O: 




- Molecule 2: Proteasome subunit alpha type-3

Chain B: 




- Molecule 2: Proteasome subunit alpha type-3

Chain P: 




- Molecule 3: Proteasome subunit alpha type-4

Chain C:  77% 13% 6%




- Molecule 3: Proteasome subunit alpha type-4

Chain Q:  78% 12% 6%




- Molecule 4: Proteasome subunit alpha type-5

Chain D:  71% 17% 10%




- Molecule 4: Proteasome subunit alpha type-5

Chain R:  71% 17% 10%




- Molecule 5: Proteasome subunit alpha type-6

Chain E:  73% 21% 6%




- Molecule 5: Proteasome subunit alpha type-6

Chain S:  73% 23% . .




- Molecule 6: Probable proteasome subunit alpha type-7

Chain F:  72% 11% . 16%




- Molecule 6: Probable proteasome subunit alpha type-7

Chain T:  71% 13% . 16%




- Molecule 7: Proteasome subunit alpha type-1

Chain G:  79% 14% . . .




- Molecule 7: Proteasome subunit alpha type-1


Chain U:  77% 16% . . .




- Molecule 8: Proteasome subunit beta type-2




- Molecule 8: Proteasome subunit beta type-2




- Molecule 9: Proteasome subunit beta type-3




- Molecule 9: Proteasome subunit beta type-3




- Molecule 10: Proteasome subunit beta type-4




- Molecule 10: Proteasome subunit beta type-4




- Molecule 11: Proteasome subunit beta type-5




- Molecule 11: Proteasome subunit beta type-5



- Molecule 12: Proteasome subunit beta type-6



- Molecule 12: Proteasome subunit beta type-6



- Molecule 13: Proteasome subunit beta type-7

Chain M:  84% 9% • 5%




- Molecule 13: Proteasome subunit beta type-7

Chain a:  83% 10% • 5%



- Molecule 14: Proteasome subunit beta type-1

Chain N:  84% 15% •



- Molecule 15: BOC-ALA-ALA-ALA-CHO

Chain 1:  75% 25%



- Molecule 15: BOC-ALA-ALA-ALA-CHO

Chain 2:  75% 25%



- Molecule 15: BOC-ALA-ALA-ALA-CHO

Chain 3:  75% 25%



• Molecule 15: BOC-ALA-ALA-ALA-CHO

Chain 4:  75% 25%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	134.99Å 300.13Å 144.80Å 90.00° 112.51° 90.00°	Depositor
Resolution (Å)	15.00 – 3.00 15.00 – 3.00	Depositor EDS
% Data completeness (in resolution range)	92.1 (15.00-3.00) 91.4 (15.00-3.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 3.01Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.178 , 0.207 0.179 , 0.204	Depositor DCC
R_{free} test set	9676 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	65.1	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 60.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	49645	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BOC, MG, 2A1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/1952	0.79	0/2642
1	O	0.43	0/1952	0.79	0/2642
2	B	0.42	0/1934	0.77	0/2618
2	P	0.42	0/1934	0.77	0/2618
3	C	0.45	0/1910	0.83	0/2586
3	Q	0.45	0/1910	0.83	1/2586 (0.0%)
4	D	0.43	0/1837	0.75	0/2475
4	R	0.43	0/1837	0.75	0/2475
5	E	0.42	0/1800	0.76	2/2433 (0.1%)
5	S	0.42	0/1800	0.76	0/2433
6	F	0.42	0/1932	0.79	0/2609
6	T	0.43	0/1932	0.79	0/2609
7	G	0.41	0/1945	0.78	0/2634
7	U	0.41	0/1945	0.78	0/2634
8	H	0.41	0/1750	0.76	0/2373
8	V	0.41	0/1750	0.76	0/2373
9	I	0.41	0/1611	0.75	0/2174
9	W	0.41	0/1611	0.75	0/2174
10	J	0.43	0/1589	0.76	0/2142
10	X	0.43	0/1589	0.75	0/2142
11	K	0.42	0/1681	0.78	0/2274
11	Y	0.42	0/1681	0.78	0/2274
12	L	0.41	0/1795	0.76	0/2420
12	Z	0.40	0/1795	0.75	0/2420
13	M	0.42	0/1855	0.77	0/2514
13	a	0.42	0/1855	0.77	0/2514
14	N	0.40	0/1541	0.75	0/2087
14	b	0.41	0/1541	0.75	0/2087
15	1	0.87	0/9	0.40	0/11
15	2	0.49	0/9	0.63	0/11
15	3	0.86	0/9	0.38	0/11
15	4	0.51	0/9	0.68	0/11

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
All	All	0.42	0/50300	0.77	3/68006 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
5	E	147	GLN	CA-C-N	5.13	124.85	119.82
5	E	147	GLN	C-N-CA	5.13	124.85	119.82
3	Q	4	ARG	N-CA-C	5.06	117.81	110.42

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1915	0	1929	11	0
1	O	1915	0	1929	14	0
2	B	1904	0	1904	24	0
2	P	1904	0	1904	24	0
3	C	1881	0	1895	21	0
3	Q	1881	0	1895	17	0
4	D	1813	0	1797	21	0
4	R	1813	0	1797	21	0
5	E	1773	0	1775	27	0
5	S	1773	0	1775	27	0
6	F	1892	0	1883	17	0
6	T	1892	0	1883	20	0
7	G	1907	0	1901	15	0
7	U	1907	0	1901	20	0
8	H	1719	0	1718	26	0
8	V	1719	0	1718	25	0
9	I	1581	0	1574	25	0
9	W	1581	0	1574	22	0
10	J	1561	0	1569	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	X	1561	0	1569	14	0
11	K	1644	0	1594	18	0
11	Y	1644	0	1594	16	0
12	L	1757	0	1711	23	0
12	Z	1757	0	1711	22	0
13	M	1824	0	1832	13	0
13	a	1824	0	1832	13	0
14	N	1512	0	1481	19	0
14	b	1512	0	1481	19	0
15	1	22	0	19	3	0
15	2	22	0	19	1	0
15	3	22	0	19	2	0
15	4	22	0	19	1	0
16	G	1	0	0	0	0
16	H	1	0	0	0	0
16	I	2	0	0	0	0
16	J	1	0	0	0	0
16	K	2	0	0	0	0
16	N	1	0	0	0	0
16	V	1	0	0	0	0
16	Y	1	0	0	0	0
16	Z	1	0	0	0	0
17	2	1	0	0	0	0
17	3	2	0	0	0	0
17	4	1	0	0	0	0
17	A	7	0	0	0	0
17	B	6	0	0	0	0
17	C	5	0	0	0	0
17	D	5	0	0	0	0
17	E	5	0	0	0	0
17	F	6	0	0	0	0
17	G	10	0	0	0	0
17	H	8	0	0	1	0
17	I	4	0	0	0	0
17	J	8	0	0	1	0
17	K	3	0	0	0	0
17	L	7	0	0	0	0
17	M	10	0	0	0	0
17	N	6	0	0	0	0
17	O	5	0	0	0	0
17	P	8	0	0	0	0
17	Q	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
17	R	3	0	0	0	0
17	S	1	0	0	0	0
17	T	7	0	0	0	0
17	U	5	0	0	0	0
17	V	11	0	0	1	0
17	W	3	0	0	0	0
17	X	7	0	0	0	0
17	Y	5	0	0	0	0
17	Z	4	0	0	0	0
17	a	16	0	0	1	0
17	b	5	0	0	0	0
All	All	49645	0	49202	495	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 495 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:K:209:ASN:ND2	10:X:130:TYR:OH	1.93	0.99
3:C:201:VAL:O	3:C:202:GLN:HB3	1.79	0.82
3:Q:201:VAL:O	3:Q:202:GLN:HB3	1.80	0.82
12:L:172:LEU:H	12:L:172:LEU:HD23	1.46	0.81
12:Z:172:LEU:HD23	12:Z:172:LEU:H	1.46	0.79

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	248/250 (99%)	240 (97%)	6 (2%)	2 (1%)	16 50

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	O	248/250 (99%)	240 (97%)	6 (2%)	2 (1%)	16	50
2	B	242/258 (94%)	226 (93%)	12 (5%)	4 (2%)	7	32
2	P	242/258 (94%)	226 (93%)	12 (5%)	4 (2%)	7	32
3	C	238/254 (94%)	226 (95%)	9 (4%)	3 (1%)	9	38
3	Q	238/254 (94%)	226 (95%)	9 (4%)	3 (1%)	9	38
4	D	231/260 (89%)	222 (96%)	8 (4%)	1 (0%)	30	65
4	R	231/260 (89%)	223 (96%)	7 (3%)	1 (0%)	30	65
5	E	229/234 (98%)	217 (95%)	10 (4%)	2 (1%)	14	48
5	S	229/234 (98%)	217 (95%)	10 (4%)	2 (1%)	14	48
6	F	241/288 (84%)	229 (95%)	11 (5%)	1 (0%)	30	65
6	T	241/288 (84%)	229 (95%)	11 (5%)	1 (0%)	30	65
7	G	239/252 (95%)	228 (95%)	10 (4%)	1 (0%)	30	65
7	U	239/252 (95%)	228 (95%)	10 (4%)	1 (0%)	30	65
8	H	224/232 (97%)	216 (96%)	8 (4%)	0	100	100
8	V	224/232 (97%)	216 (96%)	8 (4%)	0	100	100
9	I	202/205 (98%)	192 (95%)	10 (5%)	0	100	100
9	W	202/205 (98%)	192 (95%)	10 (5%)	0	100	100
10	J	193/198 (98%)	180 (93%)	12 (6%)	1 (0%)	24	60
10	X	193/198 (98%)	181 (94%)	11 (6%)	1 (0%)	24	60
11	K	210/212 (99%)	199 (95%)	9 (4%)	2 (1%)	12	45
11	Y	210/212 (99%)	199 (95%)	8 (4%)	3 (1%)	9	36
12	L	220/222 (99%)	213 (97%)	7 (3%)	0	100	100
12	Z	220/222 (99%)	213 (97%)	7 (3%)	0	100	100
13	M	231/246 (94%)	218 (94%)	13 (6%)	0	100	100
13	a	231/246 (94%)	218 (94%)	13 (6%)	0	100	100
14	N	194/196 (99%)	184 (95%)	10 (5%)	0	100	100
14	b	194/196 (99%)	184 (95%)	10 (5%)	0	100	100
15	1	2/4 (50%)	2 (100%)	0	0	100	100
15	2	2/4 (50%)	2 (100%)	0	0	100	100
15	3	2/4 (50%)	2 (100%)	0	0	100	100
15	4	2/4 (50%)	2 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	6292/6630 (95%)	5990 (95%)	267 (4%)	35 (1%)	21	56

5 of 35 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	THR
2	B	51	VAL
3	C	202	GLN
3	C	205	ALA
4	D	2	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	193 (92%)	16 (8%)	12	40
1	O	209/209 (100%)	193 (92%)	16 (8%)	12	40
2	B	203/216 (94%)	178 (88%)	25 (12%)	4	21
2	P	203/216 (94%)	178 (88%)	25 (12%)	4	21
3	C	212/226 (94%)	186 (88%)	26 (12%)	4	21
3	Q	212/226 (94%)	186 (88%)	26 (12%)	4	21
4	D	194/215 (90%)	169 (87%)	25 (13%)	4	19
4	R	194/215 (90%)	169 (87%)	25 (13%)	4	19
5	E	190/193 (98%)	166 (87%)	24 (13%)	4	20
5	S	190/193 (98%)	165 (87%)	25 (13%)	4	18
6	F	201/239 (84%)	186 (92%)	15 (8%)	12	41
6	T	201/239 (84%)	186 (92%)	15 (8%)	12	41
7	G	206/210 (98%)	180 (87%)	26 (13%)	4	20
7	U	206/210 (98%)	180 (87%)	26 (13%)	4	20
8	H	185/190 (97%)	163 (88%)	22 (12%)	5	22
8	V	185/190 (97%)	163 (88%)	22 (12%)	5	22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
9	I	172/173 (99%)	157 (91%)	15 (9%)	9	35
9	W	172/173 (99%)	157 (91%)	15 (9%)	9	35
10	J	173/175 (99%)	152 (88%)	21 (12%)	5	21
10	X	173/175 (99%)	152 (88%)	21 (12%)	5	21
11	K	169/169 (100%)	160 (95%)	9 (5%)	20	54
11	Y	169/169 (100%)	159 (94%)	10 (6%)	18	50
12	L	185/185 (100%)	169 (91%)	16 (9%)	10	36
12	Z	185/185 (100%)	169 (91%)	16 (9%)	10	36
13	M	199/208 (96%)	185 (93%)	14 (7%)	14	44
13	a	199/208 (96%)	185 (93%)	14 (7%)	14	44
14	N	162/162 (100%)	151 (93%)	11 (7%)	14	45
14	b	162/162 (100%)	151 (93%)	11 (7%)	14	45
All	All	5320/5540 (96%)	4788 (90%)	532 (10%)	7	29

5 of 532 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
10	X	17	SER
10	X	126	VAL
10	X	8	ARG
13	a	214	VAL
10	J	3	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 155 such sidechains are listed below:

Mol	Chain	Res	Type
7	U	117	GLN
12	Z	94	GLN
7	U	242	GLN
10	X	86	GLN
13	a	194	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/250 (100%)	-0.73	0 100 100	45, 63, 95, 132	0
1	O	250/250 (100%)	-0.69	0 100 100	46, 67, 106, 132	0
2	B	244/258 (94%)	-0.66	0 100 100	46, 68, 112, 159	0
2	P	244/258 (94%)	-0.63	0 100 100	45, 69, 119, 158	0
3	C	240/254 (94%)	-0.60	0 100 100	44, 69, 120, 139	0
3	Q	240/254 (94%)	-0.52	1 (0%) 88 76	52, 79, 138, 156	0
4	D	235/260 (90%)	-0.65	0 100 100	44, 69, 97, 118	0
4	R	235/260 (90%)	-0.67	0 100 100	50, 73, 102, 129	0
5	E	231/234 (98%)	-0.66	0 100 100	49, 74, 103, 134	0
5	S	231/234 (98%)	-0.59	0 100 100	53, 79, 111, 139	0
6	F	243/288 (84%)	-0.67	0 100 100	47, 68, 109, 142	0
6	T	243/288 (84%)	-0.62	0 100 100	44, 71, 112, 135	0
7	G	241/252 (95%)	-0.71	0 100 100	44, 62, 100, 142	0
7	U	241/252 (95%)	-0.69	0 100 100	46, 63, 95, 121	0
8	H	226/232 (97%)	-0.74	0 100 100	43, 60, 91, 150	0
8	V	226/232 (97%)	-0.77	0 100 100	46, 62, 90, 156	0
9	I	204/205 (99%)	-0.85	0 100 100	42, 61, 88, 112	0
9	W	204/205 (99%)	-0.84	0 100 100	41, 60, 88, 132	0
10	J	195/198 (98%)	-0.77	1 (0%) 87 72	41, 61, 90, 129	0
10	X	195/198 (98%)	-0.73	1 (0%) 87 72	43, 63, 89, 144	0
11	K	212/212 (100%)	-0.75	0 100 100	44, 61, 94, 113	0
11	Y	212/212 (100%)	-0.74	0 100 100	47, 63, 98, 127	0
12	L	222/222 (100%)	-0.79	0 100 100	43, 61, 85, 103	0
12	Z	222/222 (100%)	-0.79	0 100 100	43, 62, 91, 107	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å²)	Q<0.9	
13	M	233/246 (94%)	-0.80	0	100	100	42, 62, 86, 107	0
13	a	233/246 (94%)	-0.79	0	100	100	42, 60, 82, 105	0
14	N	196/196 (100%)	-0.81	0	100	100	41, 57, 84, 121	0
14	b	196/196 (100%)	-0.79	0	100	100	42, 58, 83, 112	0
15	1	2/4 (50%)	0.09	0	100	100	68, 68, 68, 71	0
15	2	2/4 (50%)	-0.63	0	100	100	57, 57, 57, 58	0
15	3	2/4 (50%)	-0.23	0	100	100	77, 77, 77, 81	0
15	4	2/4 (50%)	-0.84	0	100	100	65, 65, 65, 69	0
All	All	6352/6630 (95%)	-0.71	3 (0%)	100	100	41, 65, 103, 159	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
10	X	1	MET	4.4
3	Q	50	LEU	3.3
10	J	1	MET	2.4

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
16	MG	H	301	1/1	0.91	0.12	60,60,60,60	0
16	MG	Z	301	1/1	0.95	0.09	71,71,71,71	0
16	MG	I	302	1/1	0.96	0.12	67,67,67,67	0
16	MG	K	302	1/1	0.97	0.12	54,54,54,54	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
16	MG	N	201	1/1	0.97	0.03	48,48,48,48	0
16	MG	I	301	1/1	0.97	0.05	42,42,42,42	0
16	MG	V	301	1/1	0.98	0.04	68,68,68,68	0
16	MG	Y	301	1/1	0.98	0.06	50,50,50,50	0
16	MG	K	301	1/1	0.98	0.05	45,45,45,45	0
16	MG	G	301	1/1	0.99	0.03	53,53,53,53	0
16	MG	J	201	1/1	1.00	0.08	33,33,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.