



wwPDB EM Validation Summary Report ⓘ

Mar 27, 2026 – 04:38 PM UTC

PDB ID : 8QBG / pdb_00008qbg
EMDB ID : EMD-18312
Title : Stretched state - Pill in native eisosome lattice bound to plasma membrane microdomain
Authors : Kefauver, J.M.; Zou, L.; Desfosses, A.; Loewith, R.J.
Deposited on : 2023-08-24
Resolution : 3.64 Å(reported)
Based on initial model : .

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	11633 (3.14 - 4.14)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	339	5% 56% 24% 20%
1	B	339	6% 55% 25% 20%
1	C	339	16% 56% 24% 20%
1	D	339	48% 55% 24% 20%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	339	
1	F	339	
1	G	339	
1	H	339	
1	I	339	
1	J	339	
1	K	339	
1	L	339	
1	M	339	
1	N	339	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 30058 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Sphingolipid long chain base-responsive protein PIL1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	271	Total 2147	C 1346	N 374	O 425	S 2	0	0
1	B	271	Total 2147	C 1346	N 374	O 425	S 2	0	0
1	C	271	Total 2147	C 1346	N 374	O 425	S 2	0	0
1	I	271	Total 2147	C 1346	N 374	O 425	S 2	0	0
1	D	271	Total 2147	C 1346	N 374	O 425	S 2	0	0
1	J	271	Total 2147	C 1346	N 374	O 425	S 2	0	0
1	E	271	Total 2147	C 1346	N 374	O 425	S 2	0	0
1	K	271	Total 2147	C 1346	N 374	O 425	S 2	0	0
1	F	271	Total 2147	C 1346	N 374	O 425	S 2	0	0
1	L	271	Total 2147	C 1346	N 374	O 425	S 2	0	0
1	G	271	Total 2147	C 1346	N 374	O 425	S 2	0	0
1	M	271	Total 2147	C 1346	N 374	O 425	S 2	0	0
1	H	271	Total 2147	C 1346	N 374	O 425	S 2	0	0
1	N	271	Total 2147	C 1346	N 374	O 425	S 2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Spingolipid long chain base-responsive protein PIL1




- Molecule 1: Spingolipid long chain base-responsive protein PIL1




- Molecule 1: Spingolipid long chain base-responsive protein PIL1







• Molecule 1: Spingolipid long chain base-responsive protein PIL1




• Molecule 1: Spingolipid long chain base-responsive protein PIL1






• Molecule 1: Spingolipid long chain base-responsive protein PIL1




• Molecule 1: Spingolipid long chain base-responsive protein PIL1





- Molecule 1: Sphingolipid long chain base-responsive protein PIL1




- Molecule 1: Sphingolipid long chain base-responsive protein PIL1




Chain L:






• Molecule 1: Spingolipid long chain base-responsive protein PIL1



• Molecule 1: Spingolipid long chain base-responsive protein PIL1



GLU	T250	A185	Y125
PRO	T251	Q186	R126
GLU	T252	L187	L127
GLN	D253	S188	T128
GLU	A254	N189	L129
GLY	E255	I190	K130
GLY	S256	T191	S131
ALA	A257	R192	I132
VAL	L258	S193	R133
GLU	N259	K194	D134
GLU	E260	L195	I135
HIS	W261	R196	E136
GLN	T262	A197	G137
GLU	L263	A198	S138
GLY	D264	F199	Y139
HIS	S265	W200	Q140
GLN	G266	Y201	P141
GLN	A267	Q202	S142
SER	V268	H209	R143
SER	K269	P203	R144
PRO	T270	D204	D144
GLN	T271	S205	R145
THR	LEU	I206	K146
THR	SER	I207	D147
ALA	PHE	E208	K148
	LYS	H209	I149
	GLN	S210	T150
	ASP	E211	D151
	TYR	K212	K152
	GLU	I213	I153
	ASP	A214	A154
	PHE	L215	Y155
	GLU	L216	L156
	GLU	A217	K157
	GLY	G218	Y158
	GLU	Y219	K159
	GLU	G220	D160
	GLU	K221	P161
	GLU	A222	Q162
	GLU	L223	P164
	ASP	L224	K165
	GLN	E225	L166
	GLY	L226	E167
	ARG	L227	V168
	TRP	D228	L169
	SER	D229	E170
	GLU	S230	Q171
	ASP	P231	E172
	GLU		L173
	GLU	E236	V174
	GLY	T237	R175
	GLY	R238	A176
	ILE	P239	E177
	GLU	A240	A178
		Y241	E179
		D242	S180
		E245	L181
		A246	V182
			A183
		Q249	F184

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	77457	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	3.415	Depositor
Minimum map value	-2.454	Depositor
Average map value	0.016	Depositor
Map value standard deviation	0.102	Depositor
Recommended contour level	0.55	Depositor
Map size (Å)	339.712, 339.712, 339.712	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.327, 1.327, 1.327	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.16	0/2184	0.37	0/2952
1	B	0.16	0/2184	0.37	0/2952
1	C	0.16	0/2184	0.37	0/2952
1	D	0.16	0/2184	0.37	0/2952
1	E	0.16	0/2184	0.37	0/2952
1	F	0.16	0/2184	0.37	0/2952
1	G	0.16	0/2184	0.37	0/2952
1	H	0.16	0/2184	0.37	0/2952
1	I	0.16	0/2184	0.37	0/2952
1	J	0.16	0/2184	0.37	0/2952
1	K	0.16	0/2184	0.37	0/2952
1	L	0.16	0/2184	0.37	0/2952
1	M	0.16	0/2184	0.37	0/2952
1	N	0.16	0/2184	0.37	0/2952
All	All	0.16	0/30576	0.37	0/41328

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2147	0	2150	62	0
1	B	2147	0	2150	66	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2147	0	2150	59	0
1	D	2147	0	2150	62	0
1	E	2147	0	2150	60	0
1	F	2147	0	2150	59	0
1	G	2147	0	2150	60	0
1	H	2147	0	2150	63	0
1	I	2147	0	2150	63	0
1	J	2147	0	2150	67	0
1	K	2147	0	2150	64	0
1	L	2147	0	2150	62	0
1	M	2147	0	2150	58	0
1	N	2147	0	2150	62	0
All	All	30058	0	30100	783	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 783 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:165:LYS:HE2	1:A:165:LYS:HA	1.73	0.71
1:F:165:LYS:HE2	1:F:165:LYS:HA	1.73	0.71
1:J:165:LYS:HE2	1:J:165:LYS:HA	1.73	0.71
1:E:165:LYS:HE2	1:E:165:LYS:HA	1.73	0.71
1:L:165:LYS:HA	1:L:165:LYS:HE2	1.73	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	269/339 (79%)	263 (98%)	6 (2%)	0	100	100
1	B	269/339 (79%)	263 (98%)	6 (2%)	0	100	100
1	C	269/339 (79%)	263 (98%)	6 (2%)	0	100	100
1	D	269/339 (79%)	263 (98%)	6 (2%)	0	100	100
1	E	269/339 (79%)	263 (98%)	6 (2%)	0	100	100
1	F	269/339 (79%)	263 (98%)	6 (2%)	0	100	100
1	G	269/339 (79%)	263 (98%)	6 (2%)	0	100	100
1	H	269/339 (79%)	263 (98%)	6 (2%)	0	100	100
1	I	269/339 (79%)	263 (98%)	6 (2%)	0	100	100
1	J	269/339 (79%)	263 (98%)	6 (2%)	0	100	100
1	K	269/339 (79%)	263 (98%)	6 (2%)	0	100	100
1	L	269/339 (79%)	263 (98%)	6 (2%)	0	100	100
1	M	269/339 (79%)	263 (98%)	6 (2%)	0	100	100
1	N	269/339 (79%)	263 (98%)	6 (2%)	0	100	100
All	All	3766/4746 (79%)	3682 (98%)	84 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/294 (80%)	234 (100%)	0	100	100
1	B	234/294 (80%)	233 (100%)	1 (0%)	84	79
1	C	234/294 (80%)	233 (100%)	1 (0%)	84	79
1	D	234/294 (80%)	234 (100%)	0	100	100
1	E	234/294 (80%)	233 (100%)	1 (0%)	84	79
1	F	234/294 (80%)	233 (100%)	1 (0%)	84	79
1	G	234/294 (80%)	234 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	234/294 (80%)	234 (100%)	0	100	100
1	I	234/294 (80%)	234 (100%)	0	100	100
1	J	234/294 (80%)	234 (100%)	0	100	100
1	K	234/294 (80%)	234 (100%)	0	100	100
1	L	234/294 (80%)	234 (100%)	0	100	100
1	M	234/294 (80%)	234 (100%)	0	100	100
1	N	234/294 (80%)	234 (100%)	0	100	100
All	All	3276/4116 (80%)	3272 (100%)	4 (0%)	87	87

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	20	ASN
1	C	20	ASN
1	E	20	ASN
1	F	20	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	202	GLN
1	N	202	GLN
1	K	202	GLN
1	F	202	GLN
1	L	202	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-18312. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 126



Y Index: 128




Z Index: 111


6.3.2 Raw map



X Index: 130



Y Index: 128




Z Index: 110


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.55. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_18312_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 184 nm^3 ; this corresponds to an approximate mass of 166 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.275 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of $0.275 \text{ \AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.64	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	6.44	8.16	6.70

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.44 differs from the reported value 3.64 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-18312 and PDB model 8QBG. Per-residue inclusion information can be found in [section 3](#) on [page 5](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.55 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.55).

9.4 Atom inclusion [i](#)



At the recommended contour level, 60% of all backbone atoms, 44% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.55) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.4450	0.2170
A	0.7760	0.3990
B	0.7550	0.3860
C	0.6030	0.2870
D	0.3320	0.1600
E	0.1280	0.0860
F	0.5590	0.2750
G	0.4330	0.2070
H	0.2770	0.1170
I	0.5700	0.2690
J	0.4110	0.1920
K	0.2590	0.1150
L	0.5970	0.2790
M	0.3640	0.1740
N	0.1630	0.0950

1.0

0.0

<0.0