



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:41 AM UTC

PDB ID : 1QCX / pdb_00001qcx
Title : PECTIN LYASE B
Authors : Vitali, J.; Jurnak, F.
Deposited on : 1999-05-13
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	359	 84% 15% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3001 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PECTIN LYASE B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	359	Total	C	N	O	S	0	0	0
			2662	1663	462	527	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	GLN	ARG	SEE REMARK 999	UNP Q00205
A	96	TYR	-	insertion	UNP Q00205
A	281	VAL	ILE	SEE REMARK 999	UNP Q00205
A	346	ALA	ASN	SEE REMARK 999	UNP Q00205

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	339	Total	O	0	0
			339	339		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: PECTIN LYASE B

Chain A:  84% 15%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	83.70Å 88.80Å 42.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.00 – 1.70	Depositor
% Data completeness (in resolution range)	91.8 (7.00-1.70)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.162 , 0.198	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3001	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	1/2719 (0.0%)	1.19	39/3705 (1.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	237	MET	CG-SD	8.10	2.00	1.80

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	149	TYR	N-CA-C	8.97	123.14	108.52
1	A	345	GLY	N-CA-C	8.58	123.30	112.83
1	A	101	ILE	N-CA-C	7.46	117.58	110.42
1	A	347	ILE	N-CA-C	7.27	117.40	110.42
1	A	304	ALA	N-CA-C	7.11	119.64	111.11
1	A	223	VAL	N-CA-C	6.97	117.92	108.17
1	A	226	LYS	N-CA-C	6.87	120.71	109.24
1	A	295	PRO	N-CA-C	6.84	122.89	113.65
1	A	231	TYR	N-CA-C	6.83	120.06	109.07
1	A	10	PHE	N-CA-C	6.75	119.50	111.33
1	A	150	VAL	N-CA-C	-6.58	101.08	109.30
1	A	98	LYS	N-CA-C	6.49	119.17	111.71
1	A	177	GLN	N-CA-C	6.42	119.78	110.28
1	A	269	VAL	N-CA-C	6.30	116.93	108.11
1	A	144	ASP	N-CA-C	6.21	118.78	111.02
1	A	12	HIS	N-CA-C	6.11	119.57	111.75
1	A	130	SER	N-CA-C	6.11	119.81	111.39
1	A	253	PHE	N-CA-C	-6.08	98.62	108.52
1	A	178	HIS	N-CA-C	-6.05	104.24	111.69
1	A	249	VAL	CB-CA-C	-6.02	101.65	110.62
1	A	192	SER	N-CA-C	5.82	118.26	109.23
1	A	76	ILE	N-CA-C	-5.77	101.37	109.21
1	A	80	SER	N-CA-C	5.72	119.45	111.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	123	GLY	N-CA-C	5.69	122.65	115.42
1	A	314	ASN	N-CA-C	-5.65	101.07	110.17
1	A	232	ASN	N-CA-C	5.61	119.37	111.52
1	A	294	SER	CA-C-N	5.49	125.31	119.28
1	A	294	SER	C-N-CA	5.49	125.31	119.28
1	A	239	LYS	N-CA-C	-5.33	100.04	108.73
1	A	250	ASN	N-CA-C	5.30	118.51	111.30
1	A	249	VAL	N-CA-C	5.28	116.09	108.48
1	A	85	TYR	N-CA-C	5.25	119.96	113.50
1	A	113	VAL	N-CA-C	5.25	115.46	108.11
1	A	129	VAL	N-CA-C	5.23	115.20	108.82
1	A	308	GLY	N-CA-C	-5.19	108.16	115.32
1	A	42	VAL	N-CA-C	-5.16	98.51	106.72
1	A	261	PHE	N-CA-C	5.16	117.91	110.28
1	A	188	ARG	N-CA-C	5.10	116.57	108.67
1	A	118	LYS	N-CA-C	5.02	119.45	113.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2662	0	2559	16	0
2	A	339	0	0	2	0
All	All	3001	0	2559	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:210:HIS:HD2	2:A:834:HOH:O	1.93	0.51
1:A:112:ILE:HD12	1:A:137:ILE:HG12	1.93	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:353:LYS:HG3	1:A:354:ASN:ND2	2.29	0.48
1:A:176:ARG:O	1:A:210:HIS:HE1	1.98	0.47
1:A:71:GLN:HG3	1:A:206:CYS:HB3	1.98	0.46
1:A:129:VAL:HG12	1:A:159:ASP:HB3	1.98	0.46
1:A:294:SER:HA	1:A:300:ASN:ND2	2.31	0.46
1:A:74:VAL:HG22	1:A:148:LYS:HE2	1.97	0.46
1:A:294:SER:CA	1:A:300:ASN:HD21	2.29	0.46
1:A:294:SER:C	1:A:300:ASN:HD21	2.24	0.46
1:A:284:GLU:HG3	2:A:579:HOH:O	2.16	0.45
1:A:263:ILE:HB	1:A:287:ILE:HG23	1.99	0.44
1:A:294:SER:HA	1:A:300:ASN:HD21	1.82	0.44
1:A:177:GLN:NE2	1:A:236:ARG:HH11	2.16	0.43
1:A:249:VAL:HG13	1:A:272:GLU:HG3	2.00	0.43
1:A:176:ARG:O	1:A:210:HIS:CE1	2.73	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	357/359 (99%)	338 (95%)	19 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	285/285 (100%)	279 (98%)	6 (2%)	47 30

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	232	ASN
1	A	240	VAL
1	A	249	VAL
1	A	270	LEU
1	A	306	VAL
1	A	357	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	ASN
1	A	177	GLN
1	A	187	ASN
1	A	210	HIS
1	A	255	ASN
1	A	300	ASN
1	A	354	ASN
1	A	357	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.