



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:38 PM UTC

PDB ID : 3QC4 / pdb_00003qc4
Title : PDK1 in complex with DFG-OUT inhibitor xxx
Authors : Arndt, J.W.
Deposited on : 2011-01-15
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	8% 65% 20% • 12%
1	B	314	10% 70% 16% • 11%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5010 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3-phosphoinositide-dependent protein kinase 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	277	Total	C	N	O	P	S	0	2	0
			2268	1474	373	413	1	7			
1	B	279	Total	C	N	O	P	S	0	2	0
			2241	1453	366	413	1	8			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	expression tag	UNP O15530
A	-4	PRO	-	expression tag	UNP O15530
A	-3	LEU	-	expression tag	UNP O15530
A	-2	GLY	-	expression tag	UNP O15530
A	-1	SER	-	expression tag	UNP O15530
A	303	ALA	GLU	engineered mutation	UNP O15530
A	304	ALA	LYS	engineered mutation	UNP O15530
B	-5	GLY	-	expression tag	UNP O15530
B	-4	PRO	-	expression tag	UNP O15530
B	-3	LEU	-	expression tag	UNP O15530
B	-2	GLY	-	expression tag	UNP O15530
B	-1	SER	-	expression tag	UNP O15530
B	303	ALA	GLU	engineered mutation	UNP O15530
B	304	ALA	LYS	engineered mutation	UNP O15530

- Molecule 2 is 1-(3,4-difluorobenzyl)-2-oxo-N-{(1R)-2-[(2-oxo-2,3-dihydro-1H-benzimidazol-5-yl)oxy]-1-phenylethyl}-1,2-dihydropyridine-3-carboxamide (CCD ID: MP7) (formula: C₂₈H₂₂F₂N₄O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			38	28	2	4	4		
2	B	1	Total	C	F	N	O	0	0
			38	28	2	4	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	209	Total	O	0	0
			209	209		
3	B	216	Total	O	0	0
			216	216		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-phosphoinositide-dependent protein kinase 1



- Molecule 1: 3-phosphoinositide-dependent protein kinase 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	183.15Å 57.34Å 83.69Å 90.00° 114.13° 90.00°	Depositor
Resolution (Å)	30.49 – 1.80 30.49 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.0 (30.49-1.80) 98.0 (30.49-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.82 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.172 , 0.213 0.191 , 0.229	Depositor DCC
R_{free} test set	3620 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	25.9	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5010	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MP7, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.63	22/2318 (0.9%)	1.29	11/3124 (0.4%)
1	B	1.64	21/2287 (0.9%)	1.29	4/3082 (0.1%)
All	All	1.64	43/4605 (0.9%)	1.29	15/6206 (0.2%)

All (43) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	294	ILE	CA-CB	8.61	1.64	1.54
1	A	288	TYR	C-O	-7.46	1.15	1.24
1	B	319	LEU	N-CA	6.76	1.54	1.46
1	A	343	GLU	C-O	6.66	1.31	1.24
1	A	196	LEU	C-O	-6.50	1.16	1.24
1	B	288	TYR	CA-C	6.39	1.60	1.52
1	B	227	ALA	CA-CB	6.39	1.64	1.53
1	B	205	ASP	C-O	-6.29	1.17	1.24
1	A	271	ILE	CA-CB	6.13	1.62	1.54
1	A	162	ALA	CA-CB	6.11	1.60	1.53
1	B	276	VAL	C-O	5.99	1.30	1.24
1	A	171	ILE	CA-C	5.95	1.60	1.52
1	B	258	SER	C-O	-5.94	1.16	1.23
1	B	305	PHE	N-CA	5.92	1.53	1.46
1	A	309	ALA	CA-CB	5.92	1.62	1.53
1	B	249	VAL	CA-C	5.90	1.60	1.52
1	A	308	LYS	N-CA	5.86	1.53	1.46
1	B	185	TYR	N-CA	5.74	1.53	1.46
1	B	354	THR	C-O	-5.65	1.17	1.24
1	A	250	SER	C-O	-5.62	1.16	1.23
1	A	203	HIS	C-O	-5.61	1.17	1.24
1	A	165	GLY	C-O	-5.60	1.15	1.23
1	A	220	GLN	N-CA	-5.59	1.39	1.46
1	A	246	ALA	C-O	-5.55	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	227	ALA	CA-CB	5.51	1.62	1.53
1	A	286	ASN	C-O	-5.41	1.17	1.23
1	B	303	ALA	CA-CB	5.38	1.61	1.53
1	B	338	ALA	CA-CB	5.31	1.62	1.53
1	A	222	THR	N-CA	-5.30	1.39	1.45
1	A	124	VAL	CA-CB	-5.26	1.47	1.54
1	B	221	ILE	C-O	5.25	1.31	1.24
1	B	206	LEU	C-O	-5.21	1.17	1.23
1	A	283	ARG	C-O	-5.14	1.17	1.23
1	B	181	CYS	N-CA	5.14	1.52	1.46
1	B	346	THR	CA-CB	5.10	1.59	1.53
1	A	207	LYS	C-O	5.10	1.31	1.23
1	B	262	SER	N-CA	5.08	1.52	1.46
1	B	53	GLY	C-O	5.08	1.28	1.23
1	B	195	TYR	N-CA	5.08	1.52	1.46
1	B	304	ALA	C-O	5.07	1.30	1.23
1	A	206	LEU	C-O	-5.04	1.17	1.23
1	A	143	VAL	CA-CB	5.02	1.59	1.54
1	A	134	MET	CA-C	-5.01	1.45	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	272	ILE	N-CA-C	-7.66	102.98	110.72
1	A	349	ASN	N-CA-C	6.40	119.07	110.88
1	A	154	LYS	N-CA-C	6.21	118.85	109.23
1	B	133	VAL	CB-CA-C	-5.93	104.14	112.14
1	B	247	GLN	CA-CB-CG	-5.81	102.48	114.10
1	A	345	VAL	N-CA-C	5.77	116.52	109.30
1	A	355	PRO	O-C-N	5.59	123.78	121.15
1	B	203	HIS	N-CA-C	5.58	117.17	111.14
1	A	78	ARG	N-CA-C	5.48	114.61	108.25
1	A	343	GLU	N-CA-C	5.44	117.64	111.11
1	A	293	LYS	CA-CB-CG	-5.14	103.83	114.10
1	A	245	THR	N-CA-C	-5.12	100.68	109.24
1	A	308	LYS	CA-CB-CG	-5.03	104.04	114.10
1	A	285	GLY	N-CA-C	5.02	119.82	113.24
1	A	102	LEU	N-CA-C	5.00	116.73	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2268	0	2282	47	0
1	B	2241	0	2241	31	0
2	A	38	0	22	0	0
2	B	38	0	22	0	0
3	A	209	0	0	2	0
3	B	216	0	0	2	0
All	All	5010	0	4567	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:228:LYS:NZ	1:B:239:ALA:HA	1.43	1.29
1:A:78:ARG:HG3	1:A:78:ARG:HH11	1.01	1.13
1:A:131:ARG:HG2	1:A:131:ARG:O	1.44	1.07
1:B:228:LYS:HZ1	1:B:239:ALA:HA	1.00	0.94
1:B:118:ILE:HG23	1:B:118:ILE:O	1.66	0.93
1:A:78:ARG:HG3	1:A:78:ARG:NH1	1.81	0.90
1:B:228:LYS:HZ2	1:B:239:ALA:HA	1.46	0.80
1:B:118:ILE:O	1:B:118:ILE:CG2	2.30	0.80
1:A:328:GLU:HG2	3:A:488:HOH:O	1.82	0.77
1:A:112:ILE:HD12	1:A:156:TYR:CD2	2.20	0.77
1:B:228:LYS:HZ1	1:B:239:ALA:CA	1.90	0.76
1:B:228:LYS:NZ	1:B:239:ALA:CA	2.39	0.74
1:A:78:ARG:HH11	1:A:78:ARG:CG	1.93	0.73
1:A:134:MET:HE3	1:A:229:VAL:HG23	1.71	0.73
1:A:76:LYS:CE	1:A:76:LYS:HA	2.20	0.72
1:A:78:ARG:HB2	1:A:79:PRO:HD2	1.74	0.70
1:A:134:MET:HE3	1:A:229:VAL:CG2	2.20	0.70
1:B:154:LYS:HD2	1:B:156:TYR:CZ	2.27	0.70
1:B:154:LYS:HD2	1:B:156:TYR:OH	1.91	0.69
1:A:128:THR:O	1:A:128:THR:HG22	1.93	0.67
1:A:76:LYS:HA	1:A:76:LYS:HE2	1.78	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:134:MET:HE2	1:A:201:ILE:HG23	1.77	0.66
1:A:131:ARG:O	1:A:131:ARG:CG	2.30	0.65
1:B:81:ASP:OD1	1:B:81:ASP:N	2.30	0.65
1:A:104:THR:O	1:A:105:SER:CB	2.44	0.64
1:B:228:LYS:CE	1:B:239:ALA:HA	2.28	0.63
1:A:115:LYS:NZ	1:A:150:GLN:HE21	1.96	0.62
1:A:114:GLU:HA	1:A:154:LYS:HG2	1.81	0.62
1:A:104:THR:O	1:A:105:SER:HB2	1.99	0.61
1:A:128:THR:O	1:A:128:THR:CG2	2.48	0.60
1:B:77:LYS:C	1:B:78:ARG:HG3	2.26	0.60
1:A:125:PRO:O	1:A:126:TYR:HB2	2.00	0.59
1:A:112:ILE:HD12	1:A:156:TYR:HD2	1.67	0.58
1:B:77:LYS:HD3	1:B:82:PHE:HZ	1.66	0.58
1:A:110:ILE:HG22	1:A:112:ILE:CD1	2.36	0.56
1:B:111:LYS:HB3	1:B:157:PHE:HB2	1.88	0.55
1:A:119:ILE:HD12	1:A:121:GLU:HG2	1.87	0.55
1:A:110:ILE:HG22	1:A:112:ILE:HD13	1.89	0.54
1:B:117:HIS:O	1:B:118:ILE:C	2.49	0.54
1:B:143:VAL:HG21	1:B:222[A]:THR:HG22	1.90	0.53
1:B:199:LYS:HE2	3:B:364:HOH:O	2.06	0.53
1:B:87:ILE:HD12	1:B:87:ILE:H	1.73	0.52
1:B:113:LEU:HD23	1:B:155:LEU:HB2	1.92	0.52
1:A:76:LYS:HE2	1:A:76:LYS:CA	2.39	0.52
1:A:134:MET:HE1	1:A:228:LYS:C	2.35	0.52
1:A:76:LYS:HA	1:A:76:LYS:HE3	1.92	0.51
1:B:204:ARG:NH1	1:B:228:LYS:HE2	2.26	0.51
1:A:249:VAL:CG2	1:A:253:LEU:HD23	2.41	0.50
1:A:115:LYS:HZ1	1:A:150:GLN:HE21	1.59	0.50
1:A:78:ARG:NH1	1:A:78:ARG:CG	2.62	0.50
1:A:93:PHE:CE1	1:A:122:ASN:ND2	2.80	0.50
1:B:154:LYS:HG3	1:B:156:TYR:CE1	2.47	0.50
1:A:132:ASP:O	1:A:132:ASP:OD1	2.30	0.49
1:A:76:LYS:CE	1:A:76:LYS:CA	2.91	0.48
1:A:134:MET:HE1	1:A:229:VAL:N	2.30	0.47
1:B:106:ARG:NH1	3:B:542:HOH:O	2.47	0.47
1:B:77:LYS:HD3	1:B:82:PHE:CZ	2.48	0.46
1:A:134:MET:HE3	1:A:229:VAL:HG22	1.95	0.45
1:B:118:ILE:HD12	1:B:118:ILE:HA	1.59	0.45
1:A:229:VAL:HG12	1:A:230:LEU:O	2.16	0.45
1:A:78:ARG:HB2	1:A:79:PRO:CD	2.44	0.45
1:B:113:LEU:HD21	1:B:155:LEU:HD12	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:151:ASP:OD1	1:A:154:LYS:N	2.41	0.44
1:B:139:HIS:CG	1:B:140:PRO:HD2	2.53	0.43
1:A:100:ARG:HG3	1:A:105:SER:O	2.19	0.43
1:A:285:GLY:HA3	1:B:257:LYS:HD3	1.99	0.43
1:A:328:GLU:CG	3:A:488:HOH:O	2.50	0.43
1:A:139:HIS:CG	1:A:140:PRO:HD2	2.53	0.43
1:A:348:GLU:H	1:A:348:GLU:CD	2.25	0.43
1:B:228:LYS:HZ2	1:B:239:ALA:CA	2.22	0.43
1:A:214:ASN:HB3	1:A:220:GLN:OE1	2.20	0.42
1:B:79:PRO:HD3	1:B:149:PHE:CE1	2.55	0.41
1:B:215:GLU:HG2	1:B:216:ASP:N	2.36	0.41
1:A:100:ARG:HD3	1:A:105:SER:HA	2.03	0.41
1:A:257:LYS:HD3	1:B:285:GLY:HA3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/314 (87%)	266 (97%)	8 (3%)	0	100	100
1	B	272/314 (87%)	268 (98%)	4 (2%)	0	100	100
All	All	546/628 (87%)	534 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/269 (91%)	229 (94%)	15 (6%)	17	6
1	B	239/269 (89%)	226 (95%)	13 (5%)	20	8
All	All	483/538 (90%)	455 (94%)	28 (6%)	18	7

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	76	LYS
1	A	78	ARG
1	A	87	ILE
1	A	92	SER
1	A	105	SER
1	A	120	LYS
1	A	121	GLU
1	A	123	LYS
1	A	131	ARG
1	A	135	SER
1	A	143	VAL
1	A	153	GLU
1	A	224	PHE
1	A	328	GLU
1	A	340	PRO
1	B	-1	SER
1	B	78	ARG
1	B	81	ASP
1	B	87	ILE
1	B	113	LEU
1	B	115	LYS
1	B	118	ILE
1	B	127	VAL
1	B	143	VAL
1	B	154	LYS
1	B	224	PHE
1	B	228	LYS
1	B	359	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	150	GLN
1	A	292	GLN
1	B	117	HIS
1	B	150	GLN
1	B	292	GLN
1	B	349	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SEP	B	241	1	8,9,10	1.11	0	7,12,14	1.98	3 (42%)
1	SEP	A	241	1	8,9,10	1.31	1 (12%)	7,12,14	1.36	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	B	241	1	-	1/6/8/10	-
1	SEP	A	241	1	-	1/6/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	241	SEP	P-O1P	2.99	1.59	1.50

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	241	SEP	OG-CB-CA	3.13	111.19	108.14
1	B	241	SEP	O2P-P-OG	2.91	114.25	106.67
1	B	241	SEP	O3P-P-O2P	2.50	117.18	107.80
1	A	241	SEP	O2P-P-O1P	-2.11	102.63	110.83

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	241	SEP	CA-CB-OG-P
1	B	241	SEP	CA-CB-OG-P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MP7	A	1	-	42,42,42	1.38	7 (16%)	57,59,59	2.09	13 (22%)
2	MP7	B	2	-	42,42,42	1.50	9 (21%)	57,59,59	2.32	14 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MP7	A	1	-	-	4/21/21/21	0/5/5/5
2	MP7	B	2	-	-	2/21/21/21	0/5/5/5

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	MP7	C30-N20	-4.01	1.31	1.37
2	A	1	MP7	O31-C30	3.36	1.30	1.23
2	B	2	MP7	C8-C7	3.16	1.43	1.38
2	B	2	MP7	C7-C6	2.95	1.44	1.38
2	B	2	MP7	C18-C17	2.83	1.43	1.38
2	A	1	MP7	C3-C2	2.49	1.40	1.35
2	A	1	MP7	C17-C16	2.47	1.43	1.38
2	A	1	MP7	C14-N20	-2.45	1.34	1.38
2	B	2	MP7	C28-C27	2.25	1.43	1.38
2	B	2	MP7	C24-C23	-2.23	1.49	1.52
2	B	2	MP7	C1-C	2.22	1.42	1.38
2	B	2	MP7	O31-C30	2.17	1.28	1.23
2	B	2	MP7	C30-N20	-2.14	1.34	1.37
2	B	2	MP7	C22-C23	-2.10	1.49	1.52
2	A	1	MP7	C13-N19	-2.02	1.35	1.38
2	A	1	MP7	F12-C9	-2.01	1.30	1.35

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	MP7	N20-C30-N19	7.35	114.97	106.62
2	B	2	MP7	C6-C11-C10	-6.77	114.95	119.37
2	A	1	MP7	C1-C-C33	6.43	122.84	117.94
2	A	1	MP7	N20-C30-N19	6.25	113.73	106.62
2	B	2	MP7	C14-N20-C30	-5.42	104.11	110.03
2	A	1	MP7	C13-N19-C30	-5.34	104.20	110.03
2	B	2	MP7	C7-C8-C9	-4.69	113.26	119.04
2	B	2	MP7	C13-N19-C30	-4.60	105.01	110.03
2	A	1	MP7	O31-C30-N20	-4.56	121.48	126.68
2	A	1	MP7	C5-N-C4	-4.07	113.52	117.73
2	B	2	MP7	O31-C30-N20	-3.99	122.14	126.68
2	B	2	MP7	C5-C6-C11	-3.91	112.91	120.27
2	B	2	MP7	O31-C30-N19	-3.37	122.84	126.68
2	A	1	MP7	C-C33-N32	3.28	120.52	116.25
2	B	2	MP7	C4-C-C33	3.19	125.15	121.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	MP7	C-C4-N	3.13	119.31	114.52
2	A	1	MP7	C-C4-N	3.11	119.29	114.52
2	A	1	MP7	O34-C33-C	-2.94	115.91	121.04
2	A	1	MP7	C2-C3-N	-2.75	117.99	121.31
2	A	1	MP7	C5-C6-C11	-2.58	115.40	120.27
2	A	1	MP7	C14-N20-C30	-2.55	107.25	110.03
2	B	2	MP7	C5-C6-C7	2.52	125.39	120.75
2	A	1	MP7	C7-C6-C11	2.37	121.82	118.55
2	B	2	MP7	C7-C6-C11	2.29	121.70	118.55
2	B	2	MP7	C29-C24-C25	2.26	121.10	118.30
2	B	2	MP7	O-C4-N	-2.23	117.57	120.53
2	A	1	MP7	C5-N-C3	2.12	123.02	119.28

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1	MP7	N32-C23-C24-C25
2	B	2	MP7	N32-C23-C24-C25
2	B	2	MP7	N32-C23-C24-C29
2	A	1	MP7	C22-C23-C24-C29
2	A	1	MP7	N32-C23-C24-C29
2	A	1	MP7	C22-C23-C24-C25

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand MP7 A 1




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/314 (87%)	0.33	25 (9%) 15 13	17, 32, 95, 116	2 (0%)
1	B	278/314 (88%)	0.33	32 (11%) 9 8	13, 29, 94, 153	2 (0%)
All	All	554/628 (88%)	0.33	57 (10%) 12 10	13, 31, 95, 153	4 (0%)

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	118	ILE	10.7
1	B	-1	SER	4.7
1	B	129	ARG	4.7
1	B	93	PHE	4.5
1	B	116	ARG	4.4
1	B	117	HIS	4.2
1	B	359	THR	3.5
1	A	102	LEU	3.5
1	A	131	ARG	3.5
1	B	105	SER	3.4
1	A	359	THR	3.3
1	B	-4	PRO	3.2
1	B	113	LEU	3.2
1	B	102	LEU	3.1
1	A	119	ILE	3.1
1	A	93	PHE	3.1
1	A	130	GLU	3.0
1	B	231	SER	3.0
1	A	132	ASP	3.0
1	A	133	VAL	2.9
1	A	126	TYR	2.9
1	A	83	LYS	2.9
1	B	239	ALA	2.8
1	A	231	SER	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	152	ASP	2.7
1	B	127	VAL	2.7
1	B	87	ILE	2.7
1	A	300	ASP	2.7
1	B	128	THR	2.6
1	B	78	ARG	2.6
1	A	123	LYS	2.5
1	A	122	ASN	2.5
1	B	155	LEU	2.5
1	A	129	ARG	2.5
1	A	239	ALA	2.5
1	B	92	SER	2.5
1	B	103	ALA	2.4
1	B	91	GLY	2.4
1	A	104	THR	2.4
1	B	133	VAL	2.4
1	B	115	LYS	2.3
1	B	148	THR	2.3
1	A	116	ARG	2.3
1	B	52	ASP	2.3
1	A	78	ARG	2.3
1	A	127	VAL	2.3
1	B	-2	GLY	2.2
1	A	118	ILE	2.2
1	A	128	THR	2.2
1	B	104	THR	2.2
1	A	82	PHE	2.1
1	B	76	LYS	2.1
1	B	304	ALA	2.1
1	A	152	ASP	2.1
1	B	114	GLU	2.0
1	A	149	PHE	2.0
1	B	112	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	SEP	A	241	10/11	0.97	0.08	34,41,43,46	0
1	SEP	B	241	10/11	0.98	0.08	27,31,32,33	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MP7	A	1	38/38	0.96	0.07	30,35,43,46	0
2	MP7	B	2	38/38	0.96	0.06	24,31,36,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around MP7 A 1:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.