



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 09:00 PM UTC

PDB ID : 9QCK / pdb_00009qck
Title : Yeast 20S proteasome mutant: beta1_G128V (b1-propeptide deleted)
Authors : Huber, E.M.; Heinemeyer, W.; Groll, M.
Deposited on : 2025-03-04
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	% 96% .
1	O	250	% 94% 5%
2	B	258	% 84% 8% 7%
2	P	258	2% 87% 7% 5%
3	C	254	% 88% 5% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Q	254	 2% 88% 6%
4	D	260	 84% 6% 10%
4	R	260	 84% 6% 10%
5	E	234	 91% 7%
5	S	234	 91% 6%
6	F	288	 78% 6% 16%
6	T	288	 78% 6% 16%
7	G	252	 87% 8%
7	U	252	 87% 8%
8	H	232	 89% 6%
8	V	232	 89% 6%
9	I	205	 93% 6%
9	W	205	 93% 6%
10	J	198	 80% 15%
10	X	198	 82% 13%
11	K	212	 91% 7%
11	Y	212	 89% 8%
12	L	222	 92% 7%
12	Z	222	 93% 6%
13	M	246	 79% 13% 5%
13	a	246	 78% 13% 5%
14	N	196	 3% 79% 18%
14	b	196	 78% 18%

2 Entry composition

There are 18 unique types of molecules in this entry. The entry contains 49670 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit alpha type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			
1	O	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			

- Molecule 2 is a protein called Proteasome subunit alpha type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	241	Total	C	N	O	S	0	0	0
			1887	1192	317	375	3			
2	P	244	Total	C	N	O	S	0	0	0
			1904	1201	321	379	3			

- Molecule 3 is a protein called Proteasome subunit alpha type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	240	Total	C	N	O	S	0	0	0
			1881	1176	329	372	4			
3	Q	240	Total	C	N	O	S	0	0	0
			1881	1176	329	372	4			

- Molecule 4 is a protein called Proteasome subunit alpha type-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	235	Total	C	N	O	S	0	0	0
			1813	1136	304	366	7			
4	R	235	Total	C	N	O	S	0	0	0
			1813	1136	304	366	7			

- Molecule 5 is a protein called Proteasome subunit alpha type-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	231	Total	C	N	O	S	0	0	0
			1773	1114	307	348	4			
5	S	231	Total	C	N	O	S	0	0	0
			1773	1114	307	348	4			

- Molecule 6 is a protein called Probable proteasome subunit alpha type-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	243	Total	C	N	O	S	0	0	0
			1892	1203	329	356	4			
6	T	243	Total	C	N	O	S	0	0	0
			1892	1203	329	356	4			

- Molecule 7 is a protein called Proteasome subunit alpha type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	241	Total	C	N	O	S	0	0	0
			1907	1214	320	365	8			
7	U	241	Total	C	N	O	S	0	0	0
			1907	1214	320	365	8			

- Molecule 8 is a protein called Proteasome subunit beta type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	222	Total	C	N	O	S	0	0	0
			1684	1061	293	323	7			
8	V	222	Total	C	N	O	S	0	0	0
			1684	1061	293	323	7			

- Molecule 9 is a protein called Proteasome subunit beta type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			
9	W	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			

- Molecule 10 is a protein called Proteasome subunit beta type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	195	Total	C	N	O	S	0	0	0
			1561	992	264	299	6			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	X	195	Total	C	N	O	S	0	0	0
			1561	992	264	299	6			

- Molecule 11 is a protein called Proteasome subunit beta type-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			
11	Y	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			

- Molecule 12 is a protein called Proteasome subunit beta type-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			
12	Z	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			

- Molecule 13 is a protein called Proteasome subunit beta type-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			
13	a	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			

- Molecule 14 is a protein called Proteasome subunit beta type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	196	Total	C	N	O	S	0	0	0
			1515	958	250	300	7			
14	b	196	Total	C	N	O	S	0	0	0
			1515	958	250	300	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
N	128	VAL	GLY	engineered mutation	UNP P38624
b	128	VAL	GLY	engineered mutation	UNP P38624


- Molecule 15 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
15	G	1	Total Mg 1 1	0	0
15	I	1	Total Mg 1 1	0	0
15	K	1	Total Mg 1 1	0	0
15	V	1	Total Mg 1 1	0	0
15	Z	1	Total Mg 1 1	0	0

- Molecule 16 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
16	G	1	Total Cl 1 1	0	0
16	U	1	Total Cl 1 1	0	0

- Molecule 17 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
17	a	1	Total C N O S 12 6 1 4 1	0	0

- Molecule 18 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
18	A	9	Total O 9 9	0	0
18	B	14	Total O 14 14	0	0
18	C	14	Total O 14 14	0	0
18	D	9	Total O 9 9	0	0
18	E	7	Total O 7 7	0	0
18	F	14	Total O 14 14	0	0
18	G	16	Total O 16 16	0	0
18	H	19	Total O 19 19	0	0
18	I	17	Total O 17 17	0	0
18	J	10	Total O 10 10	0	0
18	K	21	Total O 21 21	0	0
18	L	15	Total O 15 15	0	0
18	M	18	Total O 18 18	0	0
18	N	9	Total O 9 9	0	0
18	O	4	Total O 4 4	0	0
18	P	14	Total O 14 14	0	0
18	Q	14	Total O 14 14	0	0
18	R	8	Total O 8 8	0	0
18	S	6	Total O 6 6	0	0
18	T	13	Total O 13 13	0	0
18	U	22	Total O 22 22	0	0
18	V	15	Total O 15 15	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
18	W	12	Total 12	O 12	0	0
18	X	13	Total 13	O 13	0	0
18	Y	11	Total 11	O 11	0	0
18	Z	14	Total 14	O 14	0	0
18	a	17	Total 17	O 17	0	0
18	b	11	Total 11	O 11	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Proteasome subunit alpha type-2




- Molecule 1: Proteasome subunit alpha type-2




- Molecule 2: Proteasome subunit alpha type-3





- Molecule 2: Proteasome subunit alpha type-3





- Molecule 3: Proteasome subunit alpha type-4





- Molecule 3: Proteasome subunit alpha type-4



- Molecule 4: Proteasome subunit alpha type-5



- Molecule 4: Proteasome subunit alpha type-5




- Molecule 5: Proteasome subunit alpha type-6



- Molecule 5: Proteasome subunit alpha type-6



- Molecule 6: Probable proteasome subunit alpha type-7



SER
ASP
SER
GLY
ILE
ASN
ALA
PRO
VAL
ALA
THR
THR
ASN
ALA
ALA
THR
THR
ASP
GLN
GLY
GLY
ASP
ILE
HIS
LEU
GLU

- Molecule 6: Probable proteasome subunit alpha type-7

Chain T:  78% 6% 16%

MET THR SER GLY T2 Q19 I34 G41 K59 R68 L77 S94 I105 P106 N123 S124 L172 E181 I195 N203 K206 L210 E211 I212 C215 N221 N244 GLY ASP ASP ASP GLU GLU ASP ASP SER ASP ASP VAL MET

SER
SER
ASP
ASP
GLY
ALA
ASN
PRO
VAL
ALA
THR
THR
ASN
ALA
ALA
THR
THR
ASP
GLN
GLY
GLY
ASP
ILE
HIS
LEU
GLU

- Molecule 7: Proteasome subunit alpha type-1

Chain G:  87% 8% 5%

MET SER GLY ALA ALA ALA SER SER G2 E13 F23 T26 L34 V43 S61 I66 S67 R68 M72 V73 V74 I78 P79 N83 L115 M125 T133 Q166 K181 V194 E215 K223 R235 Q242 ASP

- Molecule 7: Proteasome subunit alpha type-1

Chain U:  87% 8% 5%

MET SER GLY ALA ALA ALA SER SER G2 P12 E13 F23 T26 L34 V43 S61 R68 V73 I78 P79 N83 L115 M125 T133 A159 Q166 K181 V194 E215 K223 R235 Q242 ASP

- Molecule 8: Proteasome subunit beta type-2

Chain H:  89% 6% 5%

T1 V13 L34 S38 T52 V55 T56 L68 D104 P105 I113 H114 L127 M135 K153 M165 R196 Q222 ILE GLN GLU GLU VAL ASP ILE THR ALA

- Molecule 8: Proteasome subunit beta type-2

Chain V:  89% 6% 5%


T1 V13 L34 S38 T52 V55 T56 L68 D104 P105 I113 S126 L127 M135 K153 M165 R196 Q222 ILE GLN GLU GLU VAL ASP ILE THR ALA

- Molecule 9: Proteasome subunit beta type-3

Chain I:  93% 6% 1%

MET SI G9 I10 V20 N37 K38 K41 A56 T57 D58 N71 P118 A141 L171 D177 G183 D204


- Molecule 9: Proteasome subunit beta type-3

Chain W:  93% 6%




- Molecule 10: Proteasome subunit beta type-4

Chain J:  80% 15% . .



- Molecule 10: Proteasome subunit beta type-4

Chain X:  82% 13% . .



- Molecule 11: Proteasome subunit beta type-5

Chain K:  91% 7% .




- Molecule 11: Proteasome subunit beta type-5

Chain Y:  89% 8% .




- Molecule 12: Proteasome subunit beta type-6

Chain L:  92% 7% .






- Molecule 12: Proteasome subunit beta type-6




Chain Z:  93% 6% .




• Molecule 13: Proteasome subunit beta type-7




• Molecule 13: Proteasome subunit beta type-7



• Molecule 14: Proteasome subunit beta type-1



• Molecule 14: Proteasome subunit beta type-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	135.91Å 300.97Å 144.81Å 90.00° 113.54° 90.00°	Depositor
Resolution (Å)	15.00 – 2.70 15.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	98.1 (15.00-2.70) 98.1 (15.00-2.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.83 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.178 , 0.213 0.186 , 0.218	Depositor DCC
R_{free} test set	14274 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	51.1	Xtriage
Anisotropy	0.491	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 52.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	49670	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MG, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.01	0/1952	1.42	0/2642
1	O	1.01	0/1952	1.43	2/2642 (0.1%)
2	B	1.01	0/1916	1.44	1/2592 (0.0%)
2	P	1.02	0/1934	1.43	0/2618
3	C	1.02	0/1910	1.46	0/2586
3	Q	1.02	0/1910	1.46	0/2586
4	D	1.02	0/1837	1.47	2/2475 (0.1%)
4	R	1.03	0/1837	1.47	2/2475 (0.1%)
5	E	1.02	0/1800	1.43	2/2433 (0.1%)
5	S	1.03	0/1800	1.44	2/2433 (0.1%)
6	F	1.01	0/1932	1.45	4/2609 (0.2%)
6	T	1.00	0/1932	1.46	4/2609 (0.2%)
7	G	1.00	0/1945	1.42	0/2634
7	U	1.00	0/1945	1.42	0/2634
8	H	1.02	0/1715	1.39	1/2326 (0.0%)
8	V	1.02	0/1715	1.39	0/2326
9	I	1.00	0/1611	1.41	1/2174 (0.0%)
9	W	1.00	0/1611	1.41	1/2174 (0.0%)
10	J	0.98	0/1589	1.37	0/2142
10	X	1.00	0/1589	1.38	0/2142
11	K	0.99	0/1681	1.45	0/2274
11	Y	0.99	0/1681	1.44	0/2274
12	L	0.99	0/1795	1.38	0/2420
12	Z	0.99	0/1795	1.37	0/2420
13	M	0.98	0/1855	1.37	0/2514
13	a	0.99	0/1855	1.39	0/2514
14	N	0.98	0/1544	1.41	0/2092
14	b	0.99	0/1544	1.42	1/2092 (0.0%)
All	All	1.01	0/50182	1.42	23/67852 (0.0%)

There are no bond length outliers.

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	216	ARG	CB-CA-C	-6.73	102.53	113.37
9	W	183	GLY	CA-C-O	-5.66	118.54	122.22
9	I	183	GLY	CA-C-O	-5.66	118.54	122.22
6	T	77	LEU	CA-C-N	5.55	124.77	120.33
6	T	77	LEU	C-N-CA	5.55	124.77	120.33

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1915	0	1929	4	0
1	O	1915	0	1929	7	0
2	B	1887	0	1889	11	0
2	P	1904	0	1904	11	0
3	C	1881	0	1895	11	0
3	Q	1881	0	1895	13	0
4	D	1813	0	1797	7	0
4	R	1813	0	1797	7	0
5	E	1773	0	1775	12	0
5	S	1773	0	1775	12	0
6	F	1892	0	1883	7	0
6	T	1892	0	1883	7	0
7	G	1907	0	1901	11	0
7	U	1907	0	1901	13	0
8	H	1684	0	1688	4	0
8	V	1684	0	1688	5	0
9	I	1581	0	1574	9	0
9	W	1581	0	1574	10	0
10	J	1561	0	1569	27	0
10	X	1561	0	1569	25	0
11	K	1644	0	1595	14	0
11	Y	1644	0	1595	16	0
12	L	1757	0	1711	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
12	Z	1757	0	1711	10	0
13	M	1824	0	1832	27	0
13	a	1824	0	1832	26	0
14	N	1515	0	1487	30	0
14	b	1515	0	1487	24	0
15	G	1	0	0	0	0
15	I	1	0	0	0	0
15	K	1	0	0	0	0
15	V	1	0	0	0	0
15	Z	1	0	0	0	0
16	G	1	0	0	0	0
16	U	1	0	0	0	0
17	a	12	0	13	0	0
18	A	9	0	0	0	0
18	B	14	0	0	0	0
18	C	14	0	0	0	0
18	D	9	0	0	0	0
18	E	7	0	0	0	0
18	F	14	0	0	0	0
18	G	16	0	0	0	0
18	H	19	0	0	0	0
18	I	17	0	0	0	0
18	J	10	0	0	0	0
18	K	21	0	0	0	0
18	L	15	0	0	0	0
18	M	18	0	0	1	0
18	N	9	0	0	0	0
18	O	4	0	0	0	0
18	P	14	0	0	0	0
18	Q	14	0	0	0	0
18	R	8	0	0	0	0
18	S	6	0	0	0	0
18	T	13	0	0	0	0
18	U	22	0	0	0	0
18	V	15	0	0	0	0
18	W	12	0	0	0	0
18	X	13	0	0	0	0
18	Y	11	0	0	0	0
18	Z	14	0	0	0	0
18	a	17	0	0	0	0
18	b	11	0	0	0	0
All	All	49670	0	49078	326	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 326 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
13:a:119:VAL:HG23	13:a:200:ILE:HG22	1.42	1.02
13:M:119:VAL:HG23	13:M:200:ILE:HG22	1.44	0.99
3:Q:160:GLN:HA	3:Q:160:GLN:HE21	1.37	0.89
3:C:160:GLN:HE21	3:C:160:GLN:HA	1.37	0.89
10:X:3:ILE:HD12	10:X:176:PHE:CD2	2.18	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/250 (99%)	241 (97%)	7 (3%)	0	100	100
1	O	248/250 (99%)	241 (97%)	7 (3%)	0	100	100
2	B	237/258 (92%)	228 (96%)	7 (3%)	2 (1%)	16	37
2	P	242/258 (94%)	233 (96%)	8 (3%)	1 (0%)	30	54
3	C	238/254 (94%)	230 (97%)	5 (2%)	3 (1%)	9	25
3	Q	238/254 (94%)	230 (97%)	5 (2%)	3 (1%)	9	25
4	D	231/260 (89%)	228 (99%)	3 (1%)	0	100	100
4	R	231/260 (89%)	229 (99%)	2 (1%)	0	100	100
5	E	229/234 (98%)	221 (96%)	8 (4%)	0	100	100
5	S	229/234 (98%)	221 (96%)	8 (4%)	0	100	100
6	F	241/288 (84%)	234 (97%)	7 (3%)	0	100	100
6	T	241/288 (84%)	234 (97%)	7 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
7	G	239/252 (95%)	232 (97%)	7 (3%)	0	100	100
7	U	239/252 (95%)	234 (98%)	5 (2%)	0	100	100
8	H	220/232 (95%)	214 (97%)	6 (3%)	0	100	100
8	V	220/232 (95%)	214 (97%)	6 (3%)	0	100	100
9	I	202/205 (98%)	196 (97%)	6 (3%)	0	100	100
9	W	202/205 (98%)	196 (97%)	6 (3%)	0	100	100
10	J	193/198 (98%)	190 (98%)	3 (2%)	0	100	100
10	X	193/198 (98%)	190 (98%)	3 (2%)	0	100	100
11	K	210/212 (99%)	208 (99%)	2 (1%)	0	100	100
11	Y	210/212 (99%)	208 (99%)	2 (1%)	0	100	100
12	L	220/222 (99%)	214 (97%)	6 (3%)	0	100	100
12	Z	220/222 (99%)	214 (97%)	6 (3%)	0	100	100
13	M	231/246 (94%)	224 (97%)	7 (3%)	0	100	100
13	a	231/246 (94%)	224 (97%)	7 (3%)	0	100	100
14	N	194/196 (99%)	187 (96%)	7 (4%)	0	100	100
14	b	194/196 (99%)	187 (96%)	6 (3%)	1 (0%)	24	48
All	All	6271/6614 (95%)	6102 (97%)	159 (2%)	10 (0%)	43	68

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	51	VAL
3	C	202	GLN
2	P	51	VAL
3	Q	202	GLN
3	C	205	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	203 (97%)	6 (3%)	37	67
1	O	209/209 (100%)	203 (97%)	6 (3%)	37	67
2	B	202/216 (94%)	193 (96%)	9 (4%)	24	52
2	P	203/216 (94%)	196 (97%)	7 (3%)	32	62
3	C	212/226 (94%)	206 (97%)	6 (3%)	38	68
3	Q	212/226 (94%)	206 (97%)	6 (3%)	38	68
4	D	194/215 (90%)	185 (95%)	9 (5%)	24	51
4	R	194/215 (90%)	185 (95%)	9 (5%)	24	51
5	E	190/193 (98%)	185 (97%)	5 (3%)	40	70
5	S	190/193 (98%)	185 (97%)	5 (3%)	40	70
6	F	201/239 (84%)	193 (96%)	8 (4%)	28	56
6	T	201/239 (84%)	193 (96%)	8 (4%)	28	56
7	G	206/210 (98%)	197 (96%)	9 (4%)	25	53
7	U	206/210 (98%)	197 (96%)	9 (4%)	25	53
8	H	181/190 (95%)	171 (94%)	10 (6%)	19	45
8	V	181/190 (95%)	171 (94%)	10 (6%)	19	45
9	I	172/173 (99%)	170 (99%)	2 (1%)	63	84
9	W	172/173 (99%)	170 (99%)	2 (1%)	63	84
10	J	173/175 (99%)	163 (94%)	10 (6%)	18	42
10	X	173/175 (99%)	163 (94%)	10 (6%)	18	42
11	K	169/169 (100%)	162 (96%)	7 (4%)	27	56
11	Y	169/169 (100%)	162 (96%)	7 (4%)	27	56
12	L	185/185 (100%)	179 (97%)	6 (3%)	34	64
12	Z	185/185 (100%)	179 (97%)	6 (3%)	34	64
13	M	199/208 (96%)	185 (93%)	14 (7%)	14	34
13	a	199/208 (96%)	185 (93%)	14 (7%)	14	34
14	N	163/163 (100%)	145 (89%)	18 (11%)	6	15
14	b	163/163 (100%)	144 (88%)	19 (12%)	5	13
All	All	5313/5542 (96%)	5076 (96%)	237 (4%)	24	52

5 of 237 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
14	N	177	VAL
14	b	14	LEU
4	R	197	LYS
14	b	9	LYS
14	b	192	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 178 such sidechains are listed below:

Mol	Chain	Res	Type
5	S	184	ASN
10	X	55	GLN
6	T	123	ASN
8	V	30	ASN
11	Y	62	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 7 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
17	MES	a	301	-	12,12,12	0.69	0	15,16,16	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
17	MES	a	301	-	-	3/6/14/14	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
17	a	301	MES	C7-C8-S-O2S
17	a	301	MES	C7-C8-S-O3S
17	a	301	MES	C7-C8-S-O1S

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/250 (100%)	-0.41	2 (0%) 82 81	42, 54, 88, 139	0
1	O	250/250 (100%)	-0.35	2 (0%) 82 81	46, 62, 103, 144	0
2	B	241/258 (93%)	-0.28	2 (0%) 82 81	44, 61, 98, 152	0
2	P	244/258 (94%)	-0.17	5 (2%) 65 62	48, 65, 108, 148	0
3	C	240/254 (94%)	-0.28	3 (1%) 75 73	45, 66, 121, 151	0
3	Q	240/254 (94%)	-0.13	6 (2%) 58 55	50, 73, 139, 159	0
4	D	235/260 (90%)	-0.30	1 (0%) 88 87	46, 67, 95, 132	0
4	R	235/260 (90%)	-0.26	2 (0%) 81 80	46, 67, 97, 133	0
5	E	231/234 (98%)	-0.34	0 100 100	49, 66, 94, 127	0
5	S	231/234 (98%)	-0.32	1 (0%) 88 87	47, 69, 97, 125	0
6	F	243/288 (84%)	-0.40	1 (0%) 88 87	42, 60, 100, 124	0
6	T	243/288 (84%)	-0.37	2 (0%) 82 81	43, 62, 102, 124	0
7	G	241/252 (95%)	-0.44	0 100 100	42, 55, 82, 114	0
7	U	241/252 (95%)	-0.39	1 (0%) 88 87	44, 58, 92, 110	0
8	H	222/232 (95%)	-0.42	1 (0%) 87 86	41, 53, 76, 104	0
8	V	222/232 (95%)	-0.40	0 100 100	46, 56, 83, 124	0
9	I	204/205 (99%)	-0.55	0 100 100	40, 51, 73, 106	0
9	W	204/205 (99%)	-0.58	0 100 100	43, 54, 79, 106	0
10	J	195/198 (98%)	-0.49	2 (1%) 79 78	42, 56, 76, 110	0
10	X	195/198 (98%)	-0.51	2 (1%) 79 78	44, 58, 78, 122	0
11	K	212/212 (100%)	-0.51	0 100 100	42, 55, 72, 92	0
11	Y	212/212 (100%)	-0.47	0 100 100	43, 54, 73, 89	0
12	L	222/222 (100%)	-0.42	1 (0%) 87 86	43, 56, 89, 115	0
12	Z	222/222 (100%)	-0.37	1 (0%) 87 86	41, 53, 83, 127	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
13	M	233/246 (94%)	-0.43	3 (1%) 75 73	41, 56, 75, 92	0
13	a	233/246 (94%)	-0.51	3 (1%) 75 73	38, 54, 74, 96	0
14	N	196/196 (100%)	-0.26	6 (3%) 51 48	41, 55, 100, 124	0
14	b	196/196 (100%)	-0.33	2 (1%) 79 78	42, 56, 100, 117	0
All	All	6333/6614 (95%)	-0.38	49 (0%) 82 81	38, 59, 97, 159	0

The worst 5 of 49 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	Q	205	ALA	5.2
14	N	137	TYR	5.1
3	Q	50	LEU	4.7
2	P	219	ALA	3.8
10	J	1	MET	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
15	MG	I	301	1/1	0.73	0.23	109,109,109,109	0
15	MG	Z	301	1/1	0.83	0.26	137,137,137,137	0
16	CL	U	301	1/1	0.87	0.08	80,80,80,80	0
17	MES	a	301	12/12	0.88	0.14	97,108,110,111	0
15	MG	K	301	1/1	0.94	0.15	92,92,92,92	0
16	CL	G	302	1/1	0.96	0.05	57,57,57,57	0
15	MG	V	301	1/1	0.96	0.09	91,91,91,91	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
15	MG	G	301	1/1	0.96	0.08	75,75,75,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.