



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:10 AM UTC

PDB ID : 5QD1 / pdb_00005qd1
Title : Crystal structure of BACE complex with BMC011
Authors : Rondeau, J.M.; Shao, C.; Yang, H.; Burley, S.K.
Deposited on : 2017-12-01
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	402	 4% 89% 5% 6%
1	B	402	 3% 88% 6% 6%
1	C	402	 3% 87% 7% 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-secretase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	377	Total	C	N	O	S	0	0	0
			2966	1898	493	561	14			
1	B	377	Total	C	N	O	S	0	0	0
			2966	1898	493	561	14			
1	C	381	Total	C	N	O	S	0	0	0
			2993	1917	497	565	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-15	GLY	-	expression tag	UNP P56817
A	-14	PRO	-	expression tag	UNP P56817
B	-15	GLY	-	expression tag	UNP P56817
B	-14	PRO	-	expression tag	UNP P56817
C	-15	GLY	-	expression tag	UNP P56817
C	-14	PRO	-	expression tag	UNP P56817

- Molecule 2 is (10S,12S)-12-[(1R)-1-hydroxy-2-({[3-(propan-2-yl)phenyl]methyl}amino)ethyl]-17-(methoxymethyl)-10-methyl-7-oxa-2,13-diazabicyclo[13.3.1]nonadeca-1(19),15,17-trien-1 4-one (CCD ID: E61) (formula: C₃₁H₄₇N₃O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			38	31	3	4		
2	B	1	Total	C	N	O	0	0
			38	31	3	4		
2	C	1	Total	C	N	O	0	0
			38	31	3	4		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	127	Total	O	0	0
			127	127		
3	B	121	Total	O	0	0
			121	121		
3	C	133	Total	O	0	0
			133	133		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	82.85Å 104.22Å 100.83Å 90.00° 104.37° 90.00°	Depositor
Resolution (Å)	80.26 – 2.40 80.26 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (80.26-2.40) 100.0 (80.26-2.40)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.84 (at 2.40Å)	Xtriage
Refinement program	PHENIX 1.12_2829, CNX	Depositor
R, R_{free}	0.163 , 0.197 0.171 , 0.203	Depositor DCC
R_{free} test set	6516 reflections (10.02%)	wwPDB-VP
Wilson B-factor (Å ²)	40.7	Xtriage
Anisotropy	0.043	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 58.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9420	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: E61

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.37	0/3041	0.55	0/4133
1	B	0.36	0/3041	0.54	0/4133
1	C	0.35	0/3070	0.54	0/4173
All	All	0.36	0/9152	0.55	0/12439

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2966	0	2878	12	0
1	B	2966	0	2878	14	0
1	C	2993	0	2902	15	0
2	A	38	0	0	0	0
2	B	38	0	0	0	0
2	C	38	0	0	0	0
3	A	127	0	0	1	0
3	B	121	0	0	3	1
3	C	133	0	0	1	1
All	All	9420	0	8658	40	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:207:GLU:OE1	3:C:501:HOH:O	2.14	0.64
1:A:232:THR:OG1	3:A:501:HOH:O	2.15	0.63
1:B:357:SER:O	1:B:360:HIS:HB3	2.02	0.59
1:B:267:LEU:HD12	1:B:319:CYS:HB3	1.86	0.58
1:A:2:MET:HG2	1:A:90:GLY:HA2	1.90	0.54
1:A:270:TRP:O	1:A:317:ASP:HB3	2.09	0.52
1:C:2:MET:HG2	1:C:90:GLY:HA2	1.91	0.52
1:B:306:LEU:O	3:B:501:HOH:O	2.19	0.51
1:C:267:LEU:HD12	1:C:319:CYS:HB3	1.94	0.50
1:A:254:THR:HG21	1:C:254:THR:HG21	1.93	0.50
1:C:270:TRP:O	1:C:317:ASP:HB3	2.11	0.50
1:A:267:LEU:H	1:A:267:LEU:HD23	1.78	0.49
1:B:270:TRP:O	1:B:317:ASP:HB3	2.12	0.48
1:B:222:TYR:HA	1:B:223:ASP:HA	1.61	0.48
1:B:2:MET:HG2	1:B:90:GLY:HA2	1.95	0.47
1:C:202:ILE:CD1	1:C:379:MET:HG3	2.44	0.47
1:A:267:LEU:HD12	1:A:319:CYS:HB3	1.95	0.47
1:B:64:ARG:NH1	3:B:507:HOH:O	2.47	0.46
1:C:292:THR:HG21	1:C:378:ASP:HB3	1.97	0.46
1:B:174:MET:HE3	1:B:176:ILE:HD11	1.98	0.46
1:B:205:ARG:NH2	1:B:212:ASP:HB2	2.32	0.45
1:B:110:ILE:HB	1:B:113:SER:HB3	1.99	0.44
1:C:304:GLN:HG3	1:C:361:VAL:HG21	1.99	0.44
1:A:222:TYR:HA	1:A:223:ASP:HA	1.64	0.44
1:C:13:GLY:HA3	1:C:30:LEU:HD11	2.00	0.44
1:B:55:GLN:NE2	3:B:502:HOH:O	2.50	0.44
1:C:9:LYS:HD2	1:C:168:ALA:HB3	2.00	0.43
1:A:20:VAL:HG12	1:A:85:VAL:HG22	2.01	0.43
1:C:211:GLN:HE21	1:C:211:GLN:HB2	1.59	0.43
1:C:212:ASP:O	1:C:214:LYS:HG3	2.19	0.42
1:C:222:TYR:HA	1:C:223:ASP:HA	1.63	0.42
1:B:125:GLU:O	1:B:125:GLU:HG3	2.18	0.42
1:C:197:TRP:CG	1:C:198:TYR:H	2.37	0.42
1:C:363:ASP:HB3	1:C:366:ARG:O	2.20	0.42
1:B:304:GLN:HG3	1:B:361:VAL:HG11	2.02	0.41
1:A:63:LEU:HD23	1:A:63:LEU:HA	1.94	0.41
1:A:154:LEU:O	1:A:339:GLU:HA	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:MET:HE3	1:A:176:ILE:HD11	2.02	0.41
1:B:95:VAL:HG11	1:B:140:LEU:HA	2.03	0.40
1:A:197:TRP:CG	1:A:198:TYR:H	2.39	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:607:HOH:O	3:C:619:HOH:O[2_655]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	373/402 (93%)	364 (98%)	9 (2%)	0	100	100
1	B	373/402 (93%)	362 (97%)	10 (3%)	1 (0%)	36	50
1	C	377/402 (94%)	366 (97%)	11 (3%)	0	100	100
All	All	1123/1206 (93%)	1092 (97%)	30 (3%)	1 (0%)	48	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	70	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	322/342 (94%)	321 (100%)	1 (0%)	86	93
1	B	322/342 (94%)	320 (99%)	2 (1%)	78	89
1	C	324/342 (95%)	318 (98%)	6 (2%)	50	71
All	All	968/1026 (94%)	959 (99%)	9 (1%)	70	85

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	211	GLN
1	B	125	GLU
1	B	267	LEU
1	C	1	GLU
1	C	9	LYS
1	C	208	ILE
1	C	211	GLN
1	C	267	LEU
1	C	362	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	GLN
1	A	89	HIS
1	B	89	HIS
1	B	181	HIS
1	B	385	ASN
1	C	53	GLN
1	C	89	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	E61	C	401	-	40,40,40	0.44	0	50,52,52	0.68	1 (2%)
2	E61	B	401	-	40,40,40	0.55	0	50,52,52	0.75	1 (2%)
2	E61	A	401	-	40,40,40	0.45	0	50,52,52	0.77	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	E61	C	401	-	-	2/40/40/40	0/2/3/3
2	E61	B	401	-	-	2/40/40/40	0/2/3/3
2	E61	A	401	-	-	1/40/40/40	0/2/3/3

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	401	E61	C54-C58-N61	-4.54	105.99	111.95
2	A	401	E61	C54-C58-N61	-3.92	106.80	111.95
2	C	401	E61	C54-C58-N61	-3.75	107.03	111.95
2	A	401	E61	C8-C5-C3	-2.04	111.53	115.56

There are no chirality outliers.


All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	E61	C37-C40-O43-C44
2	C	401	E61	C37-C40-O43-C44
2	B	401	E61	C37-C40-O43-C44
2	C	401	E61	N1-C3-C5-C8
2	B	401	E61	C10-C13-O16-C17


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



Ligand E61 B 401





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	377/402 (93%)	-0.30	16 (4%) 40 36	26, 41, 78, 159	0
1	B	377/402 (93%)	-0.24	14 (3%) 45 41	23, 43, 93, 180	0
1	C	381/402 (94%)	-0.27	14 (3%) 45 41	28, 43, 84, 156	0
All	All	1135/1206 (94%)	-0.27	44 (3%) 43 39	23, 42, 85, 180	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	312	VAL	6.5
1	C	312	VAL	5.2
1	C	362	HIS	4.4
1	B	314	THR	4.2
1	B	312	VAL	4.1
1	C	168	ALA	4.1
1	C	314	THR	3.7
1	B	313	ALA	3.7
1	A	169	SER	3.6
1	C	159	PHE	3.6
1	A	157	ALA	3.3
1	C	361	VAL	3.2
1	A	313	ALA	3.1
1	A	362	HIS	3.1
1	C	160	PRO	3.0
1	A	254	THR	2.9
1	C	315	SER	2.9
1	A	311	ASP	2.8
1	A	365	PHE	2.8
1	B	315	SER	2.8
1	C	313	ALA	2.7
1	C	-2	SER	2.6
1	B	-2	SER	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	380	GLU	2.5
1	C	311	ASP	2.4
1	A	361	VAL	2.4
1	C	254	THR	2.3
1	A	256	LYS	2.3
1	B	258	PRO	2.3
1	A	314	THR	2.3
1	B	254	THR	2.3
1	B	316	GLN	2.3
1	A	259	ASP	2.2
1	C	316	GLN	2.2
1	B	310	GLU	2.2
1	A	315	SER	2.2
1	B	273	GLY	2.1
1	C	158	GLY	2.1
1	B	267	LEU	2.1
1	B	256	LYS	2.1
1	A	316	GLN	2.1
1	B	311	ASP	2.1
1	A	258	PRO	2.1
1	B	259	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	E61	A	401	38/38	0.95	0.10	29,33,47,66	4
2	E61	B	401	38/38	0.96	0.08	25,34,48,55	3

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	E61	C	401	38/38	0.96	0.08	29,35,45,60	3

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around E61 A 401:


2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around E61 B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.