



Full wwPDB NMR Structure Validation Report ⓘ

Mar 5, 2026 – 07:50 PM UTC

PDB ID : 7QDE / pdb_00007qde
BMRB ID : 34692
Title : NMR structure of Npl3 RRM12 bound to the AUCCAGUGGAA RNA
Authors : Allain, F.H.-T.; Clery, A.; Moursy, A.
Deposited on : 2021-11-26

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 46%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823
RNA backbone	8273	777

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	11	9% 45% 45%
2	B	169	73% 17% • 8%

2 Ensemble composition and analysis

This entry contains 10 models. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	B:120-B:275 (156)	0.97	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 1 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 8, 10
Single-model clusters	7; 9

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 3038 atoms, of which 1445 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called RNA (5'-R(*AP*UP*CP*CP*AP*GP*UP*GP*GP*AP*A)-3').

Mol	Chain	Residues	Atoms						Trace
1	A	11	Total	C	H	N	O	P	0
			355	106	121	45	73	10	

- Molecule 2 is a protein called Nucleolar protein 3.


Mol	Chain	Residues	Atoms						Trace
2	B	169	Total	C	H	N	O	S	0
			2683	856	1324	234	265	4	

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: RNA (5'-R(*AP*UP*CP*CP*AP*GP*UP*GP*GP*AP*A)-3')

Chain A: 



- Molecule 2: Nucleolar protein 3

Chain B: 




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: RNA (5'-R(*AP*UP*CP*CP*AP*GP*UP*GP*GP*AP*A)-3')

Chain A: 



- Molecule 2: Nucleolar protein 3

Chain B: 



4.2.2 Score per residue for model 2

- Molecule 1: RNA (5'-R(*AP*UP*CP*CP*AP*GP*UP*GP*GP*AP*A)-3')




- Molecule 2: Nucleolar protein 3



4.2.3 Score per residue for model 3

- Molecule 1: RNA (5'-R(*AP*UP*CP*CP*AP*GP*UP*GP*GP*AP*A)-3')



- Molecule 2: Nucleolar protein 3



4.2.4 Score per residue for model 4 (medoid)


- Molecule 1: RNA (5'-R(*AP*UP*CP*CP*AP*GP*UP*GP*GP*AP*A)-3')

Chain A:  9% 36% 55%




- Molecule 2: Nucleolar protein 3

Chain B:  73% 15% 8%



4.2.5 Score per residue for model 5


- Molecule 1: RNA (5'-R(*AP*UP*CP*CP*AP*GP*UP*GP*GP*AP*A)-3')

Chain A:  18% 45% 36%




- Molecule 2: Nucleolar protein 3

Chain B:  71% 17% 8%



4.2.6 Score per residue for model 6


- Molecule 1: RNA (5'-R(*AP*UP*CP*CP*AP*GP*UP*GP*GP*AP*A)-3')

Chain A:  9% 55% 36%



- Molecule 2: Nucleolar protein 3

Chain B:  73% 15% 8%



4.2.7 Score per residue for model 7


- Molecule 1: RNA (5'-R(*AP*UP*CP*CP*AP*GP*UP*GP*GP*AP*A)-3')

Chain A: 9% 45% 45%



- Molecule 2: Nucleolar protein 3

Chain B: 75% 15% 8%



4.2.8 Score per residue for model 8


- Molecule 1: RNA (5'-R(*AP*UP*CP*CP*AP*GP*UP*GP*GP*AP*A)-3')

Chain A: 9% 64% 27%




- Molecule 2: Nucleolar protein 3


Chain B: 74% 15% 8%



4.2.9 Score per residue for model 9


- Molecule 1: RNA (5'-R(*AP*UP*CP*CP*AP*GP*UP*GP*GP*AP*A)-3')

Chain A:  18% 36% 36% 9%




- Molecule 2: Nucleolar protein 3

Chain B:  70% 20% 8%



4.2.10 Score per residue for model 10


- Molecule 1: RNA (5'-R(*AP*UP*CP*CP*AP*GP*UP*GP*GP*AP*A)-3')

Chain A:  64% 36%



- Molecule 2: Nucleolar protein 3

Chain B:  74% 14% 5% 8%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 50 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	refinement	
CYANA	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1122
Number of shifts mapped to atoms	1122
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	46%

6 Model quality

6.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.73±0.01	0±0/262 (0.0± 0.0%)	1.04±0.03	0±0/407 (0.0± 0.0%)
2	B	0.51±0.00	0±0/1269 (0.0± 0.0%)	0.91±0.01	1±1/1717 (0.0± 0.1%)
All	All	0.55	0/15310 (0.0%)	0.94	8/21240 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.5±0.7
All	All	0	15

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	275	ASN	CA-C-N	5.45	123.58	119.66	2	4
2	B	275	ASN	C-N-CA	5.45	123.58	119.66	2	4

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	106	C	Sidechain	10
1	A	111	G	Sidechain	4
1	A	104	U	Sidechain	1

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	234	121	121	3±1
2	B	1244	1209	1209	17±3
All	All	14780	13300	13300	178

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:122:LEU:HD13	2:B:122:LEU:H	0.93	1.23	9	10
2:B:122:LEU:H	2:B:122:LEU:CD1	0.70	2.00	7	6
2:B:225:LEU:N	2:B:225:LEU:HD22	0.66	2.06	8	1
2:B:127:LEU:HD12	2:B:175:ILE:HD12	0.62	1.70	2	3
2:B:225:LEU:HD22	2:B:225:LEU:H	0.59	1.58	8	1
2:B:225:LEU:H	2:B:225:LEU:CD2	0.56	2.13	8	1
2:B:195:LEU:HD22	2:B:195:LEU:N	0.56	2.16	4	7
2:B:219:LEU:HD11	2:B:261:ILE:HD11	0.56	1.78	5	9
2:B:195:LEU:H	2:B:195:LEU:HD13	0.54	1.62	3	1
2:B:219:LEU:HD13	2:B:263:PHE:CD1	0.54	2.38	9	9
1:A:106:C:O2'	1:A:107:A:C8	0.54	2.57	1	6
1:A:108:G:C4	1:A:108:G:H5''	0.53	2.38	7	1
2:B:225:LEU:N	2:B:225:LEU:CD2	0.53	2.72	8	1
2:B:127:LEU:HD12	2:B:175:ILE:CD1	0.53	2.33	2	3
2:B:157:LEU:N	2:B:157:LEU:HD23	0.53	2.19	2	4
2:B:122:LEU:N	2:B:122:LEU:HD13	0.52	2.19	3	1
2:B:126:ARG:C	2:B:127:LEU:HD13	0.52	2.29	2	3
2:B:122:LEU:HD11	2:B:192:TYR:HB3	0.52	1.80	7	2
2:B:195:LEU:HD22	2:B:195:LEU:H	0.51	1.64	4	4
2:B:136:VAL:HG22	2:B:136:VAL:O	0.50	2.05	2	1
1:A:105:C:C5	2:B:128:PHE:CE1	0.50	3.00	8	8
2:B:199:ARG:HD2	2:B:199:ARG:C	0.50	2.32	5	2
2:B:138:GLU:H	2:B:156:ILE:HD13	0.49	1.67	9	10
2:B:195:LEU:H	2:B:195:LEU:CD2	0.48	2.20	1	3
2:B:255:LEU:C	2:B:255:LEU:HD12	0.48	2.32	4	4
2:B:272:ARG:HD3	2:B:272:ARG:H	0.48	1.69	2	3

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:112:A:C2	2:B:212:SER:HB3	0.48	2.44	3	7
2:B:195:LEU:CD2	2:B:195:LEU:H	0.48	2.22	5	1
2:B:122:LEU:HD13	2:B:122:LEU:N	0.48	2.10	7	1
2:B:157:LEU:HD23	2:B:157:LEU:N	0.47	2.24	3	5
2:B:127:LEU:HD22	2:B:175:ILE:HD12	0.47	1.86	1	7
1:A:108:G:H5''	1:A:108:G:C4	0.47	2.45	1	2
1:A:108:G:C8	2:B:229:PHE:CD1	0.47	3.03	8	1
1:A:108:G:N7	2:B:229:PHE:CG	0.46	2.83	4	1
2:B:122:LEU:HD12	2:B:193:SER:O	0.46	2.10	7	2
2:B:127:LEU:HD22	2:B:175:ILE:CD1	0.46	2.39	8	6
1:A:108:G:C5	2:B:229:PHE:CD2	0.46	3.03	9	1
2:B:130:ARG:H	2:B:131:PRO:CD	0.45	2.24	5	7
2:B:132:PHE:CE1	2:B:183:PHE:CZ	0.45	3.05	2	2
1:A:104:U:C6	2:B:130:ARG:HD2	0.45	2.46	1	1
2:B:132:PHE:CD1	2:B:136:VAL:HG21	0.45	2.46	2	1
2:B:132:PHE:CE1	2:B:136:VAL:HG21	0.45	2.47	2	1
1:A:105:C:O4'	2:B:128:PHE:CE2	0.44	2.71	2	1
2:B:255:LEU:HD12	2:B:255:LEU:C	0.44	2.37	3	6
2:B:127:LEU:HD11	2:B:171:ALA:HB1	0.44	1.87	4	1
2:B:251:LEU:HD11	2:B:272:ARG:HB3	0.43	1.89	1	2
2:B:202:ILE:CG2	2:B:251:LEU:HD13	0.43	2.43	10	1
2:B:195:LEU:N	2:B:195:LEU:CD2	0.42	2.81	1	4
2:B:183:PHE:CD1	2:B:188:LEU:HD11	0.42	2.49	10	3
1:A:110:G:H2'	1:A:111:G:H5''	0.42	1.91	5	1
2:B:202:ILE:HG22	2:B:251:LEU:HD13	0.42	1.91	6	1
2:B:223:ASN:HD22	2:B:258:LEU:CD2	0.42	2.28	1	1
2:B:130:ARG:H	2:B:131:PRO:HD2	0.41	1.75	1	1
1:A:105:C:C5'	2:B:160:PHE:CE1	0.41	3.03	1	1
1:A:111:G:C2	2:B:214:GLN:HB3	0.41	2.51	9	1
1:A:108:G:C4	2:B:229:PHE:CE2	0.41	3.09	5	1
2:B:127:LEU:HD21	2:B:171:ALA:HB1	0.41	1.91	1	2
1:A:110:G:C8	2:B:213:TRP:CH2	0.41	3.08	9	1

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	156/169 (92%)	129±1 (83±1%)	20±1 (13±1%)	7±1 (4±1%)	3	27
All	All	1560/1690 (92%)	1290 (83%)	202 (13%)	68 (4%)	3	27

All 13 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	130	ARG	10
2	B	133	PRO	10
2	B	134	LEU	10
2	B	151	MET	10
2	B	224	SER	10
2	B	237	PHE	6
2	B	262	GLU	4
2	B	206	ASN	3
2	B	123	SER	1
2	B	136	VAL	1
2	B	122	LEU	1
2	B	236	ASP	1
2	B	233	ASN	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	137/150 (91%)	121±1 (88±1%)	16±1 (12±1%)	7	50
All	All	1370/1500 (91%)	1210 (88%)	160 (12%)	7	50

All 27 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	122	LEU	10
2	B	124	ASN	10
2	B	175	ILE	10
2	B	181	LYS	10
2	B	183	PHE	10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
2	B	202	ILE	10
2	B	227	THR	10
2	B	261	ILE	10
2	B	268	ILE	10
2	B	136	VAL	9
2	B	154	VAL	9
2	B	157	LEU	9
2	B	130	ARG	8
2	B	173	LYS	7
2	B	234	THR	6
2	B	251	LEU	5
2	B	195	LEU	3
2	B	127	LEU	3
2	B	272	ARG	2
2	B	275	ASN	2
2	B	126	ARG	1
2	B	228	THR	1
2	B	214	GLN	1
2	B	203	THR	1
2	B	225	LEU	1
2	B	235	ARG	1
2	B	134	LEU	1

6.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	11/11 (100%)	8±0 (73±0%)	2±1 (18±8%)	0.10±0.00
All	All	103/110 (94%)	80 (78%)	20 (19%)	0.10

The overall RNA backbone suiteness is 0.10.

All unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	104	U	10
1	A	105	C	10
1	A	108	G	10
1	A	109	U	10
1	A	110	G	10
1	A	111	G	10
1	A	112	A	10
1	A	113	A	10

All unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	108	G	6
1	A	109	U	4
1	A	103	A	3
1	A	107	A	3
1	A	110	G	3
1	A	104	U	1

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 46% for the well-defined parts and 44% for the entire structure.

7.1 Chemical shift list 1

File name: `working_cs.cif`

Chemical shift list name: *starch_output*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1122
Number of shifts mapped to atoms	1122
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	139	-0.20 ± 0.12	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	142	-0.15 ± 0.14	None needed (< 0.5 ppm)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	146	0.22 ± 0.31	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 46%, i.e. 1077 atoms were assigned a chemical shift out of a possible 2340. 0 out of 26 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	551/773 (71%)	273/313 (87%)	135/312 (43%)	143/148 (97%)
Sidechain	457/1202 (38%)	256/774 (33%)	201/380 (53%)	0/48 (0%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	11/157 (7%)	5/78 (6%)	6/77 (8%)	0/2 (0%)
Sugar	41/121 (34%)	41/66 (62%)	0/55 (0%)	0/0 (—%)
Base	17/87 (20%)	17/54 (31%)	0/19 (0%)	0/14 (0%)
Overall	1077/2340 (46%)	592/1285 (46%)	342/843 (41%)	143/212 (67%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 44%, i.e. 1120 atoms were assigned a chemical shift out of a possible 2543. 0 out of 26 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	565/832 (68%)	280/336 (83%)	139/338 (41%)	146/158 (92%)
Sidechain	486/1332 (36%)	275/857 (32%)	211/417 (51%)	0/58 (0%)
Aromatic	11/171 (6%)	5/86 (6%)	6/81 (7%)	0/4 (0%)
Sugar	41/121 (34%)	41/66 (62%)	0/55 (0%)	0/0 (—%)
Base	17/87 (20%)	17/54 (31%)	0/19 (0%)	0/14 (0%)
Overall	1120/2543 (44%)	618/1399 (44%)	356/910 (39%)	146/234 (62%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain B:

