



Full wwPDB NMR Structure Validation Report ⓘ

Mar 8, 2026 – 12:07 PM UTC

PDB ID : 7QDW / pdb_00007qdw
BMRB ID : 34693
Title : Solution structure of the complex between plasmodial ZNHIT3 and NUFIP1 proteins
Authors : Chagot, M.E.; Quinternet, M.
Deposited on : 2021-11-30

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 94%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	72	
2	B	25	

2 Ensemble composition and analysis

This entry contains 20 models. Model 14 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *structure with the lowest restraint energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:269-A:330, B:821-B:837 (79)	0.34	14

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 6 single-model clusters were found.

Cluster number	Models
1	1, 5, 10, 16
2	9, 14, 19
3	2, 3, 6
4	4, 18
5	8, 13
Single-model clusters	7; 11; 12; 15; 17; 20

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1650 atoms, of which 828 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Zinc finger protein, putative.

Mol	Chain	Residues	Atoms						Trace
1	A	72	Total	C	H	N	O	S	0
			1209	384	607	99	116	3	

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	261	GLY	-	expression tag	UNP Q8I2Y4
A	262	PRO	-	expression tag	UNP Q8I2Y4
A	263	HIS	-	expression tag	UNP Q8I2Y4
A	264	MET	-	expression tag	UNP Q8I2Y4

- Molecule 2 is a protein called NUFIP1 domain-containing protein.


Mol	Chain	Residues	Atoms					Trace
2	B	25	Total	C	H	N	O	0
			441	145	221	31	44	

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Zinc finger protein, putative



- Molecule 2: NUFIP1 domain-containing protein




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: Zinc finger protein, putative



- Molecule 2: NUFIP1 domain-containing protein



4.2.2 Score per residue for model 2

- Molecule 1: Zinc finger protein, putative

Chain A:  81% 6% 14%



- Molecule 2: NUFIP1 domain-containing protein


Chain B:  64% . 32%



4.2.3 Score per residue for model 3


- Molecule 1: Zinc finger protein, putative

Chain A:  81% 6% 14%



- Molecule 2: NUFIP1 domain-containing protein

Chain B:  68% 32%



4.2.4 Score per residue for model 4

- Molecule 1: Zinc finger protein, putative

Chain A:  79% 7% 14%



- Molecule 2: NUFIP1 domain-containing protein


Chain B:  68% 32%



4.2.5 Score per residue for model 5

- Molecule 1: Zinc finger protein, putative

Chain A:  85% 14%



- Molecule 2: NUFIP1 domain-containing protein


Chain B:  68% 32%



4.2.6 Score per residue for model 6

- Molecule 1: Zinc finger protein, putative

Chain A:  82% 14%



- Molecule 2: NUFIP1 domain-containing protein


Chain B:  64% 32%



4.2.7 Score per residue for model 7

- Molecule 1: Zinc finger protein, putative

Chain A:  85% 14%



- Molecule 2: NUFIP1 domain-containing protein


Chain B:  68% 32%



4.2.8 Score per residue for model 8

- Molecule 1: Zinc finger protein, putative

Chain A:  82% 14%



- Molecule 2: NUFIP1 domain-containing protein

Chain B:  68% 32%



4.2.9 Score per residue for model 9

- Molecule 1: Zinc finger protein, putative

Chain A:  83% 14%



- Molecule 2: NUFIP1 domain-containing protein


Chain B:  68% 32%



4.2.10 Score per residue for model 10


- Molecule 1: Zinc finger protein, putative

Chain A:  85% 14%



- Molecule 2: NUFIP1 domain-containing protein


Chain B:  68% 32%



4.2.11 Score per residue for model 11

- Molecule 1: Zinc finger protein, putative

Chain A:  81% 6% 14%



- Molecule 2: NUFIP1 domain-containing protein


Chain B:  68% 32%



4.2.12 Score per residue for model 12

- Molecule 1: Zinc finger protein, putative

Chain A:  82% 14%



- Molecule 2: NUFIP1 domain-containing protein

Chain B:  64% 32%



4.2.13 Score per residue for model 13

- Molecule 1: Zinc finger protein, putative

Chain A:  79% 7% 14%



- Molecule 2: NUFIP1 domain-containing protein

Chain B:  64% 32%



4.2.14 Score per residue for model 14 (medoid)


- Molecule 1: Zinc finger protein, putative

Chain A: 



- Molecule 2: NUFIP1 domain-containing protein

Chain B: 



4.2.15 Score per residue for model 15

- Molecule 1: Zinc finger protein, putative

Chain A: 



- Molecule 2: NUFIP1 domain-containing protein


Chain B: 



4.2.16 Score per residue for model 16

- Molecule 1: Zinc finger protein, putative

Chain A: 



- Molecule 2: NUFIP1 domain-containing protein


Chain B: 



4.2.17 Score per residue for model 17

- Molecule 1: Zinc finger protein, putative

Chain A:  85% 14%



- Molecule 2: NUFIP1 domain-containing protein


Chain B:  68% 32%



4.2.18 Score per residue for model 18

- Molecule 1: Zinc finger protein, putative

Chain A:  81% 6% 14%



- Molecule 2: NUFIP1 domain-containing protein


Chain B:  64% 32%



4.2.19 Score per residue for model 19

- Molecule 1: Zinc finger protein, putative

Chain A:  82% 14%



- Molecule 2: NUFIP1 domain-containing protein


Chain B:  68% 32%



4.2.20 Score per residue for model 20

- Molecule 1: Zinc finger protein, putative

Chain A:  85% 14%



- Molecule 2: NUFIP1 domain-containing protein

Chain B:  68% 32%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest restraint energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
TALOS-N	structure calculation	
CYANA	structure calculation	3.98.13
Amber	refinement	14

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1310
Number of shifts mapped to atoms	1310
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	94%

6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.83±0.01	0±0/530 (0.0± 0.0%)	1.30±0.02	0±0/714 (0.0± 0.0%)
2	B	0.80±0.01	0±0/154 (0.0± 0.0%)	1.34±0.05	0±1/204 (0.2± 0.4%)
All	All	0.82	0/13680 (0.0%)	1.31	11/18360 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.2±0.4
All	All	0	5

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	837	PHE	CA-CB-CG	5.67	119.47	113.80	12	8
2	B	824	LYS	CA-C-N	5.21	127.22	120.44	1	1
2	B	824	LYS	C-N-CA	5.21	127.22	120.44	1	1
1	A	273	GLN	OE1-CD-NE2	-5.04	117.56	122.60	4	1

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	293	ARG	Sidechain	5

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	521	542	542	0±0
All	All	13440	14160	14160	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:295:VAL:HG21	1:A:318:ILE:HD12	0.45	1.87	18	4

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	62/72 (86%)	62±0 (100±1%)	0±0 (0±1%)	0±0 (0±0%)	100	100
2	B	17/25 (68%)	17±0 (99±2%)	0±0 (1±2%)	0±0 (0±0%)	100	100
All	All	1580/1940 (81%)	1574 (100%)	6 (0%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	61/70 (87%)	59±1 (96±2%)	2±1 (4±2%)	30	81
2	B	17/25 (68%)	17±0 (100±0%)	0±0 (0±0%)	100	100
All	All	1560/1900 (82%)	1514 (97%)	46 (3%)	38	86

All 9 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	309	LEU	18
1	A	320	GLN	14
1	A	308	TYR	5
1	A	277	LEU	3
1	A	278	LYS	2
1	A	284	LYS	1
1	A	303	ASN	1
1	A	288	LYS	1
1	A	330	ASP	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.6 Ligand geometry ⓘ

There are no ligands in this entry.

6.7 Other polymers ⓘ

There are no such molecules in this entry.

6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 94% for the well-defined parts and 93% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *pfZNHIT3_NUFIP1_dss_corrARG.str*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1310
Number of shifts mapped to atoms	1310
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	5

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	97	-0.43 ± 0.26	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	95	0.29 ± 0.06	None needed (< 0.5 ppm)
$^{13}\text{C}'$	96	-0.28 ± 0.15	None needed (< 0.5 ppm)
^{15}N	93	0.28 ± 0.38	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 94%, i.e. 1118 atoms were assigned a chemical shift out of a possible 1188. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	394/394 (100%)	158/158 (100%)	158/158 (100%)	78/78 (100%)
Sidechain	657/698 (94%)	440/450 (98%)	208/223 (93%)	9/25 (36%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	67/96 (70%)	35/47 (74%)	32/46 (70%)	0/3 (0%)
Overall	1118/1188 (94%)	633/655 (97%)	398/427 (93%)	87/106 (82%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 93%, i.e. 1307 atoms were assigned a chemical shift out of a possible 1405. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	478/483 (99%)	192/194 (99%)	193/194 (99%)	93/95 (98%)
Sidechain	744/791 (94%)	499/509 (98%)	236/257 (92%)	9/25 (36%)
Aromatic	85/131 (65%)	45/64 (70%)	40/63 (63%)	0/4 (0%)
Overall	1307/1405 (93%)	736/767 (96%)	469/514 (91%)	102/124 (82%)

7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	270	THR	HG1	5.84	0.08 – 2.19	22.3
1	A	300	THR	HG1	5.28	0.08 – 2.19	19.6
1	A	293	ARG	HG2	-0.75	0.26 – 2.87	-8.9
1	A	274	LYS	HG2	-0.24	0.13 – 2.61	-6.5
1	A	278	LYS	HA	2.13	2.15 – 6.37	-5.0


7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:



8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1800
Intra-residue ($ i-j =0$)	551
Sequential ($ i-j =1$)	471
Medium range ($ i-j >1$ and $ i-j <5$)	459
Long range ($ i-j \geq 5$)	179
Inter-chain	140
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	147
Number of unmapped restraints	0
Number of restraints per residue	20.1
Number of long range restraints per residue ¹	1.8

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	1.0	0.14
0.2-0.5 (Medium)	None	None
>0.5 (Large)	None	None

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins (°)	Average number of violations per model	Max (°)
1.0-10.0 (Small)	0.9	8.16
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None

9 Distance violation analysis


9.1 Summary of distance violations

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue ($i-j =0$)	551	30.6	1	0.2	0.1	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	453	25.2	1	0.2	0.1	0	0.0	0.0
Sidechain-Sidechain	98	5.4	0	0.0	0.0	0	0.0	0.0
Sequential ($i-j =1$)	471	26.2	4	0.8	0.2	0	0.0	0.0
Backbone-Backbone	107	5.9	4	3.7	0.2	0	0.0	0.0
Backbone-Sidechain	325	18.1	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	39	2.2	0	0.0	0.0	0	0.0	0.0
Medium range ($i-j >1$ & $i-j <5$)	459	25.5	1	0.2	0.1	0	0.0	0.0
Backbone-Backbone	169	9.4	1	0.6	0.1	0	0.0	0.0
Backbone-Sidechain	209	11.6	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	81	4.5	0	0.0	0.0	0	0.0	0.0
Long range ($i-j \geq 5$)	179	9.9	1	0.6	0.1	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	72	4.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	107	5.9	1	0.9	0.1	0	0.0	0.0
Inter-chain	140	7.8	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	1	0.1	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	56	3.1	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	83	4.6	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1800	100.0	7	0.4	0.4	0	0.0	0.0
Backbone-Backbone	277	15.4	5	1.8	0.3	0	0.0	0.0
Backbone-Sidechain	1115	61.9	1	0.1	0.1	0	0.0	0.0
Sidechain-Sidechain	408	22.7	1	0.2	0.1	0	0.0	0.0

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	0	0	0	1	0	1	0.1	0.1	0.0	0.1
2	0	0	0	1	0	1	0.11	0.11	0.0	0.11
3	0	1	0	1	0	2	0.12	0.13	0.02	0.12
4	1	0	0	0	0	1	0.12	0.12	0.0	0.12
5	0	0	0	0	0	0	0.0	0.0	0.0	0.0
6	1	1	0	0	0	2	0.11	0.11	0.0	0.11
7	0	0	0	1	0	1	0.12	0.12	0.0	0.12
8	0	1	0	1	0	2	0.11	0.11	0.0	0.11
9	0	0	0	1	0	1	0.11	0.11	0.0	0.11
10	0	0	0	0	0	0	0.0	0.0	0.0	0.0

Continued on next page...


Continued from previous page...

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
11	1	0	0	0	0	1	0.13	0.13	0.0	0.13
12	0	0	0	0	0	0	0.0	0.0	0.0	0.0
13	0	1	1	1	0	3	0.11	0.11	0.0	0.11
14	0	0	0	1	0	1	0.11	0.11	0.0	0.11
15	1	0	0	0	0	1	0.13	0.13	0.0	0.13
16	0	0	0	1	0	1	0.11	0.11	0.0	0.11
17	0	0	0	1	0	1	0.13	0.13	0.0	0.13
18	1	0	0	0	0	1	0.14	0.14	0.0	0.14
19	0	1	0	1	0	2	0.11	0.11	0.0	0.11
20	0	0	0	1	0	1	0.11	0.11	0.0	0.11

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model ⓘ



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1793(IR:550, SQ:467, MR:458, LR:178, IC:140) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
0	3	1	0	0	4	1	5.0
0	1	0	0	0	1	2	10.0
0	0	0	0	0	0	3	15.0
0	0	0	0	0	0	4	20.0
1	0	0	0	0	1	5	25.0
0	0	0	0	0	0	6	30.0
0	0	0	0	0	0	7	35.0
0	0	0	0	0	0	8	40.0
0	0	0	0	0	0	9	45.0
0	0	0	0	0	0	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	1	0	1	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
0	0	0	0	0	0	16	80.0
0	0	0	0	0	0	17	85.0
0	0	0	0	0	0	18	90.0
0	0	0	0	0	0	19	95.0
0	0	0	0	0	0	20	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶ Number of models with violations


9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)



9.4 Most violated distance restraints in the ensemble [i](#)

9.4.1 Histogram : Distribution of mean distance violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD11	12	0.11	0.01	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD12	12	0.11	0.01	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD13	12	0.11	0.01	0.11
(1,435)	1:315:A:ASP:H	1:315:A:ASP:HB3	5	0.13	0.01	0.13
(1,25)	1:268:A:MET:H	1:269:A:LEU:H	2	0.11	0.0	0.11

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,435)	1:315:A:ASP:H	1:315:A:ASP:HB3	18	0.14
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD11	3	0.13
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD12	3	0.13
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD13	3	0.13
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD11	17	0.13
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD12	17	0.13
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD13	17	0.13
(1,435)	1:315:A:ASP:H	1:315:A:ASP:HB3	11	0.13
(1,435)	1:315:A:ASP:H	1:315:A:ASP:HB3	15	0.13
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD11	7	0.12
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD12	7	0.12
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD13	7	0.12
(1,435)	1:315:A:ASP:H	1:315:A:ASP:HB3	4	0.12
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD11	2	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD12	2	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD13	2	0.11

Continued on next page...

Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD11	8	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD12	8	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD13	8	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD11	9	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD12	9	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD13	9	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD11	13	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD12	13	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD13	13	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD11	14	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD12	14	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD13	14	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD11	16	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD12	16	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD13	16	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD11	19	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD12	19	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD13	19	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD11	20	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD12	20	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD13	20	0.11
(1,600)	1:330:A:ASP:HA	1:332:A:THR:H	13	0.11
(1,435)	1:315:A:ASP:H	1:315:A:ASP:HB3	6	0.11
(1,25)	1:268:A:MET:H	1:269:A:LEU:H	8	0.11
(1,25)	1:268:A:MET:H	1:269:A:LEU:H	13	0.11
(1,17)	1:266:A:TYR:HA	1:267:A:ASP:H	19	0.11
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD11	1	0.1
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD12	1	0.1
(1,1087)	1:295:A:VAL:HB	1:318:A:ILE:HD13	1	0.1
(1,406)	1:312:A:TYR:H	1:313:A:ILE:HA	6	0.1
(1,24)	1:267:A:ASP:HA	1:268:A:MET:H	3	0.1

10 Dihedral-angle violation analysis [i](#)


10.1 Summary of dihedral-angle violations [i](#)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
PHI	74	50.3	3	4.1	2.0	0	0.0	0.0
PSI	73	49.7	2	2.7	1.4	0	0.0	0.0
Total	147	100.0	5	3.4	3.4	0	0.0	0.0

¹ percentage calculated with respect to total number of dihedral-angle restraints, ² percentage calculated with respect to number of restraints in a particular dihedral-angle type, ³ violated in at least one model, ⁴ violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations [i](#)




Violated and consistently violated restraints are shown using different hatch patterns in their respective categories

10.2 Dihedral-angle violation statistics for each model

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Number of violations			Mean (°)	Max (°)	SD (°)	Median (°)
	PHI	PSI	Total				
1	1	0	1	4.39	4.39	0.0	4.39
2	0	0	0	0.0	0.0	0.0	0.0
3	1	0	1	1.1	1.1	0.0	1.1
4	1	0	1	1.39	1.39	0.0	1.39
5	0	2	2	3.22	4.22	1.0	3.22
6	1	0	1	1.07	1.07	0.0	1.07
7	1	0	1	1.01	1.01	0.0	1.01
8	0	0	0	0.0	0.0	0.0	0.0
9	0	0	0	0.0	0.0	0.0	0.0
10	1	0	1	5.13	5.13	0.0	5.13
11	0	1	1	4.61	4.61	0.0	4.61
12	0	0	0	0.0	0.0	0.0	0.0
13	1	0	1	7.25	7.25	0.0	7.25
14	1	1	2	1.44	1.49	0.05	1.44
15	0	1	1	1.33	1.33	0.0	1.33
16	2	0	2	2.26	2.96	0.7	2.26
17	1	0	1	8.16	8.16	0.0	8.16
18	1	1	2	3.22	5.36	2.14	3.22
19	0	0	0	0.0	0.0	0.0	0.0
20	0	0	0	0.0	0.0	0.0	0.0

10.2.1 Bar graph : Dihedral violation statistics for each model ⓘ



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble ⓘ

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Number of violated restraints			Fraction of the ensemble	
PHI	PSI	Total	Count ¹	%
1	1	2	1	5.0
0	0	0	2	10.0
1	0	1	3	15.0
0	0	0	4	20.0
0	1	1	5	25.0
0	0	0	6	30.0
0	0	0	7	35.0
1	0	1	8	40.0
0	0	0	9	45.0
0	0	0	10	50.0
0	0	0	11	55.0


Continued on next page...

Continued from previous page...

Number of violated restraints			Fraction of the ensemble	
PHI	PSI	Total	Count ¹	%
0	0	0	12	60.0
0	0	0	13	65.0
0	0	0	14	70.0
0	0	0	15	75.0
0	0	0	16	80.0
0	0	0	17	85.0
0	0	0	18	90.0
0	0	0	19	95.0
0	0	0	20	100.0

¹ Number of models with violations

10.3.1 Bar graph : Dihedral-angle Violation statistics for the ensemble ⓘ




10.4 Most violated dihedral-angle restraints in the ensemble ⓘ

10.4.1 Histogram : Distribution of mean dihedral-angle violations ⓘ

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models

in the ensemble



10.4.2 Table: Most violated dihedral-angle restraints [i](#)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.


Key	Atom-1	Atom-2	Atom-3	Atom-4	Models ¹	Mean	SD ²	Median
(1,65)	1:303:A:ASN:C	1:304:A:ASP:N	1:304:A:ASP:CA	1:304:A:ASP:C	8	3.31	2.47	2.17
(1,78)	1:310:A:SER:N	1:310:A:SER:CA	1:310:A:SER:C	1:311:A:HIS:N	5	2.15	1.29	1.49
(1,24)	1:280:A:ASP:C	1:281:A:HIS:N	1:281:A:HIS:CA	1:281:A:HIS:C	3	4.24	2.53	4.39

¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)

10.5 All violated dihedral-angle restraints [i](#)

10.5.1 Histogram : Distribution of violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



10.5.2 Table: All violated dihedral-angle restraints [i](#)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,65)	1:303:A:ASN:C	1:304:A:ASP:N	1:304:A:ASP:CA	1:304:A:ASP:C	17	8.16
(1,24)	1:280:A:ASP:C	1:281:A:HIS:N	1:281:A:HIS:CA	1:281:A:HIS:C	13	7.25
(1,65)	1:303:A:ASN:C	1:304:A:ASP:N	1:304:A:ASP:CA	1:304:A:ASP:C	18	5.36
(1,65)	1:303:A:ASN:C	1:304:A:ASP:N	1:304:A:ASP:CA	1:304:A:ASP:C	10	5.13
(1,78)	1:310:A:SER:N	1:310:A:SER:CA	1:310:A:SER:C	1:311:A:HIS:N	11	4.61
(1,24)	1:280:A:ASP:C	1:281:A:HIS:N	1:281:A:HIS:CA	1:281:A:HIS:C	1	4.39
(1,64)	1:303:A:ASN:N	1:303:A:ASN:CA	1:303:A:ASN:C	1:304:A:ASP:N	5	4.22
(1,65)	1:303:A:ASN:C	1:304:A:ASP:N	1:304:A:ASP:CA	1:304:A:ASP:C	16	2.96
(1,78)	1:310:A:SER:N	1:310:A:SER:CA	1:310:A:SER:C	1:311:A:HIS:N	5	2.22
(1,79)	1:310:A:SER:C	1:311:A:HIS:N	1:311:A:HIS:CA	1:311:A:HIS:C	16	1.56
(1,78)	1:310:A:SER:N	1:310:A:SER:CA	1:310:A:SER:C	1:311:A:HIS:N	14	1.49
(1,65)	1:303:A:ASN:C	1:304:A:ASP:N	1:304:A:ASP:CA	1:304:A:ASP:C	4	1.39
(1,65)	1:303:A:ASN:C	1:304:A:ASP:N	1:304:A:ASP:CA	1:304:A:ASP:C	14	1.39
(1,78)	1:310:A:SER:N	1:310:A:SER:CA	1:310:A:SER:C	1:311:A:HIS:N	15	1.33
(1,65)	1:303:A:ASN:C	1:304:A:ASP:N	1:304:A:ASP:CA	1:304:A:ASP:C	3	1.1
(1,78)	1:310:A:SER:N	1:310:A:SER:CA	1:310:A:SER:C	1:311:A:HIS:N	18	1.09
(1,24)	1:280:A:ASP:C	1:281:A:HIS:N	1:281:A:HIS:CA	1:281:A:HIS:C	6	1.07
(1,65)	1:303:A:ASN:C	1:304:A:ASP:N	1:304:A:ASP:CA	1:304:A:ASP:C	7	1.01