



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:49 AM UTC

PDB ID : 2QEP / pdb_00002qep
Title : Crystal structure of the D1 domain of PTPRN2 (IA2beta)
Authors : Ugochukwu, E.; Barr, A.; Alfano, I.; Berridge, G.; Burgess-Brown, N.; Das, S.; Fedorov, O.; King, O.; Niesen, F.; Watt, S.; Savitsky, P.; Salah, E.; Pike, A.C.W.; Bunkoczi, G.; von Delft, F.; Sundstrom, M.; Edwards, A.; Arrow-smith, C.H.; Weigelt, J.; Knapp, S.; Structural Genomics Consortium (SGC)
Deposited on : 2007-06-26
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	304	
1	B	304	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4453 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Receptor-type tyrosine-protein phosphatase N2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	288	Total	C	N	O	S	0	0	0
			2222	1408	380	419	15			
1	B	287	Total	C	N	O	S	0	0	0
			2191	1391	376	410	14			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	714	MET	-	cloning artifact	UNP Q92932
A	1011	ALA	-	cloning artifact	UNP Q92932
A	1012	HIS	-	cloning artifact	UNP Q92932
A	1013	HIS	-	cloning artifact	UNP Q92932
A	1014	HIS	-	cloning artifact	UNP Q92932
A	1015	HIS	-	cloning artifact	UNP Q92932
A	1016	HIS	-	cloning artifact	UNP Q92932
A	1017	HIS	-	cloning artifact	UNP Q92932
B	714	MET	-	cloning artifact	UNP Q92932
B	1011	ALA	-	cloning artifact	UNP Q92932
B	1012	HIS	-	cloning artifact	UNP Q92932
B	1013	HIS	-	cloning artifact	UNP Q92932
B	1014	HIS	-	cloning artifact	UNP Q92932
B	1015	HIS	-	cloning artifact	UNP Q92932
B	1016	HIS	-	cloning artifact	UNP Q92932
B	1017	HIS	-	cloning artifact	UNP Q92932

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	23	Total 23	O 23	0	0
3	B	16	Total 16	O 16	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Receptor-type tyrosine-protein phosphatase N2



- Molecule 1: Receptor-type tyrosine-protein phosphatase N2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	131.54Å 136.56Å 35.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	59.23 – 2.50 59.23 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.1 (59.23-2.50) 99.2 (59.23-2.50)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.09 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.3.0037	Depositor
R, R_{free}	0.236 , 0.286 0.242 , 0.291	Depositor DCC
R_{free} test set	1186 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	38.3	Xtriage
Anisotropy	0.110	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.068 for k,h,-l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	4453	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/2277	0.87	1/3108 (0.0%)
1	B	0.71	1/2245 (0.0%)	0.90	2/3067 (0.1%)
All	All	0.71	1/4522 (0.0%)	0.88	3/6175 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	771	VAL	CA-CB	5.55	1.56	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	885	PHE	N-CA-C	6.14	118.63	108.99
1	A	885	PHE	N-CA-C	5.35	117.20	109.07
1	B	931	ASN	N-CA-C	5.17	118.75	112.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2222	0	2050	32	0
1	B	2191	0	2008	31	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	1	0	0	0	0
3	A	23	0	0	0	0
3	B	16	0	0	0	0
All	All	4453	0	4058	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:876:VAL:HG21	1:A:891:TYR:CD1	2.23	0.73
1:A:952:SER:O	1:A:956:VAL:HG23	1.95	0.66
1:A:911:TRP:CZ3	1:A:923:LEU:HD21	2.31	0.66
1:A:876:VAL:HG21	1:A:891:TYR:CE1	2.31	0.65
1:B:735:MET:HG2	1:B:1003:ALA:HB2	1.79	0.63
1:A:991:THR:HG22	1:A:993:GLU:N	2.14	0.63
1:A:839:CYS:SG	1:A:942:ILE:HG13	2.38	0.62
1:B:752:LEU:HD13	1:B:978:LEU:HD23	1.82	0.62
1:B:974:ILE:HD12	1:B:999:LEU:HD23	1.81	0.61
1:A:917:PRO:HG2	1:A:923:LEU:HD22	1.84	0.59
1:B:991:THR:HG22	1:B:993:GLU:H	1.67	0.58
1:B:974:ILE:HD12	1:B:999:LEU:CD2	2.34	0.58
1:B:947:ASP:OD1	1:B:949:ALA:N	2.38	0.56
1:B:881:TRP:CE2	1:B:929:LYS:HE2	2.40	0.55
1:A:963:ASN:O	1:A:966:ALA:HB3	2.07	0.54
1:A:808:ASP:CG	1:A:960:MET:HE3	2.33	0.54
1:A:954:THR:HG23	1:A:998:ALA:HB2	1.88	0.54
1:B:725:ILE:HG23	1:B:1010:LEU:HD22	1.90	0.54
1:B:991:THR:HG22	1:B:993:GLU:N	2.24	0.53
1:A:974:ILE:HD12	1:A:999:LEU:CD2	2.41	0.51
1:A:827:VAL:HG21	1:A:857:TYR:CE2	2.46	0.51
1:A:991:THR:HG22	1:A:993:GLU:H	1.74	0.51
1:B:960:MET:SD	1:B:981:LEU:HD21	2.50	0.51
1:B:764:VAL:HG11	1:B:786:ARG:CZ	2.41	0.50
1:B:775:ARG:NH2	1:B:823:LEU:HD23	2.27	0.50
1:B:937:ARG:O	1:B:939:CYS:N	2.46	0.49
1:A:824:PRO:HD3	1:A:857:TYR:HB2	1.95	0.48
1:A:827:VAL:HG13	1:A:859:TYR:HB2	1.96	0.48
1:B:968:GLY:O	1:B:970:LYS:N	2.47	0.48
1:B:867:LEU:HD11	1:B:895:LEU:CD1	2.43	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:834:VAL:HG22	1:A:942:ILE:HD12	1.96	0.47
1:A:752:LEU:HD13	1:A:978:LEU:HD23	1.96	0.46
1:A:974:ILE:HD12	1:A:999:LEU:HD22	1.98	0.46
1:A:844:MET:HG3	1:A:944:HIS:NE2	2.32	0.45
1:B:759:PRO:O	1:B:784:HIS:CE1	2.70	0.45
1:A:808:ASP:CB	1:A:960:MET:HE3	2.47	0.45
1:B:801:ILE:HG23	1:B:803:ALA:HB3	1.98	0.44
1:A:923:LEU:HD23	1:A:997:PHE:CE2	2.52	0.44
1:B:844:MET:HG3	1:B:944:HIS:CE1	2.53	0.43
1:B:865:SER:HB3	1:B:874:ASN:ND2	2.34	0.43
1:B:827:VAL:HG21	1:B:857:TYR:CE2	2.53	0.43
1:A:775:ARG:NH2	1:A:856:CYS:HA	2.34	0.43
1:A:735:MET:HG3	1:A:1003:ALA:HB2	2.00	0.43
1:B:839:CYS:SG	1:B:942:ILE:HG12	2.59	0.42
1:A:991:THR:CG2	1:A:993:GLU:H	2.33	0.42
1:B:779:VAL:HG11	1:B:949:ALA:HB2	2.01	0.42
1:B:974:ILE:CD1	1:B:999:LEU:HD23	2.47	0.42
1:B:787:VAL:HG11	1:B:833:MET:HG3	2.00	0.42
1:B:831:TRP:CE2	1:B:873:VAL:HG21	2.55	0.42
1:B:886:LEU:HD12	1:B:886:LEU:C	2.45	0.42
1:A:876:VAL:CG2	1:A:891:TYR:CD1	2.99	0.41
1:B:827:VAL:HG13	1:B:859:TYR:HB2	2.02	0.41
1:A:803:ALA:HB2	1:A:819:THR:OG1	2.21	0.41
1:B:775:ARG:CZ	1:B:823:LEU:HD23	2.51	0.41
1:A:764:VAL:HG11	1:A:786:ARG:CZ	2.51	0.41
1:A:822:PRO:HG3	1:A:830:PHE:CD1	2.56	0.41
1:A:957:LEU:O	1:A:961:VAL:HG23	2.19	0.41
1:A:945:CYS:SG	1:A:946:SER:N	2.90	0.41
1:A:935:ARG:HA	1:A:936:GLY:HA2	1.84	0.40
1:B:830:PHE:O	1:B:834:VAL:HG23	2.21	0.40
1:A:845:LEU:HD11	1:A:955:TYR:HB2	2.03	0.40
1:B:780:LEU:O	1:B:781:THR:C	2.65	0.40
1:B:952:SER:O	1:B:956:VAL:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	284/304 (93%)	269 (95%)	14 (5%)	1 (0%)	30	49
1	B	281/304 (92%)	264 (94%)	15 (5%)	2 (1%)	18	34
All	All	565/608 (93%)	533 (94%)	29 (5%)	3 (0%)	24	43

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	938	SER
1	A	989	VAL
1	B	989	VAL

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/270 (84%)	223 (98%)	4 (2%)	51	77
1	B	220/270 (82%)	212 (96%)	8 (4%)	31	58
All	All	447/540 (83%)	435 (97%)	12 (3%)	39	67

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	745	LEU
1	A	746	GLU
1	A	781	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	816	TYR
1	B	744	ARG
1	B	766	GLN
1	B	781	THR
1	B	806	ILE
1	B	840	VAL
1	B	866	ASN
1	B	898	ASN
1	B	942	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	766	GLN
1	A	795	HIS
1	A	879	HIS
1	B	766	GLN
1	B	802	ASN
1	B	866	ASN
1	B	874	ASN
1	B	879	HIS
1	B	896	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	288/304 (94%)	0.69	16 (5%) 30 26	31, 38, 44, 53	0
1	B	287/304 (94%)	0.66	15 (5%) 33 29	31, 38, 44, 55	0
All	All	575/608 (94%)	0.67	31 (5%) 31 27	31, 38, 44, 55	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	811	PRO	4.5
1	A	938	SER	4.5
1	A	811	PRO	4.3
1	B	1012	HIS	3.9
1	B	936	GLY	3.7
1	A	809	HIS	3.1
1	A	778	ALA	3.1
1	B	968	GLY	3.0
1	A	777	LEU	2.8
1	A	730	MET	2.8
1	B	1011	ALA	2.8
1	B	807	MET	2.7
1	A	936	GLY	2.7
1	B	1007	ASN	2.7
1	B	862	ASP	2.6
1	B	938	SER	2.6
1	A	970	LYS	2.4
1	A	1010	LEU	2.4
1	A	971	GLU	2.3
1	A	1004	GLU	2.3
1	A	807	MET	2.3
1	B	980	HIS	2.3
1	A	810	ASP	2.2
1	B	777	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	816	TYR	2.1
1	B	970	LYS	2.1
1	B	971	GLU	2.1
1	A	812	ARG	2.1
1	A	939	CYS	2.1
1	B	754	ALA	2.1
1	A	1012	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CL	A	1	1/1	0.97	0.05	43,43,43,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.