



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:09 PM UTC

PDB ID : 7QEL / pdb_00007qel
Title : Structure of the mouse 8-oxoguanine DNA Glycosylase mOGG1 in complex with ligand TH011247
Authors : Davies, J.R.; Scaletti, E.R.; Stenmark, P.
Deposited on : 2021-12-03
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	318	3% 95% ••
1	BBB	318	26% 91% 5% ••
1	CCC	318	18% 88% 9% •

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7333 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called N-glycosylase/DNA lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	312	Total	C	N	O	S	0	0	0
			2447	1560	443	433	11			
1	BBB	306	Total	C	N	O	S	0	0	0
			2340	1501	416	412	11			
1	CCC	311	Total	C	N	O	S	0	1	0
			2370	1514	423	422	11			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	8	GLY	-	expression tag	UNP O08760
AAA	10	HIS	SER	conflict	UNP O08760
BBB	8	GLY	-	expression tag	UNP O08760
BBB	10	HIS	SER	conflict	UNP O08760
CCC	8	GLY	-	expression tag	UNP O08760
CCC	10	HIS	SER	conflict	UNP O08760

- Molecule 2 is (2 {R})- {N}-(4,5,6,7-tetrahydro-1,2-benzoxazol-3-yl)spiro[2.3]hexane-2-carboxamide (CCD ID: AUS) (formula: C₁₄H₁₈N₂O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	AAA	1	Total	C	N	O	0	0
			18	14	2	2		
2	BBB	1	Total	C	N	O	0	0
			18	14	2	2		
2	CCC	1	Total	C	N	O	0	0
			18	14	2	2		

- Molecule 3 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	AAA	1	Total	Ni	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	75	Total	O	0	0
			75	75		
4	BBB	22	Total	O	0	0
			22	22		
4	CCC	24	Total	O	0	0
			24	24		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: N-glycosylase/DNA lyase



- Molecule 1: N-glycosylase/DNA lyase



- Molecule 1: N-glycosylase/DNA lyase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.35Å 81.40Å 168.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.73 – 2.50 40.73 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (40.73-2.50) 100.0 (40.73-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.252 , 0.295 0.253 , 0.294	Depositor DCC
R_{free} test set	1906 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	51.6	Xtriage
Anisotropy	0.067	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 37.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.008 for k,h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7333	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AUS, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.00	0/2517	1.47	0/3433
1	BBB	1.03	0/2407	1.49	0/3291
1	CCC	1.03	0/2437	1.48	0/3331
All	All	1.02	0/7361	1.48	0/10055

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2447	0	2357	3	0
1	BBB	2340	0	2218	4	0
1	CCC	2370	0	2222	8	0
2	AAA	18	0	0	0	0
2	BBB	18	0	0	0	0
2	CCC	18	0	0	0	0
3	AAA	1	0	0	0	0
4	AAA	75	0	0	0	0
4	BBB	22	0	0	0	0
4	CCC	24	0	0	0	0
All	All	7333	0	6797	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:CCC:132:LEU:HD21	1:CCC:256:LEU:HG	1.69	0.73
1:CCC:193:GLU:HB2	1:CCC:209:ARG:HH12	1.59	0.67
1:BBB:229:ARG:HG3	1:BBB:259:LEU:HA	1.90	0.53
1:AAA:100:PHE:O	1:AAA:131:ARG:HD3	2.09	0.52
1:CCC:113:TRP:HE3	1:CCC:120:PHE:CZ	2.31	0.48
1:CCC:188:ALA:HB1	1:CCC:213:LYS:HG2	1.97	0.46
1:CCC:114:ALA:HB1	1:CCC:121:GLN:HG2	1.97	0.45
1:AAA:229:ARG:HG3	1:AAA:259:LEU:HA	1.99	0.44
1:CCC:66:LEU:HD21	1:CCC:96:LEU:HD21	1.99	0.44
1:BBB:284:LYS:O	1:BBB:285:THR:C	2.61	0.43
1:CCC:87:ARG:HG2	1:CCC:88:PRO:HD2	2.01	0.43
1:BBB:52:PRO:O	1:BBB:53:ALA:HB3	2.20	0.42
1:CCC:100:PHE:O	1:CCC:131:ARG:HD3	2.21	0.41
1:BBB:51:SER:O	1:BBB:52:PRO:C	2.63	0.40
1:AAA:219:GLN:HB3	1:AAA:224:TRP:CG	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	308/318 (97%)	294 (96%)	14 (4%)	0	100	100
1	BBB	300/318 (94%)	277 (92%)	19 (6%)	4 (1%)	9	18
1	CCC	306/318 (96%)	283 (92%)	19 (6%)	4 (1%)	9	18
All	All	914/954 (96%)	854 (93%)	52 (6%)	8 (1%)	14	27

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	CCC	199	LEU
1	BBB	60	ALA
1	BBB	71	ASP
1	BBB	53	ALA
1	CCC	286	SER
1	CCC	283	PRO
1	BBB	52	PRO
1	CCC	190	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	251/266 (94%)	248 (99%)	3 (1%)	63	83
1	BBB	232/266 (87%)	224 (97%)	8 (3%)	32	60
1	CCC	233/266 (88%)	222 (95%)	11 (5%)	23	47
All	All	716/798 (90%)	694 (97%)	22 (3%)	35	62

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	58	VAL
1	AAA	216	LEU
1	AAA	272	TRP
1	BBB	15	THR
1	BBB	75	CYS
1	BBB	79	ARG
1	BBB	177	THR
1	BBB	206	ARG
1	BBB	227	GLN
1	BBB	271	VAL
1	BBB	272	TRP
1	CCC	26	ILE
1	CCC	93	LEU
1	CCC	128	GLN
1	CCC	162	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	CCC	194	THR
1	CCC	198	LYS
1	CCC	209	ARG
1	CCC	216	LEU
1	CCC	271	VAL
1	CCC	272	TRP
1	CCC	323	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	AUS	CCC	401	-	17,21,21	2.64	6 (35%)	19,32,32	3.77	5 (26%)
2	AUS	AAA	401	-	17,21,21	2.54	6 (35%)	19,32,32	3.63	5 (26%)
2	AUS	BBB	401	-	17,21,21	2.64	6 (35%)	19,32,32	3.74	4 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AUS	CCC	401	-	-	0/8/31/31	0/4/4/4
2	AUS	AAA	401	-	-	0/8/31/31	0/4/4/4
2	AUS	BBB	401	-	-	0/8/31/31	0/4/4/4

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	BBB	401	AUS	C02-N03	5.94	1.47	1.36
2	CCC	401	AUS	C02-N03	5.88	1.47	1.36
2	AAA	401	AUS	C02-N03	5.66	1.46	1.36
2	CCC	401	AUS	C12-C07	5.58	1.55	1.49
2	BBB	401	AUS	C12-C07	5.42	1.55	1.49
2	AAA	401	AUS	C12-C07	5.21	1.55	1.49
2	BBB	401	AUS	C04-N03	4.35	1.47	1.38
2	CCC	401	AUS	C04-N03	4.20	1.47	1.38
2	AAA	401	AUS	C04-N03	4.07	1.46	1.38
2	BBB	401	AUS	C04-N05	3.27	1.35	1.30
2	CCC	401	AUS	C04-N05	3.23	1.35	1.30
2	AAA	401	AUS	C04-N05	3.19	1.35	1.30
2	CCC	401	AUS	O06-N05	-2.73	1.37	1.42
2	AAA	401	AUS	O06-N05	-2.66	1.37	1.42
2	BBB	401	AUS	O06-N05	-2.64	1.37	1.42
2	BBB	401	AUS	O01-C02	-2.19	1.19	1.23
2	CCC	401	AUS	O01-C02	-2.10	1.19	1.23
2	AAA	401	AUS	O01-C02	-2.09	1.19	1.23

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	CCC	401	AUS	C16-C15-C13	13.61	156.87	125.86
2	BBB	401	AUS	C16-C15-C13	13.49	156.59	125.86
2	AAA	401	AUS	C16-C15-C13	13.13	155.77	125.86
2	CCC	401	AUS	C18-C15-C13	-6.11	111.93	125.86
2	BBB	401	AUS	C18-C15-C13	-5.91	112.39	125.86
2	AAA	401	AUS	C18-C15-C13	-5.80	112.65	125.86
2	BBB	401	AUS	O06-N05-C04	5.12	107.71	104.15
2	CCC	401	AUS	O06-N05-C04	4.85	107.52	104.15
2	AAA	401	AUS	O06-N05-C04	4.48	107.26	104.15

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	CCC	401	AUS	O06-C07-C08	-2.68	107.69	110.53
2	AAA	401	AUS	O06-C07-C08	-2.52	107.86	110.53
2	BBB	401	AUS	O06-C07-C08	-2.49	107.90	110.53
2	AAA	401	AUS	C16-C15-C18	-2.13	89.18	91.16
2	CCC	401	AUS	C11-C12-C07	2.01	113.56	110.26


There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	AAA	312/318 (98%)	0.44	10 (3%)	50 46	28, 45, 71, 93	1 (0%)
1	BBB	306/318 (96%)	1.41	83 (27%)	1 1	39, 71, 100, 106	0
1	CCC	311/318 (97%)	1.30	57 (18%)	3 3	34, 70, 96, 100	1 (0%)
All	All	929/954 (97%)	1.05	150 (16%)	4 4	28, 63, 95, 106	2 (0%)

All (150) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	93	LEU	4.6
1	CCC	85	VAL	4.3
1	CCC	180	GLY	4.1
1	BBB	77	VAL	4.1
1	BBB	26	ILE	4.0
1	CCC	80	GLY	3.9
1	BBB	172	GLN	3.8
1	BBB	41	SER	3.7
1	CCC	287	GLN	3.6
1	BBB	24	ALA	3.5
1	CCC	247	GLY	3.4
1	BBB	45	PHE	3.3
1	BBB	22	LEU	3.3
1	BBB	96	LEU	3.3
1	BBB	171	ILE	3.3
1	CCC	220	GLY	3.2
1	BBB	75	CYS	3.2
1	AAA	289	LYS	3.2
1	CCC	285	THR	3.2
1	BBB	86	SER	3.2
1	CCC	142	PHE	3.2
1	BBB	85	VAL	3.2
1	BBB	205	ALA	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	CCC	323	LEU	3.1
1	CCC	217	GLU	3.1
1	AAA	81	ASP	3.1
1	BBB	173	LEU	3.1
1	BBB	23	TRP	3.1
1	CCC	284	LYS	3.0
1	CCC	120	PHE	3.0
1	CCC	324	ARG	3.0
1	AAA	272	TRP	3.0
1	CCC	290	GLY	3.0
1	BBB	208	VAL	3.0
1	AAA	11	MET	3.0
1	BBB	187	LEU	3.0
1	BBB	206	ARG	3.0
1	BBB	103	ASP	3.0
1	BBB	16	LEU	2.9
1	CCC	141	LEU	2.9
1	AAA	82	ASP	2.9
1	CCC	201	LEU	2.9
1	CCC	211	SER	2.9
1	CCC	60	ALA	2.9
1	AAA	10	HIS	2.8
1	BBB	11	MET	2.8
1	BBB	10	HIS	2.8
1	BBB	190	PRO	2.8
1	BBB	74	TYR	2.8
1	BBB	73	LEU	2.8
1	CCC	203	TYR	2.7
1	CCC	268	ASP	2.7
1	BBB	38	VAL	2.7
1	BBB	321	ALA	2.7
1	BBB	146	CYS	2.7
1	CCC	116	VAL	2.7
1	BBB	69	THR	2.7
1	BBB	175	ASP	2.7
1	BBB	220	GLY	2.7
1	BBB	33	LEU	2.7
1	BBB	201	LEU	2.7
1	AAA	80	GLY	2.6
1	CCC	147	SER	2.6
1	BBB	21	ALA	2.6
1	CCC	272	TRP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	BBB	163	CYS	2.6
1	CCC	114	ALA	2.6
1	BBB	272	TRP	2.6
1	BBB	59	LEU	2.6
1	CCC	149	ASN	2.6
1	CCC	150	ASN	2.6
1	BBB	30	ARG	2.6
1	BBB	99	TYR	2.6
1	BBB	64	TRP	2.5
1	BBB	323	LEU	2.5
1	BBB	158	MET	2.5
1	BBB	291	PRO	2.5
1	BBB	127	PHE	2.5
1	CCC	26	ILE	2.5
1	BBB	222	PRO	2.5
1	BBB	141	LEU	2.5
1	CCC	125	GLN	2.5
1	CCC	188	ALA	2.5
1	BBB	65	THR	2.5
1	CCC	291	PRO	2.4
1	BBB	207	TYR	2.4
1	BBB	216	LEU	2.4
1	BBB	84	GLN	2.4
1	AAA	60	ALA	2.4
1	BBB	17	SER	2.4
1	BBB	203	TYR	2.4
1	CCC	293	PRO	2.4
1	BBB	35	LEU	2.4
1	CCC	270	HIS	2.4
1	BBB	159	VAL	2.4
1	BBB	188	ALA	2.4
1	BBB	53	ALA	2.4
1	BBB	167	GLY	2.3
1	BBB	180	GLY	2.3
1	BBB	290	GLY	2.3
1	BBB	152	ILE	2.3
1	CCC	106	LEU	2.3
1	CCC	108[A]	GLN	2.3
1	AAA	71	ASP	2.3
1	CCC	59	LEU	2.3
1	CCC	104	VAL	2.3
1	BBB	319	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	CCC	205	ALA	2.3
1	BBB	169	ARG	2.3
1	CCC	16	LEU	2.2
1	BBB	209	ARG	2.2
1	CCC	319	PHE	2.2
1	BBB	40	ALA	2.2
1	BBB	68	GLN	2.2
1	CCC	96	LEU	2.2
1	BBB	147	SER	2.2
1	CCC	317	VAL	2.2
1	BBB	97	HIS	2.2
1	BBB	185	HIS	2.2
1	BBB	155	ILE	2.2
1	BBB	63	VAL	2.1
1	BBB	165	ALA	2.1
1	CCC	196	LEU	2.1
1	CCC	225	LEU	2.1
1	AAA	270	HIS	2.1
1	CCC	230	VAL	2.1
1	CCC	158	MET	2.1
1	CCC	90	LEU	2.1
1	CCC	74	TYR	2.1
1	CCC	168	PRO	2.1
1	BBB	28	CYS	2.1
1	CCC	322	ASP	2.1
1	BBB	125	GLN	2.1
1	CCC	271	VAL	2.1
1	BBB	78	TYR	2.1
1	BBB	322	ASP	2.1
1	CCC	178	TYR	2.1
1	CCC	146	CYS	2.1
1	CCC	190	PRO	2.1
1	BBB	153	ALA	2.0
1	BBB	170	LEU	2.0
1	CCC	200	GLY	2.0
1	BBB	88	PRO	2.0
1	BBB	124	ALA	2.0
1	CCC	294	LEU	2.0
1	BBB	200	GLY	2.0
1	CCC	221	GLY	2.0
1	BBB	20	PRO	2.0
1	BBB	25	SER	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	CCC	18	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	AUS	BBB	401	18/18	0.86	0.18	71,72,77,77	0
2	AUS	AAA	401	18/18	0.89	0.13	49,51,52,52	0
2	AUS	CCC	401	18/18	0.89	0.15	77,78,84,84	0
3	NI	AAA	402	1/1	1.00	0.05	37,37,37,37	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around AUS BBB 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around AUS AAA 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around AUS CCC 401:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.