



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:30 PM UTC

PDB ID : 8QEZ / pdb_00008qez
Title : Crystal structure of the AMPA receptor GluA2-L504Y-N775S ligand binding domain in complex with L-glutamate and positive allosteric modulator BPAM395 at 1.55Å resolution
Authors : Dorosz, J.; Laulumaa, S.; Frydenvang, K.; Kastrup, J.S.
Deposited on : 2023-09-01
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	264	4% 94% 5%
1	B	264	2% 97% .
1	C	264	2% 94% 6% .

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 13824 atoms, of which 6556 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate receptor 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	264	Total	C	H	N	O	S	0	11	0
			4263	1350	2149	348	401	15			
1	B	264	Total	C	H	N	O	S	0	9	0
			4245	1342	2137	349	402	15			
1	C	264	Total	C	H	N	O	S	0	11	0
			4267	1351	2157	344	400	15			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP P19491
A	2	ALA	-	expression tag	UNP P19491
A	94	TYR	LEU	engineered mutation	UNP P19491
A	118	GLY	-	linker	UNP P19491
A	119	THR	-	linker	UNP P19491
A	242	SER	ASN	engineered mutation	UNP P19491
B	1	GLY	-	expression tag	UNP P19491
B	2	ALA	-	expression tag	UNP P19491
B	94	TYR	LEU	engineered mutation	UNP P19491
B	118	GLY	-	linker	UNP P19491
B	119	THR	-	linker	UNP P19491
B	242	SER	ASN	engineered mutation	UNP P19491
C	1	GLY	-	expression tag	UNP P19491
C	2	ALA	-	expression tag	UNP P19491
C	94	TYR	LEU	engineered mutation	UNP P19491
C	118	GLY	-	linker	UNP P19491
C	119	THR	-	linker	UNP P19491
C	242	SER	ASN	engineered mutation	UNP P19491

- Molecule 2 is GLUTAMIC ACID (CCD ID: GLU) (formula: C₅H₉NO₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			18	5	8	1	4		
2	B	1	Total	C	H	N	O	0	0
			18	5	8	1	4		
2	C	1	Total	C	H	N	O	0	0
			18	5	8	1	4		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Zn	0	0
			4	4		
3	B	1	Total	Zn	0	0
			1	1		
3	C	2	Total	Zn	0	0
			2	2		

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			7	2	3	2		
4	A	1	Total	C	H	O	0	0
			7	2	3	2		
4	A	1	Total	C	H	O	0	0
			7	2	3	2		
4	A	1	Total	C	H	O	0	0
			7	2	3	2		
4	B	1	Total	C	H	O	0	0
			7	2	3	2		
4	B	1	Total	C	H	O	0	0
			7	2	3	2		
4	B	1	Total	C	H	O	0	0
			7	2	3	2		
4	C	1	Total	C	H	O	0	0
			7	2	3	2		

- Molecule 5 is 6-chloranyl-4-cyclopropyl-2,3-dihydrothieno[3,2-e][1,2,4]thiadiazine 1,1-dioxide (CCD ID: UF5) (formula: C₈H₉ClN₂O₂S₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
5	A	1	Total	C	Cl	H	N	O	S	0	0
			24	8	1	9	2	2	2		
5	B	1	Total	C	Cl	H	N	O	S	0	0
			24	8	1	9	2	2	2		
5	C	1	Total	C	Cl	H	N	O	S	0	0
			24	8	1	9	2	2	2		

- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Cl 1 1	0	0
6	B	1	Total Cl 1 1	0	0
6	C	1	Total Cl 1 1	0	0

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	1	Total	C	H	O	0	0
			14	3	8	3		
7	C	1	Total	C	H	O	0	0
			14	3	8	3		
7	C	1	Total	C	H	O	0	0
			14	3	8	3		
7	C	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 8 is CACODYLATE ION (CCD ID: CAC') (formula: $C_2H_6AsO_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	C	1	Total	As	C	H	O	0	0
			11	1	2	6	2		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	239	Total	O	0	0
			239	239		
9	B	249	Total	O	0	4
			249	249		
9	C	301	Total	O	0	1
			302	302		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutamate receptor 2



- Molecule 1: Glutamate receptor 2



- Molecule 1: Glutamate receptor 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	114.32Å 163.42Å 47.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.17 – 1.55 49.17 – 1.55	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.17-1.55) 100.0 (49.17-1.55)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 1.55Å)	Xtriage
Refinement program	PHENIX (1.20_4459: ???)	Depositor
R, R_{free}	0.168 , 0.195 (Not available) , 0.189	Depositor DCC
R_{free} test set	6594 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	20.0	Xtriage
Anisotropy	0.136	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13824	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UF5, GOL, ACT, ZN, CAC, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/2185	0.45	0/2935
1	B	0.29	0/2170	0.47	0/2914
1	C	0.31	0/2181	0.51	0/2930
All	All	0.29	0/6536	0.48	0/8779

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2114	2149	2165	9	0
1	B	2108	2137	2144	6	0
1	C	2110	2157	2167	14	0
2	A	10	8	5	0	0
2	B	10	8	5	0	0
2	C	10	8	5	0	0
3	A	4	0	0	0	0
3	B	1	0	0	0	0
3	C	2	0	0	0	0
4	A	16	12	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	12	9	9	1	0
4	C	4	3	3	0	0
5	A	15	9	0	0	0
5	B	15	9	0	0	0
5	C	15	9	0	0	0
6	A	1	0	0	1	0
6	B	1	0	0	0	0
6	C	1	0	0	1	0
7	B	6	8	8	0	0
7	C	18	24	24	0	0
8	C	5	6	0	0	0
9	A	239	0	0	3	0
9	B	249	0	0	2	0
9	C	302	0	0	4	0
All	All	7268	6556	6547	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:C:310:CL:CL	9:C:649:HOH:O	2.39	0.77
1:A:198:GLU:OE1	9:A:401:HOH:O	2.06	0.74
1:C:198:GLU:OE1	9:C:401:HOH:O	2.07	0.72
1:A:209:MET:HE2	1:A:211:VAL:HG12	1.82	0.61
1:C:100[B]:ILE:HD11	1:C:102:PHE:CE2	2.38	0.59
1:B:77[B]:GLU:OE1	9:B:401:HOH:O	2.17	0.58
1:B:210:LYS:HE3	4:B:303:ACT:H2	1.90	0.52
1:C:100[B]:ILE:HD13	1:C:223:ALA:HB1	1.92	0.52
1:A:149:ARG:NH2	9:A:406:HOH:O	2.44	0.51
1:B:25:MET:HE2	1:B:25:MET:HA	1.93	0.51
1:B:109:LEU:C	1:B:109:LEU:HD22	2.37	0.50
1:C:109:LEU:HD22	1:C:109:LEU:C	2.38	0.49
1:A:243[B]:GLU:HG3	9:A:460:HOH:O	2.11	0.49
1:A:109:LEU:C	1:A:109:LEU:HD22	2.38	0.49
1:C:23:HIS:HA	1:C:26:LEU:HD23	1.94	0.48
1:C:152:ILE:N	1:C:152:ILE:HD12	2.28	0.48
1:B:241:LEU:HD22	1:B:246:LEU:HD22	1.95	0.48
1:C:178:VAL:HG21	1:C:200[B]:ILE:HD13	1.96	0.47
1:C:152:ILE:HD13	9:C:441:HOH:O	2.14	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:125:GLU:CG	1:C:129:LYS:HE2	2.46	0.45
1:B:81:GLY:HA2	9:B:477:HOH:O	2.17	0.45
1:C:81:GLY:HA2	9:C:433:HOH:O	2.16	0.44
1:C:101:ASP:OD1	1:C:226:LYS:NZ	2.46	0.44
1:A:95[B]:VAL:CG2	6:A:311:CL:CL	3.03	0.43
1:A:203[B]:ARG:HH22	1:A:258:LYS:HB2	1.84	0.42
1:A:137:THR:HG22	1:A:191:LEU:HB2	2.02	0.42
1:C:178:VAL:CG2	1:C:200[B]:ILE:HD13	2.52	0.40
1:A:71:TRP:HH2	1:A:95[A]:VAL:HG12	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/264 (103%)	269 (98%)	4 (2%)	0	100	100
1	B	271/264 (103%)	268 (99%)	3 (1%)	0	100	100
1	C	273/264 (103%)	271 (99%)	2 (1%)	0	100	100
All	All	817/792 (103%)	808 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/219 (105%)	224 (97%)	6 (3%)	40	13
1	B	228/219 (104%)	226 (99%)	2 (1%)	70	51
1	C	230/219 (105%)	227 (99%)	3 (1%)	61	36
All	All	688/657 (105%)	677 (98%)	11 (2%)	59	28

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	97	GLU
1	A	109	LEU
1	A	139[A]	ASP
1	A	139[B]	ASP
1	A	230[A]	LEU
1	A	230[B]	LEU
1	B	109	LEU
1	B	226	LYS
1	C	109	LEU
1	C	139	ASP
1	C	226	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 29 ligands modelled in this entry, 10 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
8	CAC	C	304	3	2,4,4	2.55	1 (50%)	4,6,6	2.01	2 (50%)
4	ACT	C	305	-	3,3,3	1.06	0	3,3,3	1.53	1 (33%)
4	ACT	B	303	-	3,3,3	1.56	1 (33%)	3,3,3	1.22	0
4	ACT	A	306	-	3,3,3	1.10	0	3,3,3	1.61	1 (33%)
7	GOL	C	307	-	5,5,5	1.21	0	5,5,5	0.67	0
4	ACT	A	309	3	3,3,3	1.73	1 (33%)	3,3,3	1.32	0
5	UF5	A	310	-	14,17,17	1.99	4 (28%)	17,27,27	2.46	3 (17%)
7	GOL	B	306	-	5,5,5	0.99	0	5,5,5	0.98	0
4	ACT	B	304	-	3,3,3	1.22	0	3,3,3	1.37	0
2	GLU	B	301	-	8,9,9	1.18	1 (12%)	8,11,11	0.86	0
2	GLU	C	301	-	8,9,9	1.01	1 (12%)	8,11,11	0.86	0
7	GOL	C	306	-	5,5,5	0.86	0	5,5,5	0.84	0
2	GLU	A	301	-	8,9,9	1.10	1 (12%)	8,11,11	0.85	0
4	ACT	A	307	3	3,3,3	1.66	1 (33%)	3,3,3	1.13	0
4	ACT	A	308	-	3,3,3	1.25	0	3,3,3	1.60	1 (33%)
5	UF5	C	309	-	14,17,17	1.62	4 (28%)	17,27,27	2.03	3 (17%)
5	UF5	B	307	-	14,17,17	1.88	4 (28%)	17,27,27	2.15	3 (17%)
7	GOL	C	308	-	5,5,5	0.62	0	5,5,5	1.24	1 (20%)
4	ACT	B	305	-	3,3,3	1.24	0	3,3,3	1.58	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLU	B	301	-	-	0/9/9/9	-
7	GOL	C	307	-	-	4/4/4/4	-
7	GOL	C	306	-	-	0/4/4/4	-
2	GLU	A	301	-	-	0/9/9/9	-
5	UF5	A	310	-	-	0/4/22/22	0/2/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	C	308	-	-	2/4/4/4	-
7	GOL	B	306	-	-	4/4/4/4	-
5	UF5	C	309	-	-	0/4/22/22	0/2/3/3
5	UF5	B	307	-	-	0/4/22/22	0/2/3/3
2	GLU	C	301	-	-	0/9/9/9	-

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	307	UF5	S2-N1	4.27	1.68	1.63
5	C	309	UF5	S2-N1	4.18	1.68	1.63
5	A	310	UF5	S2-N1	4.11	1.68	1.63
5	A	310	UF5	C3-C4	3.91	1.39	1.36
5	B	307	UF5	C3-C4	3.65	1.39	1.36
8	C	304	CAC	AS-C2	3.11	1.97	1.90
5	A	310	UF5	O1-S2	2.82	1.47	1.43
5	A	310	UF5	O2-S2	2.81	1.47	1.43
5	B	307	UF5	O2-S2	2.56	1.47	1.43
4	A	307	ACT	CH3-C	2.54	1.59	1.49
2	A	301	GLU	OXT-C	-2.40	1.23	1.30
2	B	301	GLU	OXT-C	-2.39	1.23	1.30
4	A	309	ACT	CH3-C	2.38	1.58	1.49
5	B	307	UF5	O1-S2	2.19	1.46	1.43
5	C	309	UF5	O2-S2	2.18	1.46	1.43
2	C	301	GLU	OXT-C	-2.16	1.23	1.30
5	C	309	UF5	O1-S2	2.08	1.46	1.43
5	C	309	UF5	C3-C4	2.07	1.38	1.36
4	B	303	ACT	CH3-C	2.04	1.57	1.49

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	310	UF5	C4-S2-N1	-7.45	91.59	101.30
5	B	307	UF5	C4-S2-N1	-6.66	92.62	101.30
5	A	310	UF5	O1-S2-N1	5.36	112.92	107.92
5	C	309	UF5	C4-S2-N1	-5.22	94.50	101.30
5	C	309	UF5	O1-S2-N1	4.11	111.75	107.92
5	B	307	UF5	O1-S2-N1	3.80	111.47	107.92
5	C	309	UF5	O2-S2-N1	3.72	111.39	107.92
5	B	307	UF5	O2-S2-O1	2.60	119.39	116.97
8	C	304	CAC	O1-AS-C2	-2.36	108.57	111.50
7	C	308	GOL	C3-C2-C1	-2.23	103.60	111.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	310	UF5	C5-N2-C3	2.09	123.98	119.28
4	A	306	ACT	OXT-C-O	2.05	129.63	122.03
4	C	305	ACT	OXT-C-O	2.04	129.59	122.03
8	C	304	CAC	O2-AS-C1	2.03	110.76	105.84
4	B	305	ACT	O-C-CH3	-2.03	114.20	122.53
4	A	308	ACT	O-C-CH3	-2.00	114.32	122.53

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	C	307	GOL	C1-C2-C3-O3
7	B	306	GOL	O2-C2-C3-O3
7	B	306	GOL	C1-C2-C3-O3
7	C	308	GOL	O1-C1-C2-C3
7	C	308	GOL	O1-C1-C2-O2
7	C	307	GOL	O2-C2-C3-O3
7	B	306	GOL	O1-C1-C2-C3
7	C	307	GOL	O1-C1-C2-C3
7	B	306	GOL	O1-C1-C2-O2
7	C	307	GOL	O1-C1-C2-O2

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	303	ACT	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand GLU B 301





Ligand GLU C 301





Ligand UF5 C 309





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/264 (100%)	-0.01	10 (3%) 44 52	11, 28, 57, 84	11 (4%)
1	B	264/264 (100%)	-0.15	6 (2%) 61 69	10, 26, 49, 80	9 (3%)
1	C	264/264 (100%)	-0.33	6 (2%) 61 69	8, 22, 46, 77	11 (4%)
All	All	792/792 (100%)	-0.16	22 (2%) 55 63	8, 25, 54, 84	31 (3%)

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	263	SER	3.4
1	A	263	SER	3.4
1	B	263	SER	3.2
1	A	149	ARG	2.8
1	B	264	GLY	2.8
1	C	66	ALA	2.7
1	B	258	LYS	2.7
1	C	139	ASP	2.5
1	A	131	THR	2.5
1	A	172[A]	ARG	2.5
1	A	2	ALA	2.4
1	A	154	VAL	2.4
1	B	139	ASP	2.3
1	A	258	LYS	2.3
1	A	152	ILE	2.3
1	B	262	GLY	2.3
1	C	264	GLY	2.3
1	C	140	SER	2.2
1	A	153	ALA	2.2
1	A	227	GLY	2.2
1	B	140	SER	2.1
1	C	26	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ACT	A	306	4/4	0.77	0.21	33,44,49,50	7
3	ZN	C	303	1/1	0.78	0.17	58,58,58,58	1
4	ACT	B	303	4/4	0.78	0.18	35,42,45,46	0
7	GOL	C	308	6/6	0.78	0.17	36,52,68,68	0
7	GOL	C	307	6/6	0.80	0.16	29,45,61,61	0
3	ZN	A	305	1/1	0.80	0.11	96,96,96,96	1
4	ACT	B	305	4/4	0.85	0.18	34,35,45,45	7
4	ACT	A	309	4/4	0.86	0.16	36,47,71,71	0
4	ACT	B	304	4/4	0.87	0.14	28,36,43,43	7
4	ACT	C	305	4/4	0.87	0.14	28,34,37,37	0
4	ACT	A	307	4/4	0.88	0.14	38,40,48,48	0
7	GOL	B	306	6/6	0.89	0.11	30,45,64,77	0
4	ACT	A	308	4/4	0.91	0.12	26,37,45,45	7
7	GOL	C	306	6/6	0.91	0.11	27,45,53,62	0
6	CL	B	308	1/1	0.95	0.10	38,38,38,38	1
6	CL	C	310	1/1	0.95	0.09	31,31,31,31	1
5	UF5	A	310	15/15	0.96	0.07	17,20,25,35	0
5	UF5	B	307	15/15	0.97	0.06	17,20,22,30	0
5	UF5	C	309	15/15	0.97	0.06	15,18,20,28	0
6	CL	A	311	1/1	0.97	0.06	31,31,31,31	1
2	GLU	A	301	10/10	0.98	0.04	15,19,25,25	0
2	GLU	B	301	10/10	0.98	0.04	16,19,22,22	0
3	ZN	A	303	1/1	0.98	0.05	36,36,36,36	1
3	ZN	A	304	1/1	0.98	0.04	27,27,27,27	0
2	GLU	C	301	10/10	0.99	0.04	14,16,20,20	0
3	ZN	C	302	1/1	0.99	0.02	19,19,19,19	0
3	ZN	A	302	1/1	0.99	0.02	23,23,23,23	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	CAC	C	304	5/5	0.99	0.06	18,26,27,27	11
3	ZN	B	302	1/1	1.00	0.02	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around UF5 A 310:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UF5 B 307:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UF5 C 309:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around GLU A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GLU B 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.