



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:23 PM UTC

PDB ID : 1QF8 / pdb_00001qf8
Title : TRUNCATED FORM OF CASEIN KINASE II BETA SUBUNIT (2-182)
FROM HOMO SAPIENS
Authors : Chantalat, L.; Leroy, D.; Filhol, O.; Nueda, A.; Benitez, M.J.; Chambaz, E.;
Cochet, C.; Dideberg, O.
Deposited on : 1999-04-07
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	182	 75% 16% • 8%
1	B	182	 73% 15% • 10%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3031 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CASEIN KINASE II.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	167	Total	C	N	O	S	Se	3	2	0
			1363	875	225	248	6	9			
1	B	163	Total	C	N	O	S	Se	4	3	0
			1341	859	221	247	6	8			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	52	MSE	MET	modified residue	UNP P67870
A	78	MSE	MET	modified residue	UNP P67870
A	97	MSE	MET	modified residue	UNP P67870
A	119	MSE	MET	modified residue	UNP P67870
A	132	MSE	MET	modified residue	UNP P67870
A	141	MSE	MET	modified residue	UNP P67870
A	166	MSE	MET	modified residue	UNP P67870
A	169	MSE	MET	modified residue	UNP P67870
B	52	MSE	MET	modified residue	UNP P67870
B	78	MSE	MET	modified residue	UNP P67870
B	97	MSE	MET	modified residue	UNP P67870
B	119	MSE	MET	modified residue	UNP P67870
B	132	MSE	MET	modified residue	UNP P67870
B	141	MSE	MET	modified residue	UNP P67870
B	166	MSE	MET	modified residue	UNP P67870
B	169	MSE	MET	modified residue	UNP P67870

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Mg 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	165	Total 165	O 165	0	0
4	B	159	Total 159	O 159	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: CASEIN KINASE II

Chain A: 



• Molecule 1: CASEIN KINASE II

Chain B: 



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	132.23Å 132.23Å 63.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.74	Depositor
% Data completeness (in resolution range)	99.3 (20.00-1.74)	Depositor
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 0.4	Depositor
R, R_{free}	0.194 , 0.219	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	3031	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	4/1404 (0.3%)	0.91	5/1889 (0.3%)
1	B	0.77	5/1384 (0.4%)	0.87	4/1862 (0.2%)
All	All	0.76	9/2788 (0.3%)	0.89	9/3751 (0.2%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	119	MSE	SE-CE	-11.83	1.59	1.95
1	A	119	MSE	SE-CE	-11.77	1.60	1.95
1	B	169	MSE	SE-CE	-11.44	1.61	1.95
1	A	52	MSE	SE-CE	-7.84	1.72	1.95
1	B	166	MSE	SE-CE	-7.07	1.74	1.95
1	B	141	MSE	SE-CE	-5.94	1.77	1.95
1	B	52	MSE	SE-CE	-5.73	1.78	1.95
1	A	166	MSE	SE-CE	-5.59	1.78	1.95
1	A	141	MSE	SE-CE	-5.24	1.79	1.95

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	114	CYS	N-CA-C	-5.68	106.01	113.17
1	A	171	HIS	CA-C-N	5.61	125.97	119.47
1	A	171	HIS	C-N-CA	5.61	125.97	119.47
1	A	160	GLY	N-CA-C	5.55	119.68	112.68
1	B	160	GLY	N-CA-C	5.29	118.88	112.48
1	B	144	TYR	N-CA-C	5.14	117.48	109.52
1	A	91	ASN	N-CA-C	5.13	117.27	111.11
1	B	23	CYS	N-CA-C	-5.02	102.64	110.42
1	A	114	CYS	N-CA-C	-5.01	106.95	113.16

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1363	0	1290	18	0
1	B	1341	0	1261	13	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
4	A	165	0	0	0	1
4	B	159	0	0	4	0
All	All	3031	0	2551	31	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:82:LEU:HG	4:B:338:HOH:O	1.75	0.87
1:A:178:ARG:N	1:A:179:PRO:HD3	2.06	0.70
1:A:101:TYR:CD1	1:A:120:LEU:HD13	2.31	0.64
1:A:165:HIS:O	1:A:169[A]:MSE:HG3	1.97	0.63
1:A:66:PRO:HD2	1:A:69:SER:OG	2.01	0.60
1:A:177:LYS:C	1:A:179:PRO:HD3	2.29	0.57
1:A:177:LYS:N	1:A:177:LYS:HD3	2.20	0.56
1:A:138:PRO:C	1:A:176:PRO:HG2	2.32	0.55
1:A:138:PRO:HB2	1:A:176:PRO:CG	2.39	0.53
1:A:170:VAL:C	1:A:172:PRO:HD3	2.35	0.52
1:B:7:VAL:HG12	1:B:11:SER:OG	2.11	0.51
1:B:78:MSE:HG3	4:B:338:HOH:O	2.11	0.49
1:B:78:MSE:CG	4:B:338:HOH:O	2.59	0.49
1:A:171:HIS:N	1:A:172:PRO:HD3	2.28	0.48
1:B:106:PHE:CD1	1:B:121:PRO:HD3	2.51	0.46
1:A:178:ARG:N	1:A:179:PRO:CD	2.78	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:PRO:HB2	1:A:176:PRO:HG2	1.97	0.46
1:B:27:GLU:O	1:B:31:GLN:HG3	2.17	0.45
1:B:126:ASP:OD2	1:B:161:THR:HB	2.16	0.45
1:A:167:LEU:C	1:A:167:LEU:HD23	2.44	0.43
1:A:106:PHE:CD1	1:A:121:PRO:HD3	2.54	0.43
1:B:114:CYS:HA	1:B:139:LYS:HD2	2.01	0.43
1:B:163:PHE:HB3	1:B:164:PRO:HD3	2.01	0.43
1:A:106:PHE:CG	1:A:121:PRO:HD3	2.54	0.42
1:A:163:PHE:HB3	1:A:164:PRO:HD3	2.02	0.42
1:B:158:TYR:HE1	4:B:338:HOH:O	2.02	0.42
1:A:87:TYR:OH	1:A:96:GLN:NE2	2.53	0.42
1:B:40:ASN:HA	1:B:46:TYR:CD1	2.55	0.41
1:A:69:SER:O	1:A:73:GLU:HG3	2.21	0.41
1:B:167:LEU:C	1:B:167:LEU:HD23	2.46	0.41
1:B:175:ARG:HA	1:B:176:PRO:HD3	1.96	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:67:ASN:OD1	4:A:219:HOH:O[3_554]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	165/182 (91%)	161 (98%)	3 (2%)	1 (1%)	21	8
1	B	162/182 (89%)	161 (99%)	1 (1%)	0	100	100
All	All	327/364 (90%)	322 (98%)	4 (1%)	1 (0%)	36	23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	58	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	149/153 (97%)	147 (99%)	2 (1%)	61	43
1	B	147/153 (96%)	145 (99%)	2 (1%)	59	41
All	All	296/306 (97%)	292 (99%)	4 (1%)	59	41

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	GLU
1	A	120	LEU
1	B	142	ASP
1	B	161	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	GLN
1	A	42	GLN
1	A	48	GLN
1	A	96	GLN
1	A	102	GLN
1	B	45	HIS
1	B	91	ASN
1	B	102	GLN
1	B	165	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.