



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 09:10 AM UTC

PDB ID : 1QFX / pdb_00001qfx
Title : PH 2.5 ACID PHOSPHATASE FROM ASPERGILLUS NIGER
Authors : Kostrewa, D.; Wyss, M.; D'Arcy, A.; Van Loon, A.P.G.M.
Deposited on : 1999-04-15
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	460	% 85% 10% .
1	B	460	2% 86% 10% .
2	C	7	100%
3	D	5	100%

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 7835 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (PH 2.5 ACID PHOSPHATASE).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	441	Total	C	N	O	S	0	0	0
			3440	2182	560	680	18			
1	B	443	Total	C	N	O	S	0	0	0
			3456	2190	563	685	18			

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	7	Total	C	N	O	0	0	0
			83	46	2	35			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 3 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total 6	C 3	O 3	0	0
6	A	1	Total 6	C 3	O 3	0	0
6	B	1	Total 6	C 3	O 3	0	0
6	B	1	Total 6	C 3	O 3	0	0
6	B	1	Total 6	C 3	O 3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	332	Total 332	O 332	0	0
7	B	316	Total 316	O 316	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PROTEIN (PH 2.5 ACID PHOSPHATASE)




- Molecule 1: PROTEIN (PH 2.5 ACID PHOSPHATASE)



- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	170.60Å 170.60Å 201.15Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.00 – 2.40 25.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	87.1 (25.00-2.40) 86.5 (25.00-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.87 (at 2.41Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.183 , 0.225 0.180 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	27.0	Xtriage
Anisotropy	0.312	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 65.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7835	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN, BMA, SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	0/3533	0.98	7/4820 (0.1%)
1	B	0.61	0/3549	0.99	12/4842 (0.2%)
All	All	0.61	0/7082	0.99	19/9662 (0.2%)

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	176	ASN	N-CA-C	7.23	121.81	112.92
1	A	176	ASN	N-CA-C	6.93	121.82	112.88
1	A	15	GLN	N-CA-C	6.66	118.61	111.36
1	B	414	ASN	N-CA-C	6.66	122.49	113.97
1	B	15	GLN	N-CA-C	6.63	118.59	111.36
1	A	414	ASN	N-CA-C	6.21	121.92	113.97
1	B	42	TYR	N-CA-C	-6.13	105.80	113.28
1	A	242	VAL	N-CA-C	-6.00	104.66	110.42
1	B	21	ARG	N-CA-C	5.63	120.31	113.38
1	B	242	VAL	N-CA-C	-5.62	105.03	110.42
1	B	33	ASN	N-CA-C	5.48	117.33	111.36
1	B	309	ALA	N-CA-C	5.46	119.11	112.23
1	B	205	THR	N-CA-C	-5.43	105.36	111.28
1	A	434	PHE	N-CA-C	5.42	118.69	111.75
1	B	418	TYR	N-CA-C	5.39	117.01	111.03
1	B	147	VAL	N-CA-C	5.35	113.77	107.77
1	B	351	ILE	N-CA-C	5.30	118.48	111.17
1	A	418	TYR	N-CA-C	5.29	116.90	111.03
1	A	150	PHE	N-CA-C	5.20	117.87	109.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3440	0	3207	25	0
1	B	3456	0	3217	21	0
2	C	83	0	70	0	0
3	D	61	0	52	0	0
4	A	28	0	26	0	0
4	B	28	0	26	0	0
5	A	25	0	0	0	0
5	B	30	0	0	0	0
6	A	18	0	24	0	0
6	B	18	0	24	0	0
7	A	332	0	0	3	0
7	B	316	0	0	2	0
All	All	7835	0	6646	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:111:TYR:O	1:A:112:ASN:HB2	1.88	0.72
1:B:111:TYR:O	1:B:112:ASN:HB2	1.95	0.65
1:A:174:SER:HB2	7:A:1037:HOH:O	2.01	0.60
1:B:191:ASP:HA	1:B:432:LEU:HA	1.83	0.59
1:A:145:THR:HG21	7:A:1126:HOH:O	2.01	0.59
1:B:342:ARG:HD3	7:B:979:HOH:O	2.04	0.58
1:A:43:GLY:O	1:B:262:GLN:HG3	2.06	0.56
1:A:278:TYR:OH	1:A:337:ASP:HB3	2.07	0.54
1:A:452:ALA:HB3	1:A:455:GLU:HB2	1.89	0.54
1:A:50:THR:O	1:A:51:SER:HB2	2.07	0.54
1:A:262:GLN:HG3	1:B:43:GLY:O	2.08	0.54
1:A:342:ARG:HD3	7:A:970:HOH:O	2.08	0.53
1:A:14:LYS:HB3	1:B:119:PRO:HG3	1.92	0.52
1:A:191:ASP:HA	1:A:432:LEU:HA	1.93	0.51
1:B:50:THR:O	1:B:51:SER:HB2	2.11	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:PRO:HG3	1:B:14:LYS:HB3	1.94	0.50
1:A:204:THR:HG23	1:A:205:THR:H	1.77	0.50
1:B:66:ARG:HD3	1:B:156:ARG:CD	2.42	0.50
1:B:134:LYS:HD3	1:B:168:PHE:O	2.11	0.49
1:A:106:PRO:HB2	1:A:110:TYR:CD2	2.50	0.47
1:B:36:TYR:HB2	1:B:357:MET:O	2.16	0.46
1:A:66:ARG:HD3	1:A:156:ARG:CD	2.46	0.46
1:A:62:ARG:HD2	1:A:319:ASP:HB3	1.97	0.45
1:B:278:TYR:OH	1:B:337:ASP:HB3	2.17	0.45
1:A:262:GLN:O	1:A:266:VAL:HG23	2.17	0.45
1:A:319:ASP:OD1	1:A:319:ASP:N	2.50	0.45
1:B:62:ARG:HD2	1:B:319:ASP:HB3	1.99	0.44
1:B:441:THR:HB	1:B:443:GLU:HG2	2.00	0.43
1:A:134:LYS:HD3	1:A:168:PHE:O	2.18	0.43
1:B:211:TYR:HA	1:B:271:VAL:HG13	2.00	0.43
1:A:122:GLY:O	1:A:125:ASP:HB3	2.19	0.43
1:B:107:ASN:OD1	1:B:109:CYS:HB2	2.19	0.42
1:A:36:TYR:HB2	1:A:357:MET:O	2.20	0.42
1:A:61:LYS:HG2	1:A:62:ARG:O	2.20	0.42
1:B:125:ASP:OD1	1:B:125:ASP:C	2.64	0.41
1:B:215:GLN:NE2	7:B:956:HOH:O	2.49	0.41
1:A:35:PRO:HB3	1:A:120:TYR:CD1	2.56	0.41
1:B:27:LEU:HA	1:B:30:TYR:CE2	2.56	0.41
1:B:27:LEU:HA	1:B:30:TYR:CZ	2.56	0.41
1:A:21:ARG:NH2	1:B:38:GLU:OE2	2.54	0.41
1:A:145:THR:HG22	1:A:146:VAL:N	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	435/460 (95%)	420 (97%)	15 (3%)	0	100	100
1	B	437/460 (95%)	424 (97%)	13 (3%)	0	100	100
All	All	872/920 (95%)	844 (97%)	28 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	368/384 (96%)	364 (99%)	4 (1%)	65	82
1	B	370/384 (96%)	365 (99%)	5 (1%)	59	79
All	All	738/768 (96%)	729 (99%)	9 (1%)	63	81

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	TYR
1	A	199	THR
1	A	335	ASN
1	A	409	SER
1	B	30	TYR
1	B	199	THR
1	B	308	GLU
1	B	335	ASN
1	B	409	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	176	ASN
1	A	215	GLN
1	A	454	GLN
1	B	128	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	215	GLN
1	B	258	ASN
1	B	454	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	1,2	14,14,15	0.97	1 (7%)	17,19,21	1.03	1 (5%)
2	NAG	C	2	2	14,14,15	0.56	0	17,19,21	1.06	1 (5%)
2	BMA	C	3	2	11,11,12	1.28	1 (9%)	15,15,17	1.52	3 (20%)
2	MAN	C	4	2	11,11,12	1.52	2 (18%)	15,15,17	1.25	1 (6%)
2	MAN	C	5	2	11,11,12	1.48	2 (18%)	15,15,17	1.37	4 (26%)
2	MAN	C	6	2	11,11,12	1.60	2 (18%)	15,15,17	1.49	2 (13%)
2	MAN	C	7	2	11,11,12	1.85	2 (18%)	15,15,17	1.46	2 (13%)
3	NAG	D	1	1,3	14,14,15	0.92	1 (7%)	17,19,21	0.84	0
3	NAG	D	2	3	14,14,15	0.59	0	17,19,21	1.04	2 (11%)
3	BMA	D	3	3	11,11,12	0.88	0	15,15,17	0.92	1 (6%)
3	MAN	D	4	3	11,11,12	1.47	4 (36%)	15,15,17	1.31	2 (13%)
3	MAN	D	5	3	11,11,12	1.54	4 (36%)	15,15,17	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	MAN	C	5	2	-	0/2/19/22	0/1/1/1
2	MAN	C	6	2	-	0/2/19/22	0/1/1/1
2	MAN	C	7	2	-	0/2/19/22	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	0/2/19/22	0/1/1/1
3	MAN	D	5	3	-	0/2/19/22	0/1/1/1

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	7	MAN	O5-C1	4.43	1.51	1.43
2	C	6	MAN	O5-C5	4.14	1.51	1.43
2	C	5	MAN	C2-C3	3.66	1.58	1.52
2	C	4	MAN	O5-C1	3.03	1.48	1.43
2	C	4	MAN	O2-C2	2.89	1.49	1.43
2	C	7	MAN	C2-C3	2.82	1.56	1.52
3	D	4	MAN	O5-C5	2.67	1.48	1.43
2	C	6	MAN	C2-C3	2.58	1.56	1.52
3	D	5	MAN	C2-C3	2.57	1.56	1.52
2	C	3	BMA	O5-C5	2.51	1.48	1.43
3	D	4	MAN	C2-C3	2.30	1.56	1.52
2	C	5	MAN	O2-C2	2.30	1.48	1.43
3	D	5	MAN	C1-C2	2.28	1.57	1.52
3	D	1	NAG	O5-C1	-2.27	1.39	1.43
2	C	1	NAG	C1-C2	2.19	1.55	1.52
3	D	5	MAN	C4-C5	2.19	1.57	1.53
3	D	4	MAN	C1-C2	2.15	1.57	1.52
3	D	4	MAN	O5-C1	2.07	1.47	1.43
3	D	5	MAN	O5-C5	2.04	1.47	1.43

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	7	MAN	O5-C5-C6	3.95	115.35	107.66
2	C	3	BMA	C1-O5-C5	3.79	117.26	112.19
2	C	6	MAN	O5-C5-C6	3.61	114.68	107.66
2	C	7	MAN	C1-O5-C5	3.56	116.96	112.19
2	C	4	MAN	C1-O5-C5	3.48	116.85	112.19
2	C	6	MAN	C2-C3-C4	3.46	116.94	110.86
3	D	4	MAN	C1-O5-C5	2.98	116.18	112.19
2	C	2	NAG	C3-C4-C5	-2.64	105.45	110.23
2	C	3	BMA	C3-C4-C5	2.57	114.89	110.23
3	D	2	NAG	O5-C1-C2	2.44	115.06	111.29
2	C	5	MAN	C2-C3-C4	-2.39	106.66	110.86
2	C	5	MAN	C1-C2-C3	-2.38	106.18	109.64
3	D	2	NAG	C3-C4-C5	-2.38	105.92	110.23
3	D	3	BMA	O3-C3-C2	2.14	114.42	110.05
2	C	1	NAG	C1-O5-C5	-2.11	109.36	112.19
3	D	4	MAN	C2-C3-C4	2.09	114.54	110.86
2	C	5	MAN	O5-C1-C2	-2.07	105.85	110.79
2	C	3	BMA	C1-C2-C3	2.06	112.65	109.64
2	C	5	MAN	O2-C2-C1	-2.05	104.53	109.22


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

21 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SO4	B	810	-	4,4,4	0.39	0	6,6,6	0.14	0
5	SO4	A	801	-	4,4,4	0.43	0	6,6,6	0.16	0
6	GOL	A	701	-	5,5,5	0.21	0	5,5,5	0.57	0
5	SO4	B	807	-	4,4,4	0.49	0	6,6,6	0.40	0
5	SO4	B	811	-	4,4,4	0.43	0	6,6,6	0.11	0
5	SO4	B	808	-	4,4,4	0.46	0	6,6,6	0.18	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GOL	A	702	-	5,5,5	0.22	0	5,5,5	0.98	1 (20%)
5	SO4	A	803	-	4,4,4	0.34	0	6,6,6	0.13	0
5	SO4	B	809	-	4,4,4	0.35	0	6,6,6	0.28	0
4	NAG	A	739	1	14,14,15	0.69	0	17,19,21	0.77	0
5	SO4	A	802	-	4,4,4	0.45	0	6,6,6	0.24	0
6	GOL	A	703	-	5,5,5	0.22	0	5,5,5	0.48	0
6	GOL	B	705	-	5,5,5	0.28	0	5,5,5	1.07	1 (20%)
4	NAG	B	739	1	14,14,15	0.45	0	17,19,21	0.83	0
4	NAG	B	596	1	14,14,15	1.45	2 (14%)	17,19,21	1.26	2 (11%)
5	SO4	B	806	-	4,4,4	0.49	0	6,6,6	0.21	0
6	GOL	B	704	-	5,5,5	0.33	0	5,5,5	0.38	0
4	NAG	A	596	1	14,14,15	0.81	0	17,19,21	0.75	0
6	GOL	B	706	-	5,5,5	0.34	0	5,5,5	1.55	2 (40%)
5	SO4	A	805	-	4,4,4	0.39	0	6,6,6	0.13	0
5	SO4	A	804	-	4,4,4	0.35	0	6,6,6	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	B	705	-	-	3/4/4/4	-
6	GOL	B	706	-	-	1/4/4/4	-
4	NAG	B	739	1	-	0/6/23/26	0/1/1/1
6	GOL	A	701	-	-	0/4/4/4	-
4	NAG	B	596	1	-	0/6/23/26	0/1/1/1
6	GOL	B	704	-	-	0/4/4/4	-
6	GOL	A	703	-	-	0/4/4/4	-
4	NAG	A	739	1	-	0/6/23/26	0/1/1/1
6	GOL	A	702	-	-	3/4/4/4	-
4	NAG	A	596	1	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	596	NAG	O5-C1	3.97	1.50	1.43
4	B	596	NAG	O5-C5	-2.76	1.38	1.43

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	596	NAG	O7-C7-C8	-3.44	115.93	122.05
4	B	596	NAG	C8-C7-N2	2.96	121.03	116.12
6	B	706	GOL	O2-C2-C1	2.40	119.13	109.18
6	B	706	GOL	C3-C2-C1	-2.36	103.15	111.80
6	B	705	GOL	O2-C2-C1	2.30	118.71	109.18
6	A	702	GOL	O2-C2-C1	2.10	117.89	109.18

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	702	GOL	O1-C1-C2-C3
6	B	705	GOL	O1-C1-C2-C3
6	A	702	GOL	O1-C1-C2-O2
6	A	702	GOL	O2-C2-C3-O3
6	B	705	GOL	O1-C1-C2-O2
6	B	705	GOL	O2-C2-C3-O3
6	B	706	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	441/460 (95%)	-0.52	6 (1%) 73 69	11, 22, 42, 68	0
1	B	443/460 (96%)	-0.55	8 (1%) 67 63	13, 21, 41, 76	0
All	All	884/920 (96%)	-0.53	14 (1%) 70 66	11, 22, 42, 76	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	199	THR	5.7
1	A	199	THR	3.6
1	B	89	THR	3.4
1	B	100	ASP	3.2
1	A	89	THR	2.6
1	B	198	ASP	2.5
1	A	253	PHE	2.5
1	B	253	PHE	2.2
1	B	200	ASP	2.2
1	B	204	THR	2.1
1	B	201	ASN	2.1
1	A	85	SER	2.1
1	A	229	GLY	2.0
1	A	100	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MAN	D	4	11/12	0.70	0.18	57,65,68,71	0
2	MAN	C	7	11/12	0.78	0.20	70,79,86,93	0
3	MAN	D	5	11/12	0.85	0.14	52,55,59,66	0
2	BMA	C	3	11/12	0.88	0.15	34,54,67,71	0
2	MAN	C	5	11/12	0.89	0.10	37,41,47,49	0
3	BMA	D	3	11/12	0.89	0.12	41,49,55,57	0
2	MAN	C	4	11/12	0.90	0.14	49,56,61,61	0
2	MAN	C	6	11/12	0.90	0.12	43,49,61,64	0
3	NAG	D	2	14/15	0.95	0.08	26,31,36,41	0
2	NAG	C	2	14/15	0.95	0.07	27,31,40,45	0
2	NAG	C	1	14/15	0.97	0.06	14,17,28,28	0
3	NAG	D	1	14/15	0.97	0.06	14,20,27,28	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	GOL	B	706	6/6	0.81	0.17	39,52,54,56	0
5	SO4	A	805	5/5	0.85	0.29	91,94,96,97	0
6	GOL	A	703	6/6	0.85	0.19	37,43,56,61	0
5	SO4	A	801	5/5	0.85	0.21	68,70,75,88	0
4	NAG	A	596	14/15	0.87	0.11	29,39,45,49	0
5	SO4	B	807	5/5	0.88	0.15	46,49,63,69	0
6	GOL	B	704	6/6	0.89	0.15	39,46,54,59	0
5	SO4	B	811	5/5	0.89	0.22	63,76,83,85	0
4	NAG	B	596	14/15	0.90	0.14	25,41,48,48	0
5	SO4	A	802	5/5	0.90	0.17	61,65,74,75	0
6	GOL	A	701	6/6	0.92	0.12	30,38,40,52	0
5	SO4	B	809	5/5	0.92	0.19	71,74,76,77	0
5	SO4	B	810	5/5	0.93	0.21	76,81,86,90	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	SO4	B	808	5/5	0.93	0.12	53,55,60,70	0
4	NAG	B	739	14/15	0.93	0.09	24,32,38,38	0
6	GOL	A	702	6/6	0.94	0.10	21,30,35,36	0
5	SO4	A	803	5/5	0.94	0.12	52,56,61,61	0
4	NAG	A	739	14/15	0.95	0.07	13,25,29,32	0
5	SO4	B	806	5/5	0.95	0.12	47,51,61,66	0
6	GOL	B	705	6/6	0.95	0.08	21,26,34,34	0
5	SO4	A	804	5/5	0.95	0.12	68,68,72,78	0

6.5 Other polymers [i](#)

There are no such residues in this entry.