



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 02:34 AM UTC

PDB ID : 3QFZ / pdb_00003qfz
Title : Crystal Structure of Cellvibrio gilvus Cellobiose Phosphorylase Complexed with Sulfate and 1-Deoxynojirimycin
Authors : Fushinobu, S.; Hidaka, M.; Hayashi, A.M.; Wakagi, T.; Shoun, H.; Kitaoka, M.
Deposited on : 2011-01-24
Resolution : 2.39 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	842	 91% 7% .
1	B	842	 90% 7% ..

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 13721 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cellobiose Phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	822	Total	C	N	O	S	0	0	0
			6428	4066	1102	1246	14			
1	B	822	Total	C	N	O	S	0	0	0
			6428	4066	1102	1246	14			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP O66264
A	-18	GLY	-	expression tag	UNP O66264
A	-17	SER	-	expression tag	UNP O66264
A	-16	SER	-	expression tag	UNP O66264
A	-15	HIS	-	expression tag	UNP O66264
A	-14	HIS	-	expression tag	UNP O66264
A	-13	HIS	-	expression tag	UNP O66264
A	-12	HIS	-	expression tag	UNP O66264
A	-11	HIS	-	expression tag	UNP O66264
A	-10	HIS	-	expression tag	UNP O66264
A	-9	SER	-	expression tag	UNP O66264
A	-8	SER	-	expression tag	UNP O66264
A	-7	GLY	-	expression tag	UNP O66264
A	-6	LEU	-	expression tag	UNP O66264
A	-5	VAL	-	expression tag	UNP O66264
A	-4	PRO	-	expression tag	UNP O66264
A	-3	ARG	-	expression tag	UNP O66264
A	-2	GLY	-	expression tag	UNP O66264
A	-1	SER	-	expression tag	UNP O66264
A	0	HIS	-	expression tag	UNP O66264
B	-19	MET	-	expression tag	UNP O66264
B	-18	GLY	-	expression tag	UNP O66264
B	-17	SER	-	expression tag	UNP O66264
B	-16	SER	-	expression tag	UNP O66264
B	-15	HIS	-	expression tag	UNP O66264

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP O66264
B	-13	HIS	-	expression tag	UNP O66264
B	-12	HIS	-	expression tag	UNP O66264
B	-11	HIS	-	expression tag	UNP O66264
B	-10	HIS	-	expression tag	UNP O66264
B	-9	SER	-	expression tag	UNP O66264
B	-8	SER	-	expression tag	UNP O66264
B	-7	GLY	-	expression tag	UNP O66264
B	-6	LEU	-	expression tag	UNP O66264
B	-5	VAL	-	expression tag	UNP O66264
B	-4	PRO	-	expression tag	UNP O66264
B	-3	ARG	-	expression tag	UNP O66264
B	-2	GLY	-	expression tag	UNP O66264
B	-1	SER	-	expression tag	UNP O66264
B	0	HIS	-	expression tag	UNP O66264

- Molecule 2 is beta-D-glucopyranose (CCD ID: BGC) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 1-DEOXYNOJIRIMYCIN (CCD ID: NOJ) (formula: $C_6H_{13}NO_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			11	6	1	4		
4	B	1	Total	C	N	O	0	0
			11	6	1	4		

- Molecule 5 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	398	Total	O	0	0
			398	398		
6	B	381	Total	O	0	0
			381	381		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cellobiose Phosphorylase



• Molecule 1: Cellobiose Phosphorylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	84.91Å 98.13Å 104.78Å 90.00° 102.53° 90.00°	Depositor
Resolution (Å)	36.27 – 2.39 36.27 – 2.39	Depositor EDS
% Data completeness (in resolution range)	99.3 (36.27-2.39) 99.4 (36.27-2.39)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.10 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.141 , 0.208 (Not available) , 0.179	Depositor DCC
R_{free} test set	3338 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	20.5	Xtriage
Anisotropy	0.701	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 34.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13721	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EPE, BGC, NOJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.06	1/6604 (0.0%)	1.05	7/9009 (0.1%)
1	B	1.08	1/6604 (0.0%)	1.06	5/9009 (0.1%)
All	All	1.07	2/13208 (0.0%)	1.06	12/18018 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	629	THR	CA-CB	5.59	1.61	1.53
1	A	761	ILE	CA-CB	5.00	1.61	1.54

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	608	GLY	CA-C-N	7.00	127.58	119.47
1	B	608	GLY	C-N-CA	7.00	127.58	119.47
1	A	608	GLY	CA-C-N	6.05	125.74	119.56
1	A	608	GLY	C-N-CA	6.05	125.74	119.56
1	B	713	MET	CB-CA-C	5.73	119.67	109.65

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6428	0	6071	19	0
1	B	6428	0	6071	23	0
2	A	12	0	12	1	0
2	B	12	0	12	0	0
3	A	15	0	0	0	0
3	B	10	0	0	0	0
4	A	11	0	13	0	0
4	B	11	0	13	0	0
5	A	15	0	17	1	0
6	A	398	0	0	1	0
6	B	381	0	0	3	0
All	All	13721	0	12209	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 42 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:631:HIS:HD2	1:B:696:ARG:HH12	1.12	0.97
1:B:640:TYR:H	1:B:660:ASN:HD21	1.33	0.76
1:A:99:HIS:HD2	1:A:104:SER:OG	1.81	0.64
1:B:334:MET:HG3	1:B:692:THR:CG2	2.30	0.62
1:B:645:ILE:HG22	6:B:881:HOH:O	2.03	0.58

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	776/842 (92%)	744 (96%)	32 (4%)	0	100	100
1	B	771/842 (92%)	732 (95%)	38 (5%)	1 (0%)	48	64

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1547/1684 (92%)	1476 (95%)	70 (4%)	1 (0%)	48 64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	85	TRP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	665/682 (98%)	640 (96%)	25 (4%)	29 49
1	B	665/682 (98%)	638 (96%)	27 (4%)	27 46
All	All	1330/1364 (98%)	1278 (96%)	52 (4%)	28 48

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	350	SER
1	B	503	GLU
1	B	785	ASN
1	B	362	ARG
1	B	411	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	511	GLN
1	B	631	HIS
1	B	727	ASN
1	A	727	ASN
1	A	668	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	SO4	A	3003	-	4,4,4	0.36	0	6,6,6	0.41	0
2	BGC	A	1901	-	12,12,12	0.79	0	17,17,17	1.35	2 (11%)
3	SO4	A	1902	-	4,4,4	0.38	0	6,6,6	0.66	0
4	NOJ	B	2903	-	11,11,11	0.98	1 (9%)	13,15,15	1.11	1 (7%)
4	NOJ	A	1903	-	11,11,11	1.04	1 (9%)	13,15,15	1.87	3 (23%)
3	SO4	B	2902	-	4,4,4	0.36	0	6,6,6	0.35	0
2	BGC	B	2901	-	12,12,12	0.82	0	17,17,17	1.58	4 (23%)
3	SO4	B	3002	-	4,4,4	0.18	0	6,6,6	0.39	0
3	SO4	A	3004	-	4,4,4	0.35	0	6,6,6	0.53	0
5	EPE	A	3001	-	15,15,15	0.86	1 (6%)	19,20,20	2.52	8 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	A	1901	-	-	0/2/22/22	0/1/1/1
4	NOJ	B	2903	-	-	0/2/19/19	0/1/1/1
4	NOJ	A	1903	-	-	0/2/19/19	0/1/1/1
2	BGC	B	2901	-	-	0/2/22/22	0/1/1/1
5	EPE	A	3001	-	-	3/9/19/19	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	3001	EPE	C10-S	2.74	1.81	1.77
4	B	2903	NOJ	C1-C2	2.39	1.54	1.52
4	A	1903	NOJ	C2-C3	2.28	1.56	1.52

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	3001	EPE	O3S-S-C10	5.61	116.97	106.00
5	A	3001	EPE	C5-N4-C3	5.37	120.41	108.84
4	A	1903	NOJ	C1-N5-C5	4.51	119.62	109.71
5	A	3001	EPE	C6-N1-C2	3.76	116.94	108.84
2	B	2901	BGC	O5-C1-C2	-3.36	104.38	110.30

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	3001	EPE	N4-C7-C8-O8
5	A	3001	EPE	S-C10-C9-N1
5	A	3001	EPE	C8-C7-N4-C3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1901	BGC	1	0
5	A	3001	EPE	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	822/842 (97%)	-0.67	0 100 100	17, 27, 41, 57	0
1	B	822/842 (97%)	-0.62	1 (0%) 92 90	19, 28, 42, 58	0
All	All	1644/1684 (97%)	-0.65	1 (0%) 92 90	17, 27, 41, 58	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	3004	5/5	0.75	0.13	86,86,88,89	0
5	EPE	A	3001	15/15	0.87	0.15	69,71,74,74	0
3	SO4	B	3002	5/5	0.93	0.09	59,59,60,60	0
3	SO4	A	3003	5/5	0.93	0.12	54,55,58,58	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NOJ	A	1903	11/11	0.96	0.07	26,32,34,35	0
4	NOJ	B	2903	11/11	0.98	0.04	26,28,30,32	0
2	BGC	A	1901	12/12	0.99	0.04	23,24,27,27	0
2	BGC	B	2901	12/12	0.99	0.03	17,21,24,25	0
3	SO4	A	1902	5/5	0.99	0.04	25,25,27,30	0
3	SO4	B	2902	5/5	1.00	0.03	20,21,22,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.