



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 01:55 PM UTC

PDB ID : 6QFK / pdb_00006qfk
Title : EngBF DARPin Fusion 4b G10
Authors : Ernst, P.; Pluckthun, A.; Mittl, P.R.E.
Deposited on : 2019-01-10
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1357	8% 93% 5%
2	B	15	40% 87% 13%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 12243 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PEGA domain-containing protein,PEGA domain-containing protein,EngBF DARPin fusion G10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1347	Total	C	N	O	S	0	10	0
			10412	6480	1781	2121	30			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	334	GLY	-	expression tag	UNP A0A414FD23
A	335	PRO	-	expression tag	UNP A0A414FD23
A	336	LEU	-	expression tag	UNP A0A414FD23
A	337	GLY	-	expression tag	UNP A0A414FD23
A	338	SER	-	expression tag	UNP A0A414FD23
A	339	MET	-	expression tag	UNP A0A414FD23
A	342	CYS	SER	conflict	UNP A0A414FD23
A	1309	ARG	GLN	conflict	UNP A0A414FD23


- Molecule 2 is a protein called V3-IF.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	15	Total	C	N	O	0	0	0
			113	75	20	18			

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Mn	0	0
			4	4		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
6	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1637	Total 1637	O 1637	0	0
7	B	1	Total 1	O 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PEGA domain-containing protein, PEGA domain-containing protein, EngBF DARPin fusion G10



- Molecule 2: V3-IF



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	192.89Å 192.89Å 122.92Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.25 – 2.00 29.25 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.25-2.00) 100.0 (29.25-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 2.00Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.154 , 0.176 0.158 , 0.179	Depositor DCC
R_{free} test set	8747 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	35.1	Xtriage
Anisotropy	0.118	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 57.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	12243	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN, MPD, DPR, EDO, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.83	1/10612 (0.0%)	1.18	23/14393 (0.2%)
2	B	0.74	0/109	1.18	0/145
All	All	0.83	1/10721 (0.0%)	1.18	23/14538 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	630	VAL	CA-C	5.18	1.58	1.52

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1264	LYS	CA-C-N	7.77	131.73	120.38
1	A	1264	LYS	C-N-CA	7.77	131.73	120.38
1	A	789	ASP	CA-C-N	6.64	133.92	121.97
1	A	789	ASP	C-N-CA	6.64	133.92	121.97
1	A	1537	GLN	CA-C-N	6.42	129.21	120.54
1	A	1537	GLN	C-N-CA	6.42	129.21	120.54
1	A	1603	HIS	CA-C-N	6.21	128.93	120.54
1	A	1603	HIS	C-N-CA	6.21	128.93	120.54
1	A	578	ASP	CA-CB-CG	6.13	118.73	112.60
1	A	1265	ASP	CA-CB-CG	6.10	118.70	112.60
1	A	1639	PHE	CA-CB-CG	6.00	119.80	113.80
1	A	1636	HIS	CA-C-N	5.56	127.74	120.28
1	A	1636	HIS	C-N-CA	5.56	127.74	120.28
1	A	1295	ASP	N-CA-C	-5.55	98.86	108.52
1	A	1269	LYS	N-CA-C	-5.55	105.30	112.68
1	A	368	ASP	CA-CB-CG	5.55	118.15	112.60
1	A	1555	ASP	CA-C-N	5.54	127.98	120.38
1	A	1555	ASP	C-N-CA	5.54	127.98	120.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	940	VAL	N-CA-C	5.41	115.98	108.84
1	A	970	ASP	CA-CB-CG	5.33	117.93	112.60
1	A	934	ASN	CA-CB-CG	5.17	117.77	112.60
1	A	1058	ASP	CA-CB-CG	5.08	117.68	112.60
1	A	1209	THR	N-CA-C	-5.05	105.85	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10412	0	9955	28	0
2	B	113	0	116	0	0
3	A	4	0	0	0	0
4	A	8	0	14	0	0
5	A	44	0	66	1	0
6	A	24	0	26	0	0
7	A	1637	0	0	2	0
7	B	1	0	0	0	0
All	All	12243	0	10177	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:964[B]:VAL:HG12	1:A:981:ILE:HG12	1.77	0.65
1:A:570:TRP:CD2	1:A:592:PRO:HB3	2.38	0.59
1:A:1604:LEU:HD11	1:A:1638:GLU:HG3	1.85	0.58
1:A:1545:MET:HE1	1:A:1576:VAL:HG12	1.86	0.56
1:A:940:VAL:HG21	7:A:2110:HOH:O	2.08	0.53
1:A:1364:GLY:HA2	1:A:1392:LEU:HD13	1.93	0.51
1:A:1669:ASN:HB3	1:A:1672:ILE:HD13	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1189:THR:HB	1:A:1345:GLN:HB3	1.93	0.50
1:A:1603:HIS:HB3	1:A:1606:ILE:HD12	1.94	0.50
1:A:1352:GLY:HA2	1:A:1512:ILE:HG13	1.94	0.50
1:A:1628:LEU:HD23	1:A:1660:PRO:HG2	1.95	0.49
1:A:1208:ASN:HB2	5:A:1707:EDO:H11	1.95	0.49
1:A:473:VAL:HG13	1:A:517:ILE:HD12	1.97	0.47
1:A:769[A]:VAL:HG23	1:A:811:MET:HB2	1.96	0.47
1:A:1498:LYS:HD3	7:A:2682:HOH:O	2.15	0.46
1:A:1172:ASP:H	1:A:1238:LYS:HE3	1.81	0.45
1:A:470:LEU:HD22	1:A:594:MET:HE2	2.00	0.44
1:A:1537:GLN:HB3	1:A:1540:GLU:HB3	1.99	0.44
1:A:554:THR:HA	1:A:562:SER:O	2.18	0.44
1:A:634:VAL:HG11	1:A:667:LEU:HD13	1.99	0.44
1:A:869:ASP:O	1:A:996:GLY:HA3	2.19	0.43
1:A:718:HIS:CD2	1:A:718:HIS:C	2.97	0.43
1:A:676:TYR:CD1	1:A:717:VAL:HB	2.55	0.42
1:A:634:VAL:HA	1:A:914:LYS:HG3	2.02	0.42
1:A:664:THR:HA	1:A:907:TYR:OH	2.20	0.41
1:A:1268:LEU:HD22	1:A:1347:VAL:HG23	2.02	0.41
1:A:599:ALA:HB1	1:A:602:GLU:HG2	2.03	0.41
1:A:361:VAL:HG11	1:A:364:TYR:CE1	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1355/1357 (100%)	1304 (96%)	49 (4%)	2 (0%)	48	46
2	B	13/15 (87%)	12 (92%)	1 (8%)	0	100	100
All	All	1368/1372 (100%)	1316 (96%)	50 (4%)	2 (0%)	48	46

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	790	VAL
1	A	843	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1108/1106 (100%)	1094 (99%)	14 (1%)	61	68
2	B	10/10 (100%)	8 (80%)	2 (20%)	1	0
All	All	1118/1116 (100%)	1102 (99%)	16 (1%)	59	66

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	440	LYS
1	A	632	GLU
1	A	940	VAL
1	A	941	SER
1	A	1104	VAL
1	A	1182	LYS
1	A	1227	GLU
1	A	1477	LEU
1	A	1568	GLU
1	A	1571	LEU
1	A	1604	LEU
1	A	1605	GLU
1	A	1634	GLU
1	A	1636	HIS
2	B	2	LYS
2	B	5	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	426	GLN
1	A	437	ASN
1	A	458	ASN
1	A	467	ASN
1	A	498	ASN
1	A	556	GLN
1	A	936	ASN
1	A	1055	GLN
1	A	1059	GLN
1	A	1119	ASN
1	A	1218	ASN
1	A	1244	HIS
1	A	1334	GLN
1	A	1486	GLN
1	A	1537	GLN
1	A	1547	ASN
1	A	1646	HIS
1	A	1651	ASN
1	A	1667	ASN
1	A	1669	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 18 ligands modelled in this entry, 4 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	A	1715	-	3,3,3	0.48	0	2,2,2	0.33	0
6	MES	A	1711	-	12,12,12	0.77	0	15,16,16	0.46	0
4	MPD	A	1705	-	7,7,7	0.56	0	9,10,10	0.40	0
5	EDO	A	1717	-	3,3,3	0.58	0	2,2,2	0.22	0
5	EDO	A	1716	-	3,3,3	0.45	0	2,2,2	0.46	0
5	EDO	A	1712	-	3,3,3	0.49	0	2,2,2	0.37	0
5	EDO	A	1709	-	3,3,3	0.55	0	2,2,2	0.29	0
5	EDO	A	1706	-	3,3,3	0.70	0	2,2,2	0.15	0
5	EDO	A	1707	-	3,3,3	0.55	0	2,2,2	0.18	0
6	MES	A	1708	-	12,12,12	0.77	0	15,16,16	0.37	0
5	EDO	A	1710	-	3,3,3	0.52	0	2,2,2	0.29	0
5	EDO	A	1713	-	3,3,3	0.53	0	2,2,2	0.37	0
5	EDO	A	1714	-	3,3,3	0.55	0	2,2,2	0.35	0
5	EDO	A	1718	-	3,3,3	0.53	0	2,2,2	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	1715	-	-	0/1/1/1	-
6	MES	A	1711	-	-	0/6/14/14	0/1/1/1
4	MPD	A	1705	-	-	3/5/5/5	-
5	EDO	A	1717	-	-	0/1/1/1	-
5	EDO	A	1716	-	-	0/1/1/1	-
5	EDO	A	1712	-	-	0/1/1/1	-
5	EDO	A	1709	-	-	0/1/1/1	-
5	EDO	A	1706	-	-	1/1/1/1	-
5	EDO	A	1707	-	-	0/1/1/1	-
6	MES	A	1708	-	-	0/6/14/14	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	1710	-	-	0/1/1/1	-
5	EDO	A	1713	-	-	0/1/1/1	-
5	EDO	A	1714	-	-	0/1/1/1	-
5	EDO	A	1718	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1706	EDO	O1-C1-C2-O2
4	A	1705	MPD	C1-C2-C3-C4
4	A	1705	MPD	O2-C2-C3-C4
4	A	1705	MPD	CM-C2-C3-C4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1707	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	1347/1357 (99%)	-0.18	111 (8%)	17 16	14, 37, 90, 178	10 (0%)
2	B	14/15 (93%)	1.85	6 (42%)	0 1	74, 82, 110, 120	0
All	All	1361/1372 (99%)	-0.16	117 (8%)	16 15	14, 38, 92, 178	10 (0%)

All (117) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1672	ILE	7.4
1	A	1676	LEU	6.6
1	A	1648	ALA	5.5
1	A	1640	VAL	5.3
1	A	1639	PHE	5.3
1	A	1665	ILE	5.3
1	A	1650	VAL	5.2
1	A	1604	LEU	5.2
1	A	1637	LEU	5.2
1	A	1643	LEU	5.1
1	A	1574	VAL	5.1
1	A	1606	ILE	5.0
1	A	1611	LEU	5.0
1	A	1607	VAL	4.9
1	A	1644	LEU	4.6
1	A	1571	LEU	4.5
2	B	1	PRO	4.3
1	A	1541	VAL	4.2
1	A	1664	ALA	4.2
1	A	1661	PHE	4.2
1	A	1673	ALA	4.2
1	A	1617	VAL	4.1
1	A	1615	ALA	4.0
1	A	1610	LEU	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1628	LEU	3.9
1	A	1675	VAL	3.9
1	A	1642	VAL	3.8
1	A	1573	ILE	3.7
2	B	4	ILE	3.5
1	A	1608	GLU	3.5
1	A	1601	TRP	3.5
1	A	1605	GLU	3.4
1	A	1599	ALA	3.3
2	B	14	ALA	3.3
1	A	1240	THR	3.2
1	A	1545	MET	3.2
1	A	1578	LEU	3.2
1	A	1682	LEU	3.2
1	A	1576	VAL	3.2
1	A	1577	LEU	3.1
1	A	1382	SER	3.1
1	A	1609	VAL	3.1
1	A	1603	HIS	3.0
1	A	1646	HIS	3.0
1	A	1543	ILE	3.0
1	A	1544	LEU	3.0
1	A	1618	ASN	3.0
1	A	1631	ALA	2.9
1	A	940	VAL	2.9
2	B	2	LYS	2.9
1	A	1238	LYS	2.9
1	A	1566	ALA	2.9
1	A	1638	GLU	2.9
1	A	1565	ALA	2.8
1	A	1572	GLU	2.8
1	A	1660	PRO	2.8
1	A	1535	ALA	2.8
1	A	1634	GLU	2.8
1	A	1681	LYS	2.8
1	A	1636	HIS	2.7
1	A	1663	LEU	2.7
1	A	1613	HIS	2.7
1	A	1549	ALA	2.7
1	A	1600	ALA	2.7
1	A	1581	GLY	2.7
1	A	1619	ALA	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1630	LEU	2.7
1	A	1632	ALA	2.6
1	A	1647	GLY	2.6
1	A	1680	ALA	2.6
1	A	1584	VAL	2.6
1	A	1655	CYS	2.6
1	A	1533	ALA	2.6
1	A	1645	LYS	2.6
1	A	1669	ASN	2.6
1	A	1580	THR	2.6
1	A	1265	ASP	2.6
1	A	1477	LEU	2.6
1	A	1598	ALA	2.5
1	A	1539	ASP	2.5
1	A	1478	SER	2.5
1	A	1536	GLY	2.5
1	A	1671	ASP	2.5
1	A	1641	GLU	2.5
1	A	1379	SER	2.4
1	A	1542	ARG	2.4
1	A	1616	ASP	2.4
1	A	1582	ALA	2.4
1	A	1022	ASP	2.3
1	A	1612	LYS	2.3
1	A	1527	ARG	2.3
1	A	1569	GLY	2.3
1	A	1579	LYS	2.3
1	A	1674	GLU	2.3
2	B	6	ILE	2.3
1	A	1546	ALA	2.3
1	A	1668	GLY	2.3
1	A	1684	SER	2.2
1	A	1635	GLY	2.2
1	A	1475	ALA	2.2
1	A	1525	LEU	2.2
1	A	1530	LEU	2.2
1	A	1487	ASP	2.2
1	A	1614	GLY	2.2
1	A	1670	GLU	2.2
1	A	1653	GLN	2.2
1	A	1481	ALA	2.1
1	A	1595	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1480	ALA	2.1
1	A	1532	ALA	2.1
1	A	1575	GLU	2.1
1	A	1526	ASP	2.1
1	A	1649	ASP	2.1
1	A	1570	HIS	2.1
1	A	1683	GLY	2.1
1	A	1523	LYS	2.1
2	B	5	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DPR	B	15	7/8	0.68	0.27	99,101,107,107	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	A	1718	4/4	0.76	0.21	85,86,87,88	0
6	MES	A	1711	12/12	0.78	0.30	144,145,148,149	0
5	EDO	A	1714	4/4	0.80	0.21	76,78,80,80	0
5	EDO	A	1707	4/4	0.81	0.27	81,81,82,83	0
5	EDO	A	1706	4/4	0.82	0.20	55,61,63,65	0
5	EDO	A	1712	4/4	0.84	0.26	79,80,80,81	0
4	MPD	A	1705	8/8	0.84	0.28	101,104,105,105	0
5	EDO	A	1715	4/4	0.86	0.18	65,66,68,69	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	EDO	A	1716	4/4	0.86	0.29	87,88,89,90	0
5	EDO	A	1713	4/4	0.86	0.22	88,88,88,88	0
5	EDO	A	1709	4/4	0.86	0.25	76,78,79,79	0
5	EDO	A	1717	4/4	0.88	0.18	79,80,80,80	0
5	EDO	A	1710	4/4	0.90	0.22	81,82,82,82	0
6	MES	A	1708	12/12	0.96	0.11	40,59,74,74	0
3	MN	A	1704	1/1	0.99	0.05	43,43,43,43	0
3	MN	A	1701	1/1	1.00	0.04	39,39,39,39	0
3	MN	A	1702	1/1	1.00	0.04	42,42,42,42	0
3	MN	A	1703	1/1	1.00	0.07	42,42,42,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.