



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:22 PM UTC

PDB ID : 7QFA / pdb_00007qfa
Title : Monoclinic crystal structure of PTG CBM21 in complex with beta-cyclodextrin
Authors : Semrau, M.S.; Storici, P.; Lolli, G.
Deposited on : 2021-12-05
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	134	3% 86% 10% ••
1	B	134	4% 86% 8% 6%
1	C	134	6% 80% 13% 7%
2	D	7	100%
2	K	7	43% 57%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	R	7	14% 71% 14%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3609 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Protein phosphatase 1 regulatory subunit 3C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	129	Total	C	N	O	S	0	1	0
			1069	685	176	203	5			
1	B	126	Total	C	N	O	S	0	0	0
			1031	657	172	197	5			
1	C	124	Total	C	N	O	S	0	0	0
			1018	650	169	194	5			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	SER	-	expression tag	UNP Q9UQK1
B	131	SER	-	expression tag	UNP Q9UQK1
C	131	SER	-	expression tag	UNP Q9UQK1

- Molecule 2 is an oligosaccharide called Cycloheptakis-(1-4)-(alpha-D-glucopyranose).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	D	7	Total	C	O	0	0	0
			77	42	35			
2	K	7	Total	C	O	0	0	0
			77	42	35			
2	R	7	Total	C	O	0	0	0
			77	42	35			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	90	Total	O	0	0
			90	90		
4	B	72	Total	O	0	0
			72	72		
4	C	74	Total	O	0	0
			74	74		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Protein phosphatase 1 regulatory subunit 3C




- Molecule 1: Protein phosphatase 1 regulatory subunit 3C



- Molecule 1: Protein phosphatase 1 regulatory subunit 3C



- Molecule 2: Cycloheptakis-(1-4)-(alpha-D-glucopyranose)



- Molecule 2: Cycloheptakis-(1-4)-(alpha-D-glucopyranose)



- Molecule 2: Cycloheptakis-(1-4)-(alpha-D-glucopyranose)



GLC1
GLC2
GLC3
GLC4
GLC5
GLC6
GLC7

4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	66.83Å 42.87Å 168.99Å 90.00° 100.67° 90.00°	Depositor
Resolution (Å)	41.51 – 2.00 41.51 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.7 (41.51-2.00) 98.6 (41.51-2.00)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.11.1	Depositor
R, R_{free}	0.173 , 0.212 0.176 , 0.214	Depositor DCC
R_{free} test set	1554 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	30.3	Xtriage
Anisotropy	0.436	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3609	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/1101	0.57	0/1497
1	B	0.31	0/1056	0.51	0/1432
1	C	0.32	0/1043	0.51	0/1415
All	All	0.32	0/3200	0.53	0/4344

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1069	0	1016	9	0
1	B	1031	0	983	5	0
1	C	1018	0	972	12	0
2	D	77	0	63	0	0
2	K	77	0	63	0	0
2	R	77	0	63	1	0
3	B	18	0	24	0	0
3	C	6	0	8	0	0
4	A	90	0	0	1	0
4	B	72	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	74	0	0	0	0
All	All	3609	0	3192	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:140:GLN:HE22	1:C:162:CYS:H	1.30	0.80
1:B:167:ARG:HG3	1:B:222:LEU:HB2	1.69	0.74
1:C:223:PRO:HD2	1:C:226:ILE:HD11	1.68	0.73
1:A:261:TRP:H	1:A:261:TRP:CD1	2.05	0.72
1:C:140:GLN:HB2	1:C:143:THR:HG23	1.71	0.71
1:B:167:ARG:HA	1:B:222:LEU:HD12	1.82	0.61
1:B:230:GLN:NE2	4:B:402:HOH:O	2.34	0.61
1:A:164:LEU:HD11	1:A:222:LEU:HD11	1.83	0.59
1:A:261:TRP:CD1	1:A:261:TRP:N	2.73	0.57
1:A:194:LYS:NZ	4:A:303:HOH:O	2.42	0.50
1:C:191:ASP:HA	1:C:230:GLN:HB3	2.00	0.44
1:B:239:HIS:CD2	1:B:244:VAL:HG22	2.52	0.44
1:A:233:GLU:HG2	1:A:255:ARG:HG2	2.00	0.43
1:A:261:TRP:CE2	1:C:161:ASN:HB3	2.54	0.42
1:C:140:GLN:NE2	1:C:162:CYS:H	2.07	0.42
1:C:140:GLN:HB2	1:C:143:THR:CG2	2.46	0.42
1:A:223:PRO:HD2	1:A:226:ILE:HD11	2.01	0.41
1:B:145:TYR:HD1	1:B:146:LEU:HD12	1.86	0.41
1:C:181:GLU:OE1	2:R:1:GLC:O2	2.37	0.41
1:A:159:LEU:HB2	1:A:236:ILE:HD11	2.01	0.41
1:A:204:MET:HE2	1:A:204:MET:HB3	1.88	0.41
1:C:191:ASP:HB2	1:C:230:GLN:HE21	1.86	0.41
1:C:136:LEU:HD23	1:C:256:ILE:HG12	2.03	0.40
1:C:159:LEU:HB2	1:C:236:ILE:HD11	2.02	0.40
1:C:226:ILE:HD13	1:C:232:ILE:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	128/134 (96%)	123 (96%)	5 (4%)	0	100	100
1	B	122/134 (91%)	115 (94%)	7 (6%)	0	100	100
1	C	120/134 (90%)	114 (95%)	6 (5%)	0	100	100
All	All	370/402 (92%)	352 (95%)	18 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/127 (97%)	119 (97%)	4 (3%)	33	34
1	B	119/127 (94%)	115 (97%)	4 (3%)	32	33
1	C	118/127 (93%)	115 (98%)	3 (2%)	42	45
All	All	360/381 (94%)	349 (97%)	11 (3%)	35	37

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	166	GLU
1	A	168	THR
1	A	183	LYS
1	A	261	TRP
1	B	174	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	205	LYS
1	B	212	ASP
1	B	260	GLN
1	C	167	ARG
1	C	225	VAL
1	C	259	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	177	ASN
1	B	133	ASN
1	B	151	HIS
1	B	161	ASN
1	B	252	GLN
1	C	140	GLN
1	C	155	ASN
1	C	195	ASN
1	C	230	GLN
1	C	249	ASN
1	C	258	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

21 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	D	1	2	11,11,12	1.25	1 (9%)	15,15,17	0.70	0
2	GLC	D	2	2	11,11,12	0.67	0	15,15,17	1.14	2 (13%)
2	GLC	D	3	2	11,11,12	1.18	1 (9%)	15,15,17	1.25	2 (13%)
2	GLC	D	4	2	11,11,12	1.08	1 (9%)	15,15,17	0.89	0
2	GLC	D	5	2	11,11,12	1.00	0	15,15,17	0.89	1 (6%)
2	GLC	D	6	2	11,11,12	1.08	1 (9%)	15,15,17	0.94	0
2	GLC	D	7	2	11,11,12	0.71	0	15,15,17	1.32	1 (6%)
2	GLC	K	1	2	11,11,12	1.03	0	15,15,17	1.01	0
2	GLC	K	2	2	11,11,12	1.06	0	15,15,17	0.65	0
2	GLC	K	3	2	11,11,12	0.74	0	15,15,17	1.23	1 (6%)
2	GLC	K	4	2	11,11,12	1.03	1 (9%)	15,15,17	0.82	1 (6%)
2	GLC	K	5	2	11,11,12	0.76	0	15,15,17	1.01	0
2	GLC	K	6	2	11,11,12	1.31	2 (18%)	15,15,17	1.36	2 (13%)
2	GLC	K	7	2	11,11,12	1.11	1 (9%)	15,15,17	1.00	1 (6%)
2	GLC	R	1	2	11,11,12	1.03	0	15,15,17	0.87	1 (6%)
2	GLC	R	2	2	11,11,12	1.07	1 (9%)	15,15,17	1.05	1 (6%)
2	GLC	R	3	2	11,11,12	0.80	0	15,15,17	1.29	2 (13%)
2	GLC	R	4	2	11,11,12	1.19	1 (9%)	15,15,17	1.01	1 (6%)
2	GLC	R	5	2	11,11,12	0.79	0	15,15,17	0.94	0
2	GLC	R	6	2	11,11,12	1.12	0	15,15,17	0.97	1 (6%)
2	GLC	R	7	2	11,11,12	0.97	0	15,15,17	0.98	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	D	1	2	-	0/2/19/22	0/1/1/1
2	GLC	D	2	2	-	1/2/19/22	0/1/1/1
2	GLC	D	3	2	-	0/2/19/22	0/1/1/1
2	GLC	D	4	2	-	1/2/19/22	0/1/1/1
2	GLC	D	5	2	-	0/2/19/22	0/1/1/1
2	GLC	D	6	2	-	0/2/19/22	0/1/1/1
2	GLC	D	7	2	-	0/2/19/22	0/1/1/1
2	GLC	K	1	2	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	K	2	2	-	0/2/19/22	0/1/1/1
2	GLC	K	3	2	-	0/2/19/22	0/1/1/1
2	GLC	K	4	2	-	0/2/19/22	0/1/1/1
2	GLC	K	5	2	-	0/2/19/22	0/1/1/1
2	GLC	K	6	2	-	2/2/19/22	0/1/1/1
2	GLC	K	7	2	-	0/2/19/22	0/1/1/1
2	GLC	R	1	2	-	0/2/19/22	0/1/1/1
2	GLC	R	2	2	-	1/2/19/22	0/1/1/1
2	GLC	R	3	2	-	0/2/19/22	0/1/1/1
2	GLC	R	4	2	-	0/2/19/22	0/1/1/1
2	GLC	R	5	2	-	0/2/19/22	0/1/1/1
2	GLC	R	6	2	-	0/2/19/22	0/1/1/1
2	GLC	R	7	2	-	0/2/19/22	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1	GLC	C1-C2	2.68	1.58	1.52
2	R	2	GLC	O2-C2	-2.23	1.38	1.43
2	R	4	GLC	C1-C2	2.19	1.57	1.52
2	K	7	GLC	C2-C3	2.14	1.55	1.52
2	D	3	GLC	O5-C1	-2.13	1.40	1.43
2	D	6	GLC	C1-C2	2.07	1.57	1.52
2	K	6	GLC	O2-C2	-2.05	1.39	1.43
2	K	6	GLC	C1-C2	2.05	1.57	1.52
2	D	4	GLC	C2-C3	2.05	1.55	1.52
2	K	4	GLC	O2-C2	-2.03	1.39	1.43

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	K	6	GLC	C1-O5-C5	3.68	117.12	112.19
2	D	7	GLC	C1-O5-C5	3.42	116.77	112.19
2	K	3	GLC	C1-O5-C5	3.33	116.65	112.19
2	D	3	GLC	C1-O5-C5	2.75	115.87	112.19
2	D	5	GLC	C1-O5-C5	2.55	115.60	112.19
2	D	2	GLC	C1-O5-C5	2.42	115.42	112.19
2	D	3	GLC	C6-C5-C4	-2.41	107.10	113.02
2	R	7	GLC	O4-C4-C3	-2.33	104.88	110.38
2	R	3	GLC	C1-O5-C5	2.30	115.27	112.19
2	K	4	GLC	C1-O5-C5	2.26	115.21	112.19
2	R	2	GLC	O6-C6-C5	-2.23	103.73	111.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	GLC	O6-C6-C5	-2.20	103.85	111.33
2	K	6	GLC	O5-C5-C4	2.13	116.01	110.83
2	R	4	GLC	O5-C5-C6	-2.13	103.53	107.66
2	R	3	GLC	C6-C5-C4	-2.07	107.94	113.02
2	K	7	GLC	C6-C5-C4	-2.06	107.95	113.02
2	R	1	GLC	O4-C4-C3	-2.02	105.61	110.38
2	R	6	GLC	C1-O5-C5	2.01	114.88	112.19

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	K	6	GLC	C4-C5-C6-O6
2	K	6	GLC	O5-C5-C6-O6
2	D	2	GLC	C4-C5-C6-O6
2	D	4	GLC	O5-C5-C6-O6
2	R	2	GLC	C4-C5-C6-O6

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	R	1	GLC	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.




Oligosaccharide Chain K




Bond lengths




Bond angles



Torsions



Rings



5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection.

RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	301	-	5,5,5	0.42	0	5,5,5	0.70	0
3	GOL	C	301	-	5,5,5	0.34	0	5,5,5	0.59	0
3	GOL	B	302	-	5,5,5	0.34	0	5,5,5	0.43	0
3	GOL	B	303	-	5,5,5	0.39	0	5,5,5	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	301	-	-	0/4/4/4	-
3	GOL	C	301	-	-	0/4/4/4	-
3	GOL	B	302	-	-	2/4/4/4	-
3	GOL	B	303	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	302	GOL	C1-C2-C3-O3
3	B	302	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	129/134 (96%)	-0.17	4 (3%)	51	50	19, 31, 58, 88	1 (0%)
1	B	126/134 (94%)	0.04	6 (4%)	35	34	24, 35, 66, 80	0
1	C	124/134 (92%)	0.14	8 (6%)	25	23	23, 36, 74, 86	0
All	All	379/402 (94%)	-0.00	18 (4%)	36	35	19, 34, 70, 88	1 (0%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	261	TRP	3.8
1	B	260	GLN	3.3
1	A	260	GLN	3.2
1	B	209	GLY	3.0
1	C	206	ASN	2.9
1	A	209	GLY	2.6
1	A	133	ASN	2.6
1	C	222	LEU	2.6
1	C	210	GLY	2.5
1	C	225	VAL	2.5
1	C	143	THR	2.5
1	C	259	VAL	2.4
1	B	146	LEU	2.3
1	C	227	PRO	2.2
1	C	205	LYS	2.2
1	B	206	ASN	2.1
1	B	258	HIS	2.1
1	B	133	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	D	4	11/12	0.91	0.10	27,32,41,52	0
2	GLC	D	3	11/12	0.92	0.08	25,27,31,33	0
2	GLC	D	2	11/12	0.95	0.07	20,24,30,42	0
2	GLC	D	7	11/12	0.95	0.06	17,21,27,33	0
2	GLC	D	1	11/12	0.95	0.06	19,20,27,30	0
2	GLC	D	5	11/12	0.97	0.05	20,23,30,31	0
2	GLC	D	6	11/12	0.97	0.05	18,20,24,25	0
2	GLC	K	7	11/12	-	-	22,27,34,36	0
2	GLC	K	1	11/12	-	-	27,31,42,43	0
2	GLC	K	2	11/12	-	-	32,34,42,49	0
2	GLC	K	3	11/12	-	-	30,38,43,48	0
2	GLC	K	4	11/12	-	-	26,29,32,34	0
2	GLC	K	5	11/12	-	-	25,28,32,33	0
2	GLC	K	6	11/12	-	-	27,30,34,45	0
2	GLC	R	7	11/12	-	-	22,28,37,42	0
2	GLC	R	1	11/12	-	-	22,28,36,38	0
2	GLC	R	2	11/12	-	-	23,35,38,47	0
2	GLC	R	3	11/12	-	-	22,24,26,30	0
2	GLC	R	4	11/12	-	-	17,21,26,28	0
2	GLC	R	5	11/12	-	-	23,25,29,39	0
2	GLC	R	6	11/12	-	-	22,24,29,30	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain D:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	C	301	6/6	0.83	0.17	56,62,66,70	0
3	GOL	B	303	6/6	0.88	0.14	37,55,66,67	0
3	GOL	B	302	6/6	0.88	0.12	58,60,66,67	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	B	301	6/6	0.92	0.11	34,43,47,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.