



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:12 PM UTC

PDB ID : 1QG3 / pdb_00001qg3
Title : CRYSTAL STRUCTURE OF A TANDEM PAIR OF FIBRONECTIN TYPE III DOMAINS FROM THE CYTOPLASMIC TAIL OF INTEGRIN ALPHA6 BETA4
Authors : de Pereda, J.M.; Wiche, G.; Liddington, R.C.
Deposited on : 1999-04-19
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	195	2% 79% 20% .
1	B	195	4% 77% 21% ...

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CAC	A	383	-	X	-	-
2	CAC	B	382	-	X	-	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3447 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (INTEGRIN BETA-4 SUBUNIT).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	195	Total	C	N	O	S	0	0	0
			1519	955	265	291	8			
1	B	194	Total	C	N	O	S	0	2	1
			1522	956	266	290	10			

- Molecule 2 is CACODYLATE ION (CCD ID: CAC) (formula: $C_2H_6AsO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	As	C	O	0	0
			5	1	2	2		
2	B	1	Total	As	C	O	0	0
			5	1	2	2		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	196	Total	O	0	0
			196	196		
4	B	185	Total	O	0	0
			185	185		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PROTEIN (INTEGRIN BETA-4 SUBUNIT)



• Molecule 1: PROTEIN (INTEGRIN BETA-4 SUBUNIT)



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	59.04Å 59.04Å 246.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.15 20.00 – 2.15	Depositor EDS
% Data completeness (in resolution range)	91.4 (20.00-2.15) 91.7 (20.00-2.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	17.21 (at 2.15Å)	Xtriage
Refinement program	CNS 0.5	Depositor
R, R_{free}	0.201 , 0.262 0.208 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	25.3	Xtriage
Anisotropy	0.210	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 58.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3447	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CAC, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/1559	0.89	0/2122
1	B	0.38	1/1561 (0.1%)	0.85	1/2125 (0.0%)
All	All	0.38	1/3120 (0.0%)	0.87	1/4247 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1319	GLN	C-N	-5.71	1.25	1.34

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1168	GLU	N-CA-C	-5.06	106.80	112.87

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1519	0	1475	43	0
1	B	1522	0	1479	33	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	10	0	0	0	0
3	B	5	0	0	0	0
4	A	196	0	0	2	1
4	B	185	0	0	6	0
All	All	3447	0	2954	76	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1225:ARG:HH11	1:B:1310:ARG:HH22	0.97	0.97
1:A:1193:GLU:HB3	1:A:1210:LEU:HD11	1.53	0.90
1:B:1225:ARG:HH11	1:B:1310:ARG:NH2	1.75	0.83
1:A:1225:ARG:H	1:A:1225:ARG:HD2	1.43	0.82
1:B:1225:ARG:NH1	1:B:1310:ARG:HH22	1.78	0.81
1:A:1261:ASN:HD21	1:A:1265:ARG:HB3	1.48	0.77
1:A:1138:ALA:O	1:A:1142:LYS:HG3	1.87	0.75
1:A:1299:LYS:HE3	1:A:1309:GLU:HG2	1.71	0.72
1:B:1223:PRO:O	1:B:1310:ARG:HD3	1.91	0.70
1:A:1277:ASN:OD1	1:A:1279:LYS:HD3	1.92	0.69
1:B:1261:ASN:HB3	1:B:1267:ILE:HG21	1.74	0.69
1:A:1225:ARG:H	1:A:1225:ARG:CD	2.06	0.69
1:A:1132:ASN:HD22	1:A:1148:LEU:HD22	1.58	0.67
1:A:1225:ARG:HD2	1:A:1225:ARG:N	2.08	0.67
1:B:1230:VAL:HG22	1:B:1316:LEU:HD22	1.76	0.66
1:B:1270[A]:MET:HE1	4:B:251:HOH:O	1.95	0.66
1:B:1319:GLN:HE21	1:B:1319:GLN:HA	1.60	0.65
1:A:1261:ASN:ND2	1:A:1265:ARG:HB3	2.12	0.65
1:B:1132:ASN:HB2	4:B:181:HOH:O	2.03	0.58
1:B:1270[B]:MET:HE2	4:B:268:HOH:O	2.03	0.58
1:B:1284:LEU:CD1	1:B:1286:GLU:HG3	2.34	0.58
1:A:1301:ARG:HD2	4:A:217:HOH:O	2.04	0.56
1:B:1319:GLN:HA	1:B:1319:GLN:NE2	2.19	0.56
1:A:1279:LYS:N	1:A:1279:LYS:HD2	2.20	0.56
1:A:1195:LYS:HB3	1:A:1210:LEU:HD22	1.88	0.55
1:B:1267:ILE:HG13	1:B:1267:ILE:O	2.08	0.54
1:A:1162:TRP:CH2	1:A:1210:LEU:HD21	2.43	0.54
1:A:1155:MET:HE2	1:A:1200:GLY:C	2.34	0.53
1:B:1214:ARG:HD3	4:B:353:HOH:O	2.08	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1295:ARG:HD2	1:A:1311:GLU:OE2	2.09	0.52
1:B:1280:ASN:HB3	4:B:277:HOH:O	2.10	0.52
1:B:1188:PRO:HB3	1:B:1218:GLU:HG2	1.91	0.52
1:B:1194:MET:HG2	1:B:1211:VAL:HG23	1.91	0.51
1:A:1278:PRO:HG2	1:A:1279:LYS:HD2	1.94	0.50
1:A:1193:GLU:HB3	1:A:1210:LEU:CD1	2.34	0.50
1:A:1274:LEU:C	1:A:1274:LEU:HD13	2.37	0.50
1:B:1167:SER:C	1:B:1169:SER:N	2.67	0.49
1:A:1155:MET:HE2	1:A:1200:GLY:HA2	1.95	0.48
1:B:1265:ARG:O	1:B:1267:ILE:HG23	2.13	0.48
1:A:1162:TRP:HH2	1:A:1210:LEU:HD21	1.78	0.48
1:B:1313:ILE:N	1:B:1313:ILE:HD12	2.29	0.48
1:A:1167:SER:C	1:A:1169:SER:N	2.69	0.48
1:B:1194:MET:O	1:B:1210:LEU:HD12	2.14	0.47
1:B:1167:SER:C	1:B:1169:SER:H	2.22	0.47
1:A:1167:SER:C	1:A:1169:SER:H	2.22	0.47
1:A:1225:ARG:CD	1:A:1225:ARG:N	2.73	0.46
1:A:1265:ARG:HH11	1:A:1265:ARG:HG2	1.81	0.46
1:A:1129:ALA:HB2	1:A:1206:PRO:HG2	1.97	0.45
1:A:1155:MET:HE2	1:A:1200:GLY:CA	2.47	0.45
1:B:1254:GLU:OE1	1:B:1299:LYS:HE2	2.17	0.45
1:A:1220:PRO:HG3	1:A:1302:ASN:ND2	2.31	0.45
1:A:1260:VAL:HG11	1:A:1295:ARG:HB2	1.99	0.45
1:A:1279:LYS:HD2	1:A:1279:LYS:H	1.81	0.45
1:A:1274:LEU:HD13	1:A:1275:VAL:N	2.31	0.45
1:B:1189:TYR:CD2	1:B:1248:GLY:HA2	2.53	0.44
1:B:1297:THR:HA	1:B:1310:ARG:O	2.18	0.44
1:A:1256:CYS:SG	1:A:1272:LYS:HG2	2.59	0.43
1:B:1284:LEU:HD13	1:B:1286:GLU:HG3	1.99	0.43
1:B:1295:ARG:HD2	1:B:1311:GLU:OE1	2.19	0.43
1:A:1194:MET:HG2	1:A:1211:VAL:HG23	2.01	0.43
1:B:1129:ALA:HB2	1:B:1206:PRO:HG2	1.99	0.43
1:A:1236:THR:HB	1:A:1316:LEU:HD21	2.01	0.43
1:A:1131:GLN:NE2	4:A:188:HOH:O	2.40	0.42
1:A:1136:LYS:HB2	1:A:1136:LYS:HE3	1.88	0.42
1:A:1279:LYS:N	1:A:1279:LYS:CD	2.82	0.42
1:B:1256[A]:CYS:HB3	1:B:1270[A]:MET:HE3	2.02	0.42
1:A:1301:ARG:HB2	1:A:1306:TRP:CD2	2.55	0.42
1:A:1202:GLN:N	1:A:1202:GLN:CD	2.77	0.42
1:B:1156:GLY:HA3	1:B:1199:TYR:CZ	2.56	0.41
1:B:1225:ARG:CD	1:B:1310:ARG:NH2	2.84	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1265:ARG:HG2	1:A:1265:ARG:NH1	2.36	0.41
1:A:1257:TYR:HA	1:A:1295:ARG:O	2.21	0.41
1:A:1277:ASN:HA	1:A:1278:PRO:HD2	1.82	0.41
1:B:1142:LYS:HG2	1:B:1184:THR:HG22	2.02	0.41
1:B:1127:LEU:N	4:B:305:HOH:O	2.54	0.41
1:A:1132:ASN:ND2	1:A:1148:LEU:HD22	2.30	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:208:HOH:O	4:A:208:HOH:O[8_555]	1.24	0.96

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	193/195 (99%)	189 (98%)	4 (2%)	0	100	100
1	B	194/195 (100%)	188 (97%)	5 (3%)	1 (0%)	24	20
All	All	387/390 (99%)	377 (97%)	9 (2%)	1 (0%)	36	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1319	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	164/165 (99%)	162 (99%)	2 (1%)	63	70
1	B	165/165 (100%)	158 (96%)	7 (4%)	26	25
All	All	329/330 (100%)	320 (97%)	9 (3%)	38	41

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1148	LEU
1	A	1210	LEU
1	B	1151	SER
1	B	1222	GLU
1	B	1225	ARG
1	B	1237	GLN
1	B	1249	GLU
1	B	1265	ARG
1	B	1271	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1132	ASN
1	A	1229	ASN
1	A	1237	GLN
1	A	1280	ASN
1	A	1319	GLN
1	B	1164	GLN
1	B	1237	GLN
1	B	1264	ASN
1	B	1319	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	CAC	A	383	-	2,4,4	3.48	2 (100%)	4,6,6	4.48	4 (100%)
3	SO4	A	384	-	4,4,4	0.35	0	6,6,6	0.07	0
3	SO4	B	386	-	4,4,4	0.41	0	6,6,6	0.05	0
2	CAC	B	382	-	2,4,4	3.48	2 (100%)	4,6,6	4.48	4 (100%)
3	SO4	A	385	-	4,4,4	0.34	0	6,6,6	0.08	0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	382	CAC	AS-C1	3.98	1.99	1.90
2	A	383	CAC	AS-C1	3.96	1.99	1.90
2	A	383	CAC	AS-C2	2.92	1.97	1.90
2	B	382	CAC	AS-C2	2.90	1.97	1.90

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	382	CAC	O1-AS-C1	-7.87	101.72	111.50
2	A	383	CAC	O1-AS-C1	-7.86	101.72	111.50
2	B	382	CAC	O2-AS-C2	2.66	112.29	105.84
2	A	383	CAC	O2-AS-C2	2.65	112.26	105.84
2	A	383	CAC	O2-AS-C1	2.60	112.15	105.84
2	B	382	CAC	O2-AS-C1	2.59	112.13	105.84
2	A	383	CAC	O1-AS-C2	-2.16	108.82	111.50
2	B	382	CAC	O1-AS-C2	-2.14	108.84	111.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	195/195 (100%)	0.18	3 (1%) 72 76	14, 24, 38, 45	0
1	B	194/195 (99%)	0.29	8 (4%) 41 46	10, 23, 47, 54	2 (1%)
All	All	389/390 (99%)	0.24	11 (2%) 55 59	10, 23, 42, 54	2 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1152	GLY	4.8
1	B	1318	THR	3.2
1	A	1280	ASN	2.6
1	B	1320	PRO	2.6
1	A	1201	ALA	2.4
1	B	1166	ASP	2.3
1	A	1126	ASP	2.3
1	B	1270[A]	MET	2.3
1	B	1169	SER	2.3
1	B	1232	SER	2.3
1	B	1167	SER	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	385	5/5	0.93	0.10	42,42,44,45	0
2	CAC	B	382	5/5	0.94	0.20	62,64,66,66	5
3	SO4	B	386	5/5	0.94	0.09	41,42,43,44	0
3	SO4	A	384	5/5	0.96	0.10	38,42,44,44	0
2	CAC	A	383	5/5	0.98	0.19	64,68,68,68	5

6.5 Other polymers [i](#)

There are no such residues in this entry.