



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:13 AM UTC

PDB ID : 1QGH / pdb_00001qgh
Title : THE X-RAY STRUCTURE OF THE UNUSUAL DODECAMERIC FER-
RITIN FROM LISTERIA INNOCUA, REVEALS A NOVEL INTERSUB-
UNIT IRON BINDING SITE.
Authors : Ilari, A.; Stefanini, S.; Chiancone, E.; Tsernoglou, D.
Deposited on : 1999-04-27
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	156	65% 24% 6% . .
1	B	156	64% 24% 8% .
1	C	156	64% 27% 5% .
1	D	156	63% 26% 7% .
1	E	156	64% 28% . .
1	F	156	60% 28% 8% .
1	G	156	67% 21% 7% . .
1	H	156	60% 31% . . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	I	156	56%31%8% . .
1	J	156	62%31% . . .
1	K	156	59%28%8% . .
1	L	156	60%31% . . .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 15155 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NON-HEME IRON-CONTAINING FERRITIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	150	Total	C	N	O	S	32	0	0
			1222	784	196	235	7			
1	B	150	Total	C	N	O	S	57	0	0
			1222	784	196	235	7			
1	C	150	Total	C	N	O	S	44	0	0
			1222	784	196	235	7			
1	D	150	Total	C	N	O	S	47	0	0
			1222	784	196	235	7			
1	E	150	Total	C	N	O	S	46	0	0
			1222	784	196	235	7			
1	F	150	Total	C	N	O	S	42	0	0
			1222	784	196	235	7			
1	G	150	Total	C	N	O	S	41	0	0
			1222	784	196	235	7			
1	H	150	Total	C	N	O	S	34	0	0
			1222	784	196	235	7			
1	I	150	Total	C	N	O	S	45	0	0
			1222	784	196	235	7			
1	J	150	Total	C	N	O	S	44	0	0
			1222	784	196	235	7			
1	K	150	Total	C	N	O	S	43	0	0
			1222	784	196	235	7			
1	L	150	Total	C	N	O	S	50	0	0
			1222	784	196	235	7			

- Molecule 2 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		
2	B	1	Total	Fe	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	1	Total 1	Fe 1	0	0
2	D	1	Total 1	Fe 1	0	0
2	E	1	Total 1	Fe 1	0	0
2	F	1	Total 1	Fe 1	0	0
2	G	1	Total 1	Fe 1	0	0
2	H	1	Total 1	Fe 1	0	0
2	I	1	Total 1	Fe 1	0	0
2	J	1	Total 1	Fe 1	0	0
2	K	1	Total 1	Fe 1	0	0
2	L	1	Total 1	Fe 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	38	Total 38	O 38	0	0
3	B	22	Total 22	O 22	0	0
3	C	36	Total 36	O 36	0	0
3	D	28	Total 28	O 28	0	0
3	E	50	Total 50	O 50	0	0
3	F	42	Total 42	O 42	0	0
3	G	54	Total 54	O 54	0	0
3	H	44	Total 44	O 44	0	0
3	I	33	Total 33	O 33	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	J	46	Total 46	O 46	0	0
3	K	41	Total 41	O 41	0	0
3	L	45	Total 45	O 45	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: NON-HEME IRON-CONTAINING FERRITIN




• Molecule 1: NON-HEME IRON-CONTAINING FERRITIN



• Molecule 1: NON-HEME IRON-CONTAINING FERRITIN




• Molecule 1: NON-HEME IRON-CONTAINING FERRITIN






• Molecule 1: NON-HEME IRON-CONTAINING FERRITIN

Chain E: 64% 28% . .




• Molecule 1: NON-HEME IRON-CONTAINING FERRITIN

Chain F: 60% 28% 8% .




• Molecule 1: NON-HEME IRON-CONTAINING FERRITIN

Chain G: 67% 21% 7% . .



• Molecule 1: NON-HEME IRON-CONTAINING FERRITIN

Chain H: 60% 31% . . .



• Molecule 1: NON-HEME IRON-CONTAINING FERRITIN


Chain I: 56% 31% 8% . .





• Molecule 1: NON-HEME IRON-CONTAINING FERRITIN

Chain J: 62% 31% . . .



• Molecule 1: NON-HEME IRON-CONTAINING FERRITIN

Chain K: 59% 28% 8% . . .



• Molecule 1: NON-HEME IRON-CONTAINING FERRITIN

Chain L: 60% 31% . . .



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	87.54Å 137.50Å 173.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.35	Depositor
% Data completeness (in resolution range)	91.0 (10.00-2.35)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
Refinement program	TNT 5E	Depositor
R, R_{free}	0.219 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	15155	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	1/1249 (0.1%)	1.49	12/1684 (0.7%)
1	B	0.83	1/1249 (0.1%)	1.53	11/1684 (0.7%)
1	C	0.88	2/1249 (0.2%)	1.42	9/1684 (0.5%)
1	D	0.84	1/1249 (0.1%)	1.54	12/1684 (0.7%)
1	E	0.92	1/1249 (0.1%)	1.47	8/1684 (0.5%)
1	F	0.84	1/1249 (0.1%)	1.53	12/1684 (0.7%)
1	G	0.86	0/1249	1.41	6/1684 (0.4%)
1	H	0.90	2/1249 (0.2%)	1.43	9/1684 (0.5%)
1	I	0.86	1/1249 (0.1%)	1.55	20/1684 (1.2%)
1	J	0.86	3/1249 (0.2%)	1.48	9/1684 (0.5%)
1	K	0.84	1/1249 (0.1%)	1.48	11/1684 (0.7%)
1	L	0.92	2/1249 (0.2%)	1.52	15/1684 (0.9%)
All	All	0.87	16/14988 (0.1%)	1.49	134/20208 (0.7%)

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	99	MET	SD-CE	-6.32	1.63	1.79
1	L	118	GLU	C-N	-6.28	1.25	1.33
1	A	8	ASP	N-CA	6.16	1.53	1.45
1	H	8	ASP	N-CA	5.72	1.52	1.46
1	C	32	TRP	NE1-CE2	-5.46	1.31	1.37

The worst 5 of 134 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	156	GLU	CA-C-O	11.68	140.65	120.80
1	E	30	ILE	CA-C-O	11.21	133.05	121.17
1	A	91	LYS	N-CA-C	-10.67	95.36	110.08
1	L	7	VAL	O-C-N	-9.02	108.56	123.00
1	L	156	GLU	CB-CG-CD	8.69	127.38	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1222	0	1186	35	0
1	B	1222	0	1186	37	0
1	C	1222	0	1186	37	0
1	D	1222	0	1186	45	0
1	E	1222	0	1186	41	0
1	F	1222	0	1186	60	0
1	G	1222	0	1186	49	0
1	H	1222	0	1186	41	0
1	I	1222	0	1186	51	0
1	J	1222	0	1186	37	0
1	K	1222	0	1186	72	0
1	L	1222	0	1186	43	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
2	I	1	0	0	0	0
2	J	1	0	0	0	0
2	K	1	0	0	0	0
2	L	1	0	0	0	0
3	A	38	0	0	3	0
3	B	22	0	0	0	0
3	C	36	0	0	3	0
3	D	28	0	0	5	0
3	E	50	0	0	7	0
3	F	42	0	0	7	0
3	G	54	0	0	7	0
3	H	44	0	0	3	0
3	I	33	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	J	46	0	0	6	0
3	K	41	0	0	10	0
3	L	45	0	0	4	0
All	All	15155	0	14232	521	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 521 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:94:THR:HG22	1:G:97:GLN:H	1.11	1.12
1:F:94:THR:HB	1:F:97:GLN:HG3	1.20	1.12
1:F:94:THR:HG22	1:F:96:ASP:N	1.69	1.08
1:F:7:VAL:HG12	1:F:8:ASP:H	1.22	1.04
1:I:94:THR:HB	1:I:97:GLN:HG3	1.40	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	148/156 (95%)	145 (98%)	3 (2%)	0	100	100
1	B	148/156 (95%)	146 (99%)	2 (1%)	0	100	100
1	C	148/156 (95%)	146 (99%)	2 (1%)	0	100	100
1	D	148/156 (95%)	142 (96%)	6 (4%)	0	100	100
1	E	148/156 (95%)	145 (98%)	3 (2%)	0	100	100
1	F	148/156 (95%)	143 (97%)	5 (3%)	0	100	100
1	G	148/156 (95%)	146 (99%)	2 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	148/156 (95%)	146 (99%)	2 (1%)	0	100	100
1	I	148/156 (95%)	145 (98%)	3 (2%)	0	100	100
1	J	148/156 (95%)	145 (98%)	3 (2%)	0	100	100
1	K	148/156 (95%)	144 (97%)	4 (3%)	0	100	100
1	L	148/156 (95%)	143 (97%)	5 (3%)	0	100	100
All	All	1776/1872 (95%)	1736 (98%)	40 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	133/139 (96%)	119 (90%)	14 (10%)	6	6
1	B	133/139 (96%)	122 (92%)	11 (8%)	10	11
1	C	133/139 (96%)	120 (90%)	13 (10%)	7	7
1	D	133/139 (96%)	119 (90%)	14 (10%)	6	6
1	E	133/139 (96%)	120 (90%)	13 (10%)	7	7
1	F	133/139 (96%)	119 (90%)	14 (10%)	6	6
1	G	133/139 (96%)	118 (89%)	15 (11%)	5	5
1	H	133/139 (96%)	118 (89%)	15 (11%)	5	5
1	I	133/139 (96%)	118 (89%)	15 (11%)	5	5
1	J	133/139 (96%)	121 (91%)	12 (9%)	9	9
1	K	133/139 (96%)	118 (89%)	15 (11%)	5	5
1	L	133/139 (96%)	119 (90%)	14 (10%)	6	6
All	All	1596/1668 (96%)	1431 (90%)	165 (10%)	7	6

5 of 165 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	28	HIS
1	K	42	LEU
1	I	65	LEU
1	J	42	LEU
1	K	156	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 58 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	29	GLN
1	L	28	HIS
1	H	129	ASN
1	L	15	HIS
1	J	129	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.