



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:51 PM UTC

PDB ID : 2QG5 / pdb_00002qg5
Title : Cryptosporidium parvum calcium dependent protein kinase cgd7_1840
Authors : Lunin, V.V.; Wernimont, A.K.; Lew, J.; Wasney, G.; Kozieradzki, I.; Vedadi, M.; Bochkarev, A.; Arrowsmith, C.H.; Sundstrom, M.; Weigelt, J.; Edwards, A.E.; Hui, R.; Artz, J.; Amani, M.; Structural Genomics Consortium (SGC)
Deposited on : 2007-06-28
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	294	2% 77% 17% • •
1	B	294	5% 73% 19% • •
1	D	294	3% 74% 18% • •

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7227 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Calcium/calmodulin-dependent protein kinase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	285	Total	C	N	O	P	S	0	5	0
			2365	1509	412	431	1	12			
1	B	281	Total	C	N	O	P	S	0	4	0
			2334	1490	407	424	1	12			
1	D	284	Total	C	N	O	P	S	0	2	0
			2337	1492	406	426	1	12			

There are 57 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP Q5CYL9
A	2	HIS	-	expression tag	UNP Q5CYL9
A	3	HIS	-	expression tag	UNP Q5CYL9
A	4	HIS	-	expression tag	UNP Q5CYL9
A	5	HIS	-	expression tag	UNP Q5CYL9
A	6	HIS	-	expression tag	UNP Q5CYL9
A	7	HIS	-	expression tag	UNP Q5CYL9
A	8	SER	-	expression tag	UNP Q5CYL9
A	9	SER	-	expression tag	UNP Q5CYL9
A	10	GLY	-	expression tag	UNP Q5CYL9
A	11	ARG	-	expression tag	UNP Q5CYL9
A	12	GLU	-	expression tag	UNP Q5CYL9
A	13	ASN	-	expression tag	UNP Q5CYL9
A	14	LEU	-	expression tag	UNP Q5CYL9
A	15	TYR	-	expression tag	UNP Q5CYL9
A	16	PHE	-	expression tag	UNP Q5CYL9
A	17	GLN	-	expression tag	UNP Q5CYL9
A	18	GLY	-	expression tag	UNP Q5CYL9
A	230	SEP	SER	modified residue	UNP Q5CYL9
B	1	MET	-	expression tag	UNP Q5CYL9
B	2	HIS	-	expression tag	UNP Q5CYL9
B	3	HIS	-	expression tag	UNP Q5CYL9
B	4	HIS	-	expression tag	UNP Q5CYL9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	5	HIS	-	expression tag	UNP Q5CYL9
B	6	HIS	-	expression tag	UNP Q5CYL9
B	7	HIS	-	expression tag	UNP Q5CYL9
B	8	SER	-	expression tag	UNP Q5CYL9
B	9	SER	-	expression tag	UNP Q5CYL9
B	10	GLY	-	expression tag	UNP Q5CYL9
B	11	ARG	-	expression tag	UNP Q5CYL9
B	12	GLU	-	expression tag	UNP Q5CYL9
B	13	ASN	-	expression tag	UNP Q5CYL9
B	14	LEU	-	expression tag	UNP Q5CYL9
B	15	TYR	-	expression tag	UNP Q5CYL9
B	16	PHE	-	expression tag	UNP Q5CYL9
B	17	GLN	-	expression tag	UNP Q5CYL9
B	18	GLY	-	expression tag	UNP Q5CYL9
B	230	SEP	SER	modified residue	UNP Q5CYL9
D	1	MET	-	expression tag	UNP Q5CYL9
D	2	HIS	-	expression tag	UNP Q5CYL9
D	3	HIS	-	expression tag	UNP Q5CYL9
D	4	HIS	-	expression tag	UNP Q5CYL9
D	5	HIS	-	expression tag	UNP Q5CYL9
D	6	HIS	-	expression tag	UNP Q5CYL9
D	7	HIS	-	expression tag	UNP Q5CYL9
D	8	SER	-	expression tag	UNP Q5CYL9
D	9	SER	-	expression tag	UNP Q5CYL9
D	10	GLY	-	expression tag	UNP Q5CYL9
D	11	ARG	-	expression tag	UNP Q5CYL9
D	12	GLU	-	expression tag	UNP Q5CYL9
D	13	ASN	-	expression tag	UNP Q5CYL9
D	14	LEU	-	expression tag	UNP Q5CYL9
D	15	TYR	-	expression tag	UNP Q5CYL9
D	16	PHE	-	expression tag	UNP Q5CYL9
D	17	GLN	-	expression tag	UNP Q5CYL9
D	18	GLY	-	expression tag	UNP Q5CYL9
D	230	SEP	SER	modified residue	UNP Q5CYL9

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	81	Total O 81 81	0	0
2	B	47	Total O 47 47	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	63	Total	O	0	0
			63	63		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Calcium/calmodulin-dependent protein kinase



- Molecule 1: Calcium/calmodulin-dependent protein kinase



- Molecule 1: Calcium/calmodulin-dependent protein kinase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	138.18Å 86.48Å 87.97Å 90.00° 96.37° 90.00°	Depositor
Resolution (Å)	38.87 – 2.30 38.87 – 2.30	Depositor EDS
% Data completeness (in resolution range)	88.4 (38.87-2.30) 88.4 (38.87-2.30)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.220 , 0.286 (Not available) , 0.256	Depositor DCC
R_{free} test set	2104 reflections (4.59%)	wwPDB-VP
Wilson B-factor (Å ²)	37.6	Xtriage
Anisotropy	0.274	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 45.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7227	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	0/2413	0.99	4/3256 (0.1%)
1	B	0.73	1/2382 (0.0%)	0.94	3/3215 (0.1%)
1	D	0.78	0/2386	0.96	7/3221 (0.2%)
All	All	0.77	1/7181 (0.0%)	0.96	14/9692 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	D	0	2
All	All	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	64	VAL	CA-CB	5.14	1.59	1.53

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	168	ILE	N-CA-C	8.51	118.80	111.90
1	A	8	SER	N-CA-C	7.61	121.05	111.69
1	B	281	TRP	N-CA-C	-7.21	105.01	113.88
1	A	10	GLY	N-CA-C	7.21	125.18	112.64
1	D	168	ILE	N-CA-C	6.71	120.10	113.53
1	D	177	LYS	CA-C-N	6.67	127.21	119.47
1	D	177	LYS	C-N-CA	6.67	127.21	119.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	202	GLY	CA-C-N	6.25	126.72	119.47
1	D	202	GLY	C-N-CA	6.25	126.72	119.47
1	B	25	ASN	N-CA-C	5.74	123.03	110.80
1	D	188	THR	CA-C-N	5.42	126.61	119.84
1	D	188	THR	C-N-CA	5.42	126.61	119.84
1	B	64	VAL	CB-CA-C	5.20	116.85	111.23
1	A	94	ASP	N-CA-C	-5.06	102.40	109.69

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	9	SER	Peptide
1	D	6	HIS	Peptide
1	D	9	SER	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2365	0	2341	27	0
1	B	2334	0	2302	36	0
1	D	2337	0	2311	33	0
2	A	81	0	0	0	0
2	B	47	0	0	0	0
2	D	63	0	0	1	0
All	All	7227	0	6954	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261[A]:ARG:HH21	1:A:261[A]:ARG:HG2	1.28	0.95

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:136:VAL:HG11	1:D:149:LEU:HD11	1.66	0.77
1:D:53:ARG:HD2	2:D:319:HOH:O	1.86	0.74
1:B:87[A]:ARG:HH21	1:B:87[A]:ARG:CG	2.00	0.74
1:D:11:ARG:HA	1:D:14:LEU:HG	1.69	0.73
1:D:75:GLU:HG3	1:D:78:LYS:HE3	1.72	0.71
1:B:82:HIS:CD2	1:B:84:ASN:H	2.09	0.70
1:B:136:VAL:HG11	1:B:149:LEU:HD11	1.71	0.70
1:D:192:VAL:HG13	1:D:196:VAL:HB	1.74	0.70
1:D:195:GLN:NE2	1:D:265:LYS:O	2.25	0.69
1:A:192:VAL:HG13	1:A:196:VAL:HB	1.75	0.68
1:D:204:GLU:HG3	1:D:267:PRO:HB3	1.76	0.67
1:B:117:LYS:O	1:B:118:ARG:HB2	1.94	0.65
1:A:261[A]:ARG:HH21	1:A:261[A]:ARG:CG	2.08	0.63
1:B:42:VAL:HG22	1:B:57:LYS:HG2	1.80	0.62
1:A:189:PRO:HB3	1:A:233:MET:HE3	1.82	0.61
1:D:36:ARG:HG3	1:D:41:GLU:HB3	1.82	0.61
1:D:93:GLU:HB3	1:D:98:ILE:HG12	1.84	0.59
1:B:30:LEU:HB3	1:B:43:LYS:HD2	1.84	0.59
1:B:87[A]:ARG:HH21	1:B:87[A]:ARG:HG2	1.68	0.59
1:A:261[A]:ARG:HG2	1:A:261[A]:ARG:NH2	2.07	0.58
1:B:82:HIS:HD2	1:B:84:ASN:H	1.48	0.58
1:B:24:ILE:HG12	1:B:30:LEU:HD11	1.84	0.58
1:B:216:VAL:HG12	1:B:222:PRO:HA	1.85	0.58
1:A:32:ASN:H	1:A:32:ASN:HD22	1.50	0.57
1:B:222:PRO:HG2	1:B:225:SER:HB3	1.84	0.57
1:B:95:ASN:HD22	1:B:96:THR:HG23	1.69	0.57
1:B:157:LEU:HG	1:B:166:LYS:HE3	1.88	0.56
1:A:282:PHE:O	1:A:286:LEU:HB2	2.07	0.55
1:A:132:VAL:HG11	1:A:213:MET:CE	2.37	0.55
1:B:12:GLU:HB2	1:B:14:LEU:HD23	1.89	0.54
1:B:5:HIS:HB2	1:B:92:PHE:CD2	2.43	0.54
1:D:72:GLN:O	1:D:76:ILE:HG12	2.09	0.53
1:B:4:HIS:CE1	1:B:16:PHE:HB2	2.45	0.52
1:D:110:LEU:HD23	1:D:151:PRO:HB3	1.92	0.52
1:B:72:GLN:HG3	1:B:175[B]:ARG:NH2	2.24	0.51
1:B:6:HIS:HB3	1:B:19:SER:OG	2.11	0.50
1:A:117:LYS:O	1:A:118:ARG:HG2	2.12	0.50
1:A:237:ARG:O	1:A:265:LYS:HE3	2.11	0.50
1:B:87[A]:ARG:HH21	1:B:87[A]:ARG:HG3	1.76	0.50
1:A:32:ASN:H	1:A:32:ASN:ND2	2.09	0.50
1:B:87[A]:ARG:HG2	1:B:87[A]:ARG:NH2	2.24	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:31:GLU:HB3	1:D:44:ILE:HB	1.93	0.50
1:A:7:HIS:ND1	1:A:8:SER:N	2.59	0.49
1:B:245:GLU:O	1:B:249:LEU:HB2	2.13	0.49
1:D:214:MET:HE1	1:D:258:LEU:HD23	1.94	0.49
1:D:279[B]:HIS:HD2	1:D:281:TRP:H	1.59	0.49
1:D:58:LYS:HE2	1:D:99:TYR:OH	2.13	0.48
1:A:132:VAL:HG11	1:A:213:MET:HE3	1.94	0.48
1:A:9:SER:CB	1:A:95:ASN:HD21	2.27	0.48
1:D:4:HIS:O	1:D:19:SER:HB2	2.13	0.48
1:D:172:LEU:HD13	1:D:186:VAL:HG13	1.95	0.48
1:B:177:LYS:HD2	1:B:180:LYS:HD2	1.95	0.47
1:A:82:HIS:CD2	1:A:84:ASN:H	2.32	0.47
1:A:129:MET:HG3	1:A:214:MET:HG3	1.95	0.47
1:D:250:ASN:H	1:D:250:ASN:HD22	1.63	0.47
1:D:7:HIS:ND1	1:D:8:SER:N	2.63	0.47
1:A:177[A]:LYS:HB3	1:A:180:LYS:HB2	1.97	0.46
1:B:87[A]:ARG:CG	1:B:87[A]:ARG:NH2	2.67	0.46
1:D:70:PHE:CE2	1:D:74:ILE:HD11	2.51	0.46
1:B:229:ASP:O	1:B:233:MET:HG2	2.16	0.45
1:B:76:ILE:O	1:B:80:LEU:HG	2.16	0.45
1:A:5:HIS:HB3	1:A:22:GLY:HA3	1.99	0.45
1:A:75:GLU:HG3	1:A:78:LYS:HE2	1.98	0.45
1:D:91:THR:HG23	1:D:98:ILE:HD11	1.99	0.45
1:A:9:SER:HB3	1:A:95:ASN:HD21	1.81	0.45
1:B:81:ASP:OD1	1:B:87[A]:ARG:NH2	2.49	0.45
1:B:72:GLN:HG3	1:B:175[B]:ARG:HH22	1.82	0.45
1:D:267:PRO:HD2	1:D:268:LYS:HE3	1.99	0.44
1:A:261[B]:ARG:HD2	1:A:271:ILE:HD12	1.99	0.44
1:A:65:GLU:HG3	1:A:66:ASP:N	2.33	0.44
1:A:117:LYS:HD3	1:A:119:VAL:O	2.18	0.44
1:B:87[B]:ARG:HA	1:B:87[B]:ARG:HD2	1.57	0.44
1:A:192:VAL:HG12	1:A:197:LEU:HD13	1.99	0.44
1:D:230:SEP:HA	1:D:233:MET:HE3	1.98	0.44
1:D:75:GLU:HA	1:D:78:LYS:HG2	1.99	0.43
1:D:195:GLN:NE2	1:D:267:PRO:HD3	2.33	0.43
1:D:268:LYS:HD2	1:D:268:LYS:H	1.82	0.43
1:D:214:MET:HE1	1:D:258:LEU:CD2	2.49	0.43
1:A:155:LEU:HD22	1:A:168:ILE:HG21	2.00	0.43
1:A:118:ARG:HG3	1:A:119:VAL:HG23	2.00	0.43
1:D:128:ILE:HG23	1:D:165:LEU:HD22	1.99	0.42
1:D:279[B]:HIS:CD2	1:D:281:TRP:H	2.37	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:86:ILE:HD12	1:B:102:MET:HG3	2.00	0.42
1:B:193:SER:HB3	1:B:196:VAL:HG23	2.02	0.42
1:B:200:LEU:CD1	1:D:268:LYS:HD3	2.50	0.42
1:D:86:ILE:HB	1:D:168:ILE:HG22	2.02	0.41
1:B:92:PHE:HB2	1:B:99:TYR:HB2	2.02	0.41
1:B:212:VAL:O	1:B:216:VAL:HG13	2.19	0.41
1:B:197:LEU:HD23	1:B:237:ARG:HG3	2.03	0.41
1:B:197:LEU:HD11	1:B:233:MET:SD	2.60	0.41
1:D:127:ARG:NH2	1:D:161:PRO:O	2.52	0.40
1:A:24:ILE:HD13	1:A:30:LEU:HD11	2.04	0.40
1:D:176:PHE:CZ	1:D:203:PRO:HG3	2.56	0.40
1:B:242:THR:O	1:B:244:PRO:HD3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	287/294 (98%)	276 (96%)	11 (4%)	0	100	100
1	B	280/294 (95%)	262 (94%)	16 (6%)	2 (1%)	18	23
1	D	283/294 (96%)	267 (94%)	16 (6%)	0	100	100
All	All	850/882 (96%)	805 (95%)	43 (5%)	2 (0%)	43	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	25	ASN
1	B	118	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	256/262 (98%)	229 (90%)	27 (10%)	6	8
1	B	253/262 (97%)	227 (90%)	26 (10%)	7	8
1	D	254/262 (97%)	232 (91%)	22 (9%)	9	12
All	All	763/786 (97%)	688 (90%)	75 (10%)	8	10

All (75) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LEU
1	A	20	THR
1	A	24	ILE
1	A	32	ASN
1	A	64	VAL
1	A	72	GLN
1	A	75	GLU
1	A	117	LYS
1	A	136	VAL
1	A	149	LEU
1	A	150	LYS
1	A	157	LEU
1	A	159	ASP
1	A	184	THR
1	A	186	VAL
1	A	192	VAL
1	A	197	LEU
1	A	200	LEU
1	A	232	VAL
1	A	235	LYS
1	A	246	LYS
1	A	249	LEU
1	A	260	ARG
1	A	261[A]	ARG
1	A	261[B]	ARG
1	A	263	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	273	SER
1	B	3	HIS
1	B	13	ASN
1	B	14	LEU
1	B	25	ASN
1	B	41	GLU
1	B	47	GLN
1	B	64	VAL
1	B	86	ILE
1	B	87[A]	ARG
1	B	87[B]	ARG
1	B	95	ASN
1	B	98	ILE
1	B	102	MET
1	B	109	GLU
1	B	113	ARG
1	B	157	LEU
1	B	197	LEU
1	B	198	GLU
1	B	200	LEU
1	B	216	VAL
1	B	238	GLU
1	B	240	THR
1	B	263	LEU
1	B	278	GLU
1	B	280	GLU
1	B	286	LEU
1	D	11	ARG
1	D	12	GLU
1	D	17	GLN
1	D	20	THR
1	D	75	GLU
1	D	87	ARG
1	D	91	THR
1	D	98	ILE
1	D	121	ARG
1	D	157	LEU
1	D	177	LYS
1	D	186	VAL
1	D	192	VAL
1	D	195	GLN
1	D	197	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	204	GLU
1	D	250	ASN
1	D	260	ARG
1	D	263	LEU
1	D	268	LYS
1	D	278	GLU
1	D	285	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	HIS
1	A	5	HIS
1	A	13	ASN
1	A	32	ASN
1	A	47	GLN
1	A	82	HIS
1	A	84	ASN
1	A	95	ASN
1	A	116	HIS
1	A	153	ASN
1	B	4	HIS
1	B	82	HIS
1	B	95	ASN
1	D	13	ASN
1	D	82	HIS
1	D	84	ASN
1	D	250	ASN
1	D	275	GLN
1	D	285	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SEP	A	230	1	8,9,10	1.83	3 (37%)	7,12,14	1.24	1 (14%)
1	SEP	B	230	1	8,9,10	1.73	2 (25%)	7,12,14	1.13	1 (14%)
1	SEP	D	230	1	8,9,10	1.84	2 (25%)	7,12,14	1.24	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	230	1	-	1/6/8/10	-
1	SEP	B	230	1	-	0/6/8/10	-
1	SEP	D	230	1	-	0/6/8/10	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	230	SEP	P-O1P	4.05	1.63	1.50
1	A	230	SEP	P-O1P	3.89	1.62	1.50
1	B	230	SEP	P-O1P	3.70	1.62	1.50
1	A	230	SEP	P-O3P	2.14	1.62	1.54
1	A	230	SEP	P-O2P	2.08	1.62	1.54
1	B	230	SEP	P-O2P	2.04	1.62	1.54
1	D	230	SEP	P-O2P	2.03	1.62	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	230	SEP	OG-CB-CA	2.62	110.69	108.14
1	D	230	SEP	OG-CB-CA	2.61	110.68	108.14
1	A	230	SEP	OG-CB-CA	2.16	110.24	108.14

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	230	SEP	CB-OG-P-O1P

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	230	SEP	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	284/294 (96%)	0.15	5 (1%) 67 69	11, 32, 57, 68	5 (1%)
1	B	280/294 (95%)	0.47	14 (5%) 34 35	12, 38, 68, 81	4 (1%)
1	D	283/294 (96%)	0.30	10 (3%) 47 49	15, 35, 58, 73	2 (0%)
All	All	847/882 (96%)	0.31	29 (3%) 48 50	11, 35, 62, 81	11 (1%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	25	ASN	4.0
1	B	7	HIS	3.8
1	D	10	GLY	3.7
1	A	229	ASP	3.2
1	B	179	GLY	2.9
1	A	24	ILE	2.9
1	B	33	THR	2.8
1	B	34	ILE	2.8
1	A	10	GLY	2.8
1	D	179	GLY	2.8
1	D	160	SER	2.7
1	A	9	SER	2.7
1	D	9	SER	2.6
1	B	32	ASN	2.6
1	B	19	SER	2.6
1	B	3	HIS	2.5
1	D	68	ASP	2.5
1	D	239	GLY	2.4
1	D	249	LEU	2.4
1	B	6	HIS	2.3
1	B	31	GLU	2.3
1	D	8	SER	2.3
1	D	31	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	46	VAL	2.2
1	A	177[A]	LYS	2.1
1	B	39	TRP	2.1
1	B	178	PRO	2.1
1	D	18	GLY	2.1
1	B	24	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	A	230	10/11	0.85	0.12	31,34,48,49	0
1	SEP	D	230	10/11	0.86	0.10	33,37,54,55	0
1	SEP	B	230	10/11	0.89	0.08	55,57,61,62	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.