



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:23 AM UTC

PDB ID : 3QGE / pdb_00003qge

Title : Crystal structure of the hepatitis C virus NS5B RNA-dependent RNA polymerase complex with (2E)-3-(4-{{(1-{{(13-cyclohexyl-6-oxo-6,7-dihydro-5H-indolo[1,2-d][1,4]benzodiazepin-10-yl)carbonyl}amino}cyclopentyl)carbonyl}amino}phenyl)prop-2-enoic acid and (2R)-4-(2,6-dimethoxypyrimidin-4-yl)-N-(4-methoxybenzyl)-1-{{[4-(trifluoromethoxy)phenyl]sulfonyl}piperazine-2-carboxamide

Authors : Sheriff, S.

Deposited on : 2011-01-24

Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	574	
1	B	574	

Ideal geometry (proteins) : Engh & Huber (2001)
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8489 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called RNA-directed RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	541	Total	C	N	O	S	0	0	0
			4187	2640	739	776	32			
1	B	531	Total	C	N	O	S	0	0	0
			4107	2591	727	757	32			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP Q9WMX2
B	0	MET	-	initiating methionine	UNP Q9WMX2

- Molecule 2 is (2E)-3-(4-(((1-(((13-cyclohexyl-6-oxo-6,7-dihydro-5H-indolo[1,2-d][1,4]benzodiazepin-10-yl)carbonyl]amino}cyclopentyl)carbonyl]amino}phenyl)prop-2-enoic acid (CCD ID: 23E) (formula: C₃₈H₃₈N₄O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			47	38	4	5		
2	B	1	Total	C	N	O	0	0
			47	38	4	5		


- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is (2R)-4-(2,6-dimethoxypyrimidin-4-yl)-N-(4-methoxybenzyl)-1-{[4-(tr

ifluoromethoxy)phenyl)sulfonyl}piperazine-2-carboxamide (CCD ID: 26F) (formula: $C_{26}H_{28}F_3N_5O_7S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	B	1	Total	C	F	N	O	S	0	0
			42	26	3	5	7	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	4	Total	O	0	0
			4	4		
5	B	10	Total	O	0	0
			10	10		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA-directed RNA polymerase



• Molecule 1: RNA-directed RNA polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.70Å 91.40Å 232.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.61 – 3.00 31.61 – 3.00	Depositor EDS
% Data completeness (in resolution range)	90.4 (31.61-3.00) 90.3 (31.61-3.00)	Depositor EDS
R_{merge}	0.23	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.25 (at 3.00Å)	Xtriage
Refinement program	BUSTER-TNT BUSTER 2.9.7, BUSTER 2.9.7	Depositor
R, R_{free}	0.236 , 0.272 0.237 , 0.265	Depositor DCC
R_{free} test set	981 reflections (3.70%)	wwPDB-VP
Wilson B-factor (Å ²)	40.8	Xtriage
Anisotropy	0.556	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 48.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	8489	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 26F, 23E, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	1/4278 (0.0%)	1.25	10/5806 (0.2%)
1	B	0.84	1/4195 (0.0%)	1.25	8/5693 (0.1%)
All	All	0.83	2/8473 (0.0%)	1.25	18/11499 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	173	MET	SD-CE	-8.28	1.58	1.79
1	A	173	MET	SD-CE	-7.42	1.60	1.79

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	261	TYR	N-CA-C	6.44	117.96	111.07
1	B	244	ASP	N-CA-C	-6.13	99.87	109.25
1	A	244	ASP	N-CA-C	-5.97	100.12	109.25
1	A	40	THR	CB-CA-C	5.97	119.56	109.84
1	A	530	VAL	CA-C-N	5.95	128.17	120.44
1	A	530	VAL	C-N-CA	5.95	128.17	120.44
1	B	530	VAL	N-CA-C	5.92	117.73	109.55
1	B	546	ASP	N-CA-C	5.82	118.14	109.59
1	B	535	LYS	N-CA-C	5.66	117.12	111.07
1	A	552	VAL	N-CA-C	-5.49	106.32	111.48
1	A	175	LEU	N-CA-C	5.43	121.75	113.61
1	A	530	VAL	N-CA-C	5.42	117.03	109.55
1	B	175	LEU	N-CA-C	5.29	121.55	113.61
1	A	78	VAL	N-CA-C	5.18	115.72	108.89
1	A	559	ASP	CA-CB-CG	5.14	117.74	112.60
1	B	310	ASP	CA-CB-CG	5.13	117.73	112.60
1	B	545	LEU	N-CA-C	5.06	117.00	108.96

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	361	GLU	CB-CG-CD	5.02	121.13	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4187	0	4181	24	0
1	B	4107	0	4100	23	0
2	A	47	0	37	0	0
2	B	47	0	37	1	0
3	A	20	0	0	0	0
3	B	25	0	0	0	0
4	B	42	0	28	4	0
5	A	4	0	0	0	0
5	B	10	0	0	0	0
All	All	8489	0	8383	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:170:CYS:HA	1:A:173:MET:HE3	1.46	0.97
1:B:170:CYS:HA	1:B:173:MET:HE3	1.45	0.95
1:A:247:PRO:HG3	1:B:234:ARG:NH1	1.94	0.82
1:B:398:GLU:HG2	1:B:403:THR:HG21	1.60	0.82
1:B:447:ILE:HD13	4:B:602:26F:H6	1.63	0.78
1:B:234:ARG:NH1	1:B:258:GLU:OE2	2.17	0.78
1:A:40:THR:HG22	1:A:143:GLU:H	1.54	0.73
1:A:234:ARG:NH1	1:A:258:GLU:OE2	2.24	0.70
1:A:172:LYS:HE3	1:A:560:ILE:HD13	1.79	0.63
1:A:308:LEU:HB3	1:A:311:CYS:SG	2.41	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:197:PRO:O	1:A:201:VAL:HG23	2.01	0.61
1:B:197:PRO:O	1:B:201:VAL:HG23	2.02	0.60
1:A:563:SER:O	1:A:564:LEU:HG	2.01	0.60
1:B:308:LEU:HB3	1:B:311:CYS:SG	2.41	0.60
1:A:314:LEU:HB3	1:A:321:VAL:HG23	1.87	0.57
1:B:234:ARG:HG3	1:B:262:ILE:HD11	1.86	0.57
1:A:219:TYR:HB3	1:A:320:LEU:HD23	1.87	0.56
1:B:219:TYR:HB3	1:B:320:LEU:HD23	1.87	0.56
1:B:314:LEU:HB3	1:B:321:VAL:HG23	1.89	0.54
1:B:447:ILE:CD1	4:B:602:26F:H6	2.34	0.53
1:B:361:GLU:HG3	1:B:370:VAL:O	2.10	0.52
1:A:361:GLU:HG3	1:A:370:VAL:O	2.10	0.51
1:B:234:ARG:HH12	1:B:258:GLU:CD	2.18	0.51
1:A:448:TYR:CE2	1:A:551:PHE:HD1	2.28	0.51
1:A:446:GLN:O	1:A:447:ILE:HD13	2.13	0.49
1:B:405:VAL:HG13	1:B:445:CYS:HA	1.95	0.49
1:A:230:GLU:HG2	1:A:262:ILE:HG23	1.96	0.48
1:B:197:PRO:HA	4:B:602:26F:H4	1.95	0.47
1:A:82:LEU:HD13	1:A:249:ALA:HB2	1.97	0.47
1:A:40:THR:CG2	1:A:141:LYS:O	2.63	0.46
1:B:309:GLN:O	1:B:324:CYS:HB2	2.16	0.45
1:A:268:ASN:HB3	1:A:274:CYS:SG	2.56	0.45
1:B:440:GLU:H	1:B:440:GLU:CD	2.25	0.44
1:B:82:LEU:HD13	1:B:249:ALA:HB2	1.99	0.44
1:B:531:ARG:HE	1:B:531:ARG:HB2	1.57	0.44
2:B:601:23E:H37A	2:B:601:23E:H24	2.00	0.44
1:A:440:GLU:H	1:A:440:GLU:CD	2.27	0.43
1:A:201:VAL:HG22	1:A:384:LEU:HD13	2.01	0.43
1:A:182:LEU:HD12	1:A:243:CYS:SG	2.58	0.42
1:A:247:PRO:HG3	1:B:234:ARG:HH11	1.82	0.42
1:B:447:ILE:HD13	4:B:602:26F:C6	2.42	0.42
1:B:201:VAL:HG22	1:B:384:LEU:HD13	2.01	0.42
1:A:40:THR:HG23	1:A:141:LYS:O	2.20	0.42
1:A:141:LYS:HG2	1:A:143:GLU:HG3	2.03	0.41
1:B:309:GLN:HB2	1:B:325:GLU:HB3	2.02	0.41
1:A:234:ARG:HH12	1:A:258:GLU:CD	2.30	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	537/574 (94%)	526 (98%)	8 (2%)	3 (1%)	21	56
1	B	525/574 (92%)	507 (97%)	15 (3%)	3 (1%)	21	56
All	All	1062/1148 (92%)	1033 (97%)	23 (2%)	6 (1%)	21	56

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	550	TRP
1	A	147	VAL
1	B	147	VAL
1	B	544	GLN
1	A	536	LEU
1	B	536	LEU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	451/486 (93%)	428 (95%)	23 (5%)	21	55
1	B	440/486 (90%)	419 (95%)	21 (5%)	23	57
All	All	891/972 (92%)	847 (95%)	44 (5%)	22	56

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	THR
1	A	43	ARG
1	A	46	SER
1	A	47	LEU
1	A	86	GLU
1	A	90	LYS
1	A	151	LYS
1	A	159	LEU
1	A	262	ILE
1	A	278	ARG
1	A	307	LYS
1	A	308	LEU
1	A	321	VAL
1	A	336	LEU
1	A	361	GLU
1	A	372	VAL
1	A	380	ARG
1	A	399	THR
1	A	440	GLU
1	A	535	LYS
1	A	547	LEU
1	A	556	SER
1	A	563	SER
1	B	46	SER
1	B	47	LEU
1	B	86	GLU
1	B	100	LYS
1	B	151	LYS
1	B	159	LEU
1	B	160	ILE
1	B	262	ILE
1	B	308	LEU
1	B	321	VAL
1	B	336	LEU
1	B	361	GLU
1	B	379	LYS
1	B	399	THR
1	B	403	THR
1	B	440	GLU
1	B	469	LEU
1	B	531	ARG
1	B	535	LYS
1	B	547	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	555	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	142	ASN
1	A	374	HIS
1	B	142	ASN
1	B	402	HIS
1	B	446	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	658	-	4,4,4	0.34	0	6,6,6	0.30	0
2	23E	B	601	-	52,53,53	1.60	7 (13%)	68,77,77	1.27	10 (14%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	653	-	4,4,4	0.55	0	6,6,6	0.16	0
3	SO4	B	653	-	4,4,4	0.55	0	6,6,6	0.09	0
3	SO4	A	654	-	4,4,4	0.29	0	6,6,6	0.07	0
3	SO4	B	656	-	4,4,4	0.60	0	6,6,6	0.32	0
2	23E	A	601	-	52,53,53	1.64	6 (11%)	68,77,77	1.25	7 (10%)
3	SO4	B	657	-	4,4,4	0.80	0	6,6,6	0.30	0
3	SO4	A	651	-	4,4,4	0.50	0	6,6,6	0.28	0
3	SO4	B	652	-	4,4,4	0.42	0	6,6,6	0.19	0
4	26F	B	602	-	45,45,45	1.82	9 (20%)	61,65,65	1.47	6 (9%)
3	SO4	A	652	-	4,4,4	0.37	0	6,6,6	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	23E	A	601	-	-	0/28/57/57	0/6/7/7
2	23E	B	601	-	-	2/28/57/57	0/6/7/7
4	26F	B	602	-	-	2/36/49/49	0/4/4/4

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	602	26F	S42-N30	6.11	1.72	1.63
2	B	601	23E	O15-C14	5.32	1.36	1.23
2	A	601	23E	O15-C14	5.14	1.35	1.23
2	A	601	23E	C30-N26	-4.78	1.31	1.39
2	B	601	23E	C32-C30	4.42	1.51	1.47
4	B	602	26F	O33-S42	4.39	1.48	1.43
4	B	602	26F	O34-S42	3.99	1.47	1.43
2	A	601	23E	C28-C29	3.96	1.50	1.44
2	B	601	23E	C30-N26	-3.85	1.33	1.39
2	A	601	23E	C32-C30	3.46	1.50	1.47
2	B	601	23E	C28-C29	3.24	1.49	1.44
4	B	602	26F	C15-N28	2.84	1.37	1.33
4	B	602	26F	C3-C11	2.78	1.43	1.38
2	B	601	23E	O20-C14	-2.68	1.23	1.30
2	A	601	23E	O20-C14	-2.61	1.23	1.30
4	B	602	26F	C9-C15	2.55	1.42	1.38
2	B	601	23E	C27-N26	-2.54	1.35	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	602	26F	C5-C12	2.45	1.43	1.38
2	A	601	23E	C28-C27	-2.45	1.37	1.41
4	B	602	26F	C4-C11	2.24	1.42	1.38
4	B	602	26F	C9-C14	2.24	1.42	1.39
2	B	601	23E	C9-C12	2.16	1.53	1.47

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	602	26F	C25-N31-C17	5.41	129.73	122.29
4	B	602	26F	C18-C19-N30	4.57	113.99	109.00
4	B	602	26F	C21-C20-N29	3.70	114.92	109.36
2	A	601	23E	C27-N26-C30	3.50	111.17	108.37
2	A	601	23E	C27-C28-C29	-3.48	104.09	107.02
2	A	601	23E	C24-C28-C27	3.11	122.83	119.24
2	B	601	23E	O6-C3-C2	-3.01	116.48	120.99
2	B	601	23E	C23-C24-C28	-2.94	116.01	120.86
2	B	601	23E	C27-N26-C30	2.92	110.71	108.37
2	B	601	23E	C27-C28-C29	-2.89	104.59	107.02
2	B	601	23E	C24-C28-C27	2.75	122.41	119.24
4	B	602	26F	O34-S42-C13	-2.42	105.10	108.10
2	A	601	23E	O6-C3-C2	-2.40	117.40	120.99
2	B	601	23E	O47-C46-N45	-2.39	118.77	121.10
2	A	601	23E	C23-C24-C28	-2.38	116.93	120.86
2	A	601	23E	C28-C29-C31	-2.35	121.01	126.87
2	B	601	23E	C34-C33-C31	-2.32	107.01	111.24
4	B	602	26F	C16-N28-C15	2.26	116.00	114.04
4	B	602	26F	C9-C14-N27	-2.25	119.01	123.34
2	A	601	23E	C17-C2-C16	-2.11	100.74	103.39
2	B	601	23E	C28-C29-C30	-2.03	105.32	107.28
2	B	601	23E	C44-C46-N45	2.01	123.00	117.55
2	B	601	23E	C35-C34-C33	-2.00	107.30	111.42

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	601	23E	C12-C13-C14-O20
2	B	601	23E	C12-C13-C14-O15
4	B	602	26F	C6-C12-O36-C26
4	B	602	26F	C5-C12-O36-C26

There are no ring outliers.


2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	23E	1	0
4	B	602	26F	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



Ligand 23E A 601



Ligand 26F B 602



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	541/574 (94%)	0.27	14 (2%) 57 34	20, 40, 74, 125	0
1	B	531/574 (92%)	0.28	28 (5%) 32 16	14, 37, 94, 130	0
All	All	1072/1148 (93%)	0.28	42 (3%) 43 24	14, 39, 79, 130	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	556	SER	6.5
1	A	147	VAL	5.0
1	B	547	LEU	4.3
1	B	552	VAL	3.9
1	B	404	PRO	3.8
1	B	551	PHE	3.6
1	B	147	VAL	3.6
1	B	554	GLY	3.5
1	B	555	TYR	3.4
1	A	149	PRO	3.4
1	B	550	TRP	3.4
1	B	149	PRO	3.4
1	B	405	VAL	3.2
1	A	154	ARG	3.2
1	A	558	GLY	3.1
1	A	532	THR	3.1
1	B	451	CYS	3.0
1	A	148	GLN	3.0
1	B	545	LEU	2.9
1	A	564	LEU	2.9
1	B	151	LYS	2.7
1	B	543	SER	2.6
1	B	14	CYS	2.6
1	A	14	CYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	94	PRO	2.5
1	B	542	ALA	2.4
1	B	150	GLU	2.4
1	B	544	GLN	2.4
1	B	546	ASP	2.3
1	A	152	GLY	2.3
1	B	135	ASP	2.3
1	A	531	ARG	2.3
1	B	541	ALA	2.3
1	A	151	LYS	2.3
1	A	41	THR	2.3
1	B	403	THR	2.3
1	B	153	GLY	2.2
1	B	399	THR	2.2
1	B	553	ALA	2.2
1	A	535	LYS	2.1
1	A	543	SER	2.1
1	B	146	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	654	5/5	0.62	0.18	137,141,142,143	0
3	SO4	B	657	5/5	0.69	0.17	87,92,93,93	0
3	SO4	B	652	5/5	0.74	0.18	99,103,105,105	0
3	SO4	B	658	5/5	0.76	0.16	104,108,109,110	0
3	SO4	A	651	5/5	0.77	0.14	98,102,103,104	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	652	5/5	0.77	0.18	88,92,93,93	0
3	SO4	B	653	5/5	0.81	0.11	86,90,91,91	0
3	SO4	B	656	5/5	0.89	0.12	71,75,77,77	0
3	SO4	A	653	5/5	0.89	0.08	67,71,72,72	0
2	23E	B	601	47/47	0.91	0.10	14,33,44,55	0
2	23E	A	601	47/47	0.91	0.10	20,31,54,67	0
4	26F	B	602	42/42	0.93	0.10	22,40,61,64	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 23E B 601:

2mF_o-DF_c (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
 and green (positive)






Electron density around 23E A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 26F B 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.