



wwPDB EM Validation Summary Report ⓘ

Mar 9, 2026 – 05:43 AM UTC

PDB ID : 6QG6 / pdb_00006qg6
EMDB ID : EMD-4548
Title : Structure of eIF2B-eIF2 (phosphorylated at Ser51) complex (model D)
Authors : Llacer, J.L.; Gordiyenko, Y.; Ramakrishnan, V.
Deposited on : 2019-01-10
Resolution : 10.40 Å (reported)
Based on initial models : 6FYX, 5B04

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 10.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.













Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	109 (9.90 - 10.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	305	
1	B	305	
2	C	381	
2	D	381	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	578	
3	F	578	
4	G	651	
4	H	651	
5	I	712	
5	J	712	
6	K	304	
6	L	304	
7	M	527	
7	N	527	
8	O	285	
8	P	285	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 36982 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Translation initiation factor eIF-2B subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	302	Total	C	N	O	S	0	0
			2351	1504	394	443	10		
1	B	302	Total	C	N	O	S	0	0
			2351	1504	394	443	10		

- Molecule 2 is a protein called Translation initiation factor eIF-2B subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	345	Total	C	N	O	S	0	0
			2665	1694	463	502	6		
2	D	345	Total	C	N	O	S	0	0
			2665	1694	463	502	6		

- Molecule 3 is a protein called Translation initiation factor eIF-2B subunit gamma.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	E	267	Total	C	N	O	S	0	0
			2164	1391	363	400	10		
3	F	267	Total	C	N	O	S	0	0
			2164	1391	363	400	10		

- Molecule 4 is a protein called Translation initiation factor eIF-2B subunit delta.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	G	355	Total	C	N	O	S	0	0
			2744	1738	474	521	11		
4	H	355	Total	C	N	O	S	0	0
			2744	1738	474	521	11		

- Molecule 5 is a protein called Translation initiation factor eIF-2B subunit epsilon.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	I	431	Total	C	N	O	S	0	0
			3406	2147	573	666	20		
5	J	431	Total	C	N	O	S	0	0
			3406	2147	573	666	20		

- Molecule 6 is a protein called Eukaryotic translation initiation factor 2 subunit alpha.

Mol	Chain	Residues	Atoms						AltConf	Trace
6	K	246	Total	C	N	O	P	S	0	0
			1974	1259	324	382	1	8		
6	L	246	Total	C	N	O	P	S	0	0
			1974	1259	324	382	1	8		

- Molecule 7 is a protein called Eukaryotic translation initiation factor 2 subunit gamma.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	M	408	Total	C	N	O	S	0	0
			3044	1934	546	548	16		
7	N	408	Total	C	N	O	S	0	0
			3044	1934	546	548	16		


- Molecule 8 is a protein called Eukaryotic translation initiation factor 2 subunit beta.

Mol	Chain	Residues	Atoms				AltConf	Trace
8	O	17	Total	C	N	O	0	0
			143	96	24	23		
8	P	17	Total	C	N	O	0	0
			143	96	24	23		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Translation initiation factor eIF-2B subunit alpha




- Molecule 1: Translation initiation factor eIF-2B subunit alpha




- Molecule 2: Translation initiation factor eIF-2B subunit beta







• Molecule 2: Translation initiation factor eIF-2B subunit beta




• Molecule 3: Translation initiation factor eIF-2B subunit gamma







- Molecule 4: Translation initiation factor eIF-2B subunit delta




- Molecule 5: Translation initiation factor eIF-2B subunit epsilon






• Molecule 5: Translation initiation factor eIF-2B subunit epsilon




• Molecule 6: Eukaryotic translation initiation factor 2 subunit alpha






• Molecule 6: Eukaryotic translation initiation factor 2 subunit alpha



• Molecule 7: Eukaryotic translation initiation factor 2 subunit gamma



ARG
LEU
PHE
PHE
MET
VAL
CYS
LYS
SER
CYS
GLY
SER
THR
ARG
SER
VAL
SER
SER
ILE
LYS
THR
GLY
PHE
GLN
ALA
THR
VAL
GLY
LYS
ARG
ARG
MET

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	12575	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	45	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	3500	Depositor
Magnification	104478	Depositor
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.079	Depositor
Minimum map value	-0.021	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.0125	Depositor
Map size ($\text{\AA}$)	375.2, 375.2, 375.2	wwPDB
Map dimensions	280, 280, 280	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.34, 1.34, 1.34	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.59	0/2395	1.04	7/3251 (0.2%)
1	B	0.63	0/2395	1.02	4/3251 (0.1%)
2	C	0.62	0/2714	1.12	3/3693 (0.1%)
2	D	0.59	0/2714	1.05	1/3693 (0.0%)
3	E	0.66	0/2209	0.96	2/2989 (0.1%)
3	F	0.68	0/2209	0.98	1/2989 (0.0%)
4	G	0.61	0/2781	1.06	3/3747 (0.1%)
4	H	0.59	0/2781	1.09	3/3747 (0.1%)
5	I	0.63	0/3468	1.02	5/4704 (0.1%)
5	J	0.64	1/3468 (0.0%)	0.98	1/4704 (0.0%)
6	K	0.65	0/1989	1.03	1/2675 (0.0%)
6	L	0.66	0/1989	1.02	2/2675 (0.1%)
7	M	0.70	0/3087	0.97	2/4173 (0.0%)
7	N	0.70	0/3087	0.98	4/4173 (0.1%)
8	O	0.75	0/146	1.05	0/196
8	P	0.61	0/146	1.05	0/196
All	All	0.64	1/37578 (0.0%)	1.02	39/50856 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	J	242	VAL	CA-CB	5.42	1.56	1.54

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	404	GLY	N-CA-C	7.62	119.14	111.21
7	M	300	LYS	N-CA-C	7.41	119.05	110.97
5	I	57	VAL	CA-C-N	7.16	128.79	119.84
5	I	57	VAL	C-N-CA	7.16	128.79	119.84
7	N	121	SER	N-CA-C	7.12	114.62	108.78

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2351	0	2367	96	0
1	B	2351	0	2367	117	0
2	C	2665	0	2625	45	0
2	D	2665	0	2626	30	0
3	E	2164	0	2154	23	0
3	F	2164	0	2154	15	0
4	G	2744	0	2819	29	0
4	H	2744	0	2819	38	0
5	I	3406	0	3359	46	0
5	J	3406	0	3359	50	0
6	K	1974	0	2020	17	0
6	L	1974	0	2019	10	0
7	M	3044	0	3126	22	0
7	N	3044	0	3126	21	0
8	O	143	0	148	0	0
8	P	143	0	148	0	0
All	All	36982	0	37236	508	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 508 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:149:PHE:CE1	1:A:174:ILE:CD1	2.26	1.17
1:A:149:PHE:CE1	1:A:174:ILE:HD13	1.80	1.16
1:B:283:LEU:HB3	1:B:291:THR:OG1	1.42	1.16
1:A:149:PHE:HE1	1:A:174:ILE:CD1	1.58	1.14
1:B:291:THR:H	1:B:292:PRO:CD	1.65	1.09

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	300/305 (98%)	240 (80%)	52 (17%)	8 (3%)	4	25
1	B	300/305 (98%)	245 (82%)	44 (15%)	11 (4%)	2	20
2	C	339/381 (89%)	296 (87%)	36 (11%)	7 (2%)	5	30
2	D	339/381 (89%)	296 (87%)	33 (10%)	10 (3%)	3	23
3	E	253/578 (44%)	206 (81%)	41 (16%)	6 (2%)	4	27
3	F	253/578 (44%)	204 (81%)	42 (17%)	7 (3%)	4	24
4	G	349/651 (54%)	277 (79%)	58 (17%)	14 (4%)	2	18
4	H	349/651 (54%)	293 (84%)	41 (12%)	15 (4%)	2	17
5	I	427/712 (60%)	358 (84%)	51 (12%)	18 (4%)	2	17
5	J	427/712 (60%)	357 (84%)	49 (12%)	21 (5%)	1	16
6	K	237/304 (78%)	198 (84%)	33 (14%)	6 (2%)	4	26
6	L	237/304 (78%)	200 (84%)	34 (14%)	3 (1%)	9	42
7	M	398/527 (76%)	329 (83%)	61 (15%)	8 (2%)	6	31
7	N	398/527 (76%)	341 (86%)	52 (13%)	5 (1%)	9	42
8	O	15/285 (5%)	13 (87%)	1 (7%)	1 (7%)	1	12
8	P	15/285 (5%)	12 (80%)	2 (13%)	1 (7%)	1	12
All	All	4636/7486 (62%)	3865 (83%)	630 (14%)	141 (3%)	5	23

5 of 141 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	61	PRO
1	B	158	PRO
1	B	291	THR
2	C	276	SER
3	E	100	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/265 (97%)	215 (84%)	42 (16%)	2	10
1	B	257/265 (97%)	219 (85%)	38 (15%)	3	13
2	C	286/338 (85%)	245 (86%)	41 (14%)	3	13
2	D	286/338 (85%)	248 (87%)	38 (13%)	4	14
3	E	249/529 (47%)	233 (94%)	16 (6%)	16	37
3	F	249/529 (47%)	230 (92%)	19 (8%)	12	33
4	G	305/561 (54%)	277 (91%)	28 (9%)	8	27
4	H	305/561 (54%)	272 (89%)	33 (11%)	6	21
5	I	389/649 (60%)	353 (91%)	36 (9%)	8	26
5	J	389/649 (60%)	358 (92%)	31 (8%)	11	31
6	K	219/273 (80%)	184 (84%)	35 (16%)	2	11
6	L	219/273 (80%)	187 (85%)	32 (15%)	3	13
7	M	319/449 (71%)	302 (95%)	17 (5%)	20	41
7	N	319/449 (71%)	302 (95%)	17 (5%)	20	41
8	O	16/246 (6%)	16 (100%)	0	100	100
8	P	16/246 (6%)	16 (100%)	0	100	100
All	All	4080/6620 (62%)	3657 (90%)	423 (10%)	9	22

5 of 423 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	H	339	THR
5	I	345	ILE
7	M	402	VAL
4	H	429	VAL
5	I	106	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 96 such sidechains are listed below:

Mol	Chain	Res	Type
5	I	289	GLN
6	K	23	ASN
5	I	313	GLN
5	J	96	ASN
6	L	103	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	SEP	L	52	6	8,9,10	0.67	0	7,12,14	1.41	1 (14%)
6	SEP	K	52	6	8,9,10	0.63	0	7,12,14	1.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	SEP	L	52	6	-	2/6/8/10	-
6	SEP	K	52	6	-	2/6/8/10	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	L	52	SEP	OG-CB-CA	2.74	110.81	108.14

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	K	52	SEP	N-CA-CB-OG
6	K	52	SEP	C-CA-CB-OG
6	L	52	SEP	N-CA-CB-OG
6	L	52	SEP	C-CA-CB-OG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-4548. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 140



Y Index: 140




Z Index: 140


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 96



Y Index: 126




Z Index: 143


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0125. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1275 nm³; this corresponds to an approximate mass of 1151 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.096 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-4548 and PDB model 6QG6. Per-residue inclusion information can be found in section 3 on page 6.


9.1 Map-model overlay [i](#)



X



Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.0125 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0125).

9.4 Atom inclusion [i](#)



At the recommended contour level, 95% of all backbone atoms, 94% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.0125) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9370	 0.0680
A	 0.9780	 0.0750
B	 0.9470	 0.0500
C	 0.9600	 0.0750
D	 0.9550	 0.0730
E	 0.9330	 0.0530
F	 0.9520	 0.0620
G	 0.9290	 0.0800
H	 0.9330	 0.0830
I	 0.9810	 0.0750
J	 0.9730	 0.0820
K	 0.8810	 0.0640
L	 0.9780	 0.0780
M	 0.7610	 0.0470
N	 0.9780	 0.0470
O	 0.4270	 0.0780
P	 1.0000	 0.0400

