



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:29 PM UTC

PDB ID : 6QGC / pdb_00006qgc
Title : PETase from Ideonella sakaiensis without ligand
Authors : Palm, G.J.; Reisky, L.; Boettcher, D.; Mueller, H.; Michels, E.A.P.; Walczak, C.; Berndt, L.; Weiss, M.S.; Bornscheuer, U.T.; Weber, G.
Deposited on : 2019-01-10
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	290	
1	B	290	
1	C	290	
1	D	290	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CL	B	301	-	-	X	-
3	CL	D	301	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 8171 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Poly(ethylene terephthalate) hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	262	Total	C	N	O	S	0	0	0
			1926	1189	344	382	11			
1	B	261	Total	C	N	O	S	0	1	0
			1923	1189	343	380	11			
1	C	262	Total	C	N	O	S	0	1	0
			1927	1191	344	381	11			
1	D	261	Total	C	N	O	S	0	1	0
			1922	1188	343	380	11			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	B	1	Total	Cl	0	0
			1	1		
3	C	1	Total	Cl	0	0
			1	1		
3	D	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	105	Total	O	0	0
			105	105		
4	B	103	Total	O	0	0
			103	103		
4	C	120	Total	O	0	0
			120	120		
4	D	126	Total	O	0	0
			126	126		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Poly(ethylene terephthalate) hydrolase




- Molecule 1: Poly(ethylene terephthalate) hydrolase




- Molecule 1: Poly(ethylene terephthalate) hydrolase



- Molecule 1: Poly(ethylene terephthalate) hydrolase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.12Å 78.77Å 140.12Å 90.00° 92.56° 90.00°	Depositor
Resolution (Å)	48.73 – 2.00 48.73 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.4 (48.73-2.00) 97.4 (48.73-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0230	Depositor
R, R_{free}	0.213 , 0.261 0.221 , 0.264	Depositor DCC
R_{free} test set	3607 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	22.3	Xtriage
Anisotropy	0.512	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.130 for h,-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8171	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.25	7/1971 (0.4%)	1.26	3/2684 (0.1%)
1	B	1.28	11/1971 (0.6%)	1.28	4/2686 (0.1%)
1	C	1.24	8/1975 (0.4%)	1.27	10/2691 (0.4%)
1	D	1.29	10/1970 (0.5%)	1.30	7/2684 (0.3%)
All	All	1.26	36/7887 (0.5%)	1.28	24/10745 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	2
1	C	0	6
1	D	0	3
All	All	0	15

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	106	PHE	C-O	-7.12	1.15	1.23
1	D	104	HIS	CE1-NE2	6.80	1.39	1.32
1	A	161	MET	C-O	-6.79	1.16	1.24
1	D	221	SER	C-O	-6.46	1.15	1.24
1	A	30	ASN	N-CA	6.06	1.54	1.46
1	B	271	PHE	C-O	6.04	1.31	1.24
1	C	201	PHE	C-O	-6.01	1.16	1.24
1	B	209	ALA	N-CA	5.93	1.52	1.46
1	C	169	SER	C-O	5.81	1.30	1.24
1	C	121	SER	N-CA	5.78	1.53	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	81	ILE	C-O	-5.78	1.18	1.24
1	D	110	THR	N-CA	5.68	1.53	1.46
1	C	122	SER	C-O	5.58	1.30	1.24
1	B	154	MET	N-CA	5.57	1.52	1.45
1	B	115	SER	N-CA	5.56	1.52	1.46
1	A	44	GLU	C-O	-5.54	1.16	1.24
1	A	220	ASP	C-O	-5.43	1.17	1.24
1	D	194	VAL	N-CA	5.42	1.52	1.46
1	A	280	ARG	NE-CZ	5.40	1.39	1.33
1	B	262	MET	N-CA	5.34	1.53	1.46
1	D	65	ALA	C-O	-5.32	1.17	1.23
1	A	239	CYS	N-CA	5.28	1.53	1.46
1	B	73	ASN	C-O	-5.24	1.18	1.24
1	B	94	ILE	N-CA	5.24	1.51	1.46
1	A	86	GLY	N-CA	5.23	1.50	1.45
1	C	54	SER	C-O	-5.23	1.17	1.23
1	C	165	GLY	C-O	5.20	1.30	1.23
1	B	184	PRO	C-O	-5.18	1.17	1.23
1	C	110	THR	N-CA	5.18	1.52	1.46
1	B	99	PRO	C-O	-5.16	1.17	1.24
1	D	165	GLY	N-CA	5.13	1.52	1.45
1	D	64	GLY	N-CA	5.12	1.52	1.45
1	B	166	SER	N-CA	5.07	1.52	1.46
1	C	44	GLU	N-CA	5.07	1.52	1.46
1	B	137	LEU	N-CA	5.06	1.52	1.46
1	D	220	ASP	CG-OD2	5.04	1.34	1.25

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	263	ASP	CA-CB-CG	8.12	120.72	112.60
1	C	30	ASN	CB-CA-C	7.87	117.96	110.17
1	D	220	ASP	CA-CB-CG	6.75	119.35	112.60
1	D	270	THR	CA-CB-OG1	-6.28	100.19	109.60
1	B	154	MET	CA-C-N	6.15	126.87	121.58
1	B	154	MET	C-N-CA	6.15	126.87	121.58
1	A	280	ARG	NE-CZ-NH2	6.13	124.72	119.20
1	D	154	MET	CA-C-N	5.86	127.03	121.46
1	D	154	MET	C-N-CA	5.86	127.03	121.46
1	C	135	ALA	CA-C-N	5.57	127.74	120.28
1	C	135	ALA	C-N-CA	5.57	127.74	120.28
1	B	220	ASP	CA-CB-CG	5.45	118.05	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	86	GLY	CA-C-N	5.41	128.79	121.05
1	C	86	GLY	C-N-CA	5.41	128.79	121.05
1	C	163	GLY	CA-C-N	5.30	125.98	119.99
1	C	163	GLY	C-N-CA	5.30	125.98	119.99
1	B	261	PHE	CB-CA-C	5.28	118.48	109.24
1	C	195	THR	N-CA-C	-5.24	106.94	113.38
1	A	51	THR	CA-CB-OG1	-5.11	101.94	109.60
1	C	120	PRO	N-CA-C	5.10	120.53	113.65
1	D	197	PRO	CB-CA-C	-5.09	105.09	111.71
1	C	279	THR	CA-CB-OG1	-5.07	102.00	109.60
1	D	132	ARG	NE-CZ-NH1	-5.05	116.45	121.50
1	A	184	PRO	N-CA-C	5.01	119.18	111.21

There are no chirality outliers.

All (15) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	123	ARG	Sidechain
1	A	132	ARG	Sidechain
1	A	285	ARG	Sidechain
1	A	53	ARG	Sidechain
1	B	123	ARG	Sidechain
1	B	280	ARG	Sidechain
1	C	123	ARG	Sidechain
1	C	224	ARG	Sidechain
1	C	280	ARG	Sidechain
1	C	285	ARG	Sidechain
1	C	59	ARG	Sidechain
1	C	90	ARG	Sidechain
1	D	132	ARG	Sidechain
1	D	267	ARG	Sidechain
1	D	34	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1926	0	1852	19	0
1	B	1923	0	1854	12	0
1	C	1927	0	1854	13	0
1	D	1922	0	1852	7	0
2	A	5	0	0	0	0
2	C	10	0	0	0	0
3	A	1	0	0	1	0
3	B	1	0	0	2	0
3	C	1	0	0	0	0
3	D	1	0	0	2	0
4	A	105	0	0	0	0
4	B	103	0	0	2	0
4	C	120	0	0	4	0
4	D	126	0	0	4	0
All	All	8171	0	7412	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246:ASN:OD1	1:A:280:ARG:HD2	1.76	0.85
1:D:246:ASN:OD1	1:D:280:ARG:HD2	1.85	0.76
1:A:145:ILE:HA	1:A:148:LYS:HD2	1.73	0.70
3:B:301:CL:CL	4:B:450:HOH:O	2.46	0.70
1:B:246:ASN:OD1	1:B:280:ARG:HD2	1.93	0.67
1:B:113[B]:THR:HG23	1:B:126:GLN:NE2	2.12	0.65
3:D:301:CL:CL	4:D:428:HOH:O	2.51	0.64
1:B:113[B]:THR:HG23	1:B:126:GLN:HE22	1.67	0.59
1:A:246:ASN:OD1	1:A:280:ARG:CD	2.52	0.58
1:D:161:MET:HG3	3:D:301:CL:CL	2.43	0.56
1:A:216:LEU:HD11	1:A:285:ARG:HH21	1.70	0.56
1:A:230:LEU:C	1:A:230:LEU:HD23	2.29	0.56
1:C:83:ILE:HG12	1:C:157:MET:HB2	1.87	0.55
1:B:113[B]:THR:CG2	1:B:118:ASP:OD2	2.55	0.55
1:B:161:MET:HG3	3:B:301:CL:CL	2.45	0.54
1:B:113[B]:THR:HG22	1:B:118:ASP:OD2	2.09	0.53
1:C:246:ASN:OD1	1:C:280:ARG:HD3	2.08	0.53
1:A:203:CYS:O	1:A:204:GLU:C	2.52	0.51
1:A:219:TYR:CD1	1:A:229:PHE:HB2	2.46	0.50
1:B:181:PRO:HD2	1:B:199:LEU:O	2.12	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:183:ALA:N	1:C:184:PRO:CD	2.75	0.49
1:A:180:ALA:HA	1:A:199:LEU:O	2.13	0.49
1:D:119:GLN:HB3	1:D:120:PRO:CD	2.42	0.49
1:C:180:ALA:HA	1:C:199:LEU:O	2.12	0.49
1:C:177:LYS:HG2	4:C:495:HOH:O	2.13	0.47
1:A:119:GLN:HB3	1:A:120:PRO:CD	2.45	0.47
1:C:279:THR:HG22	4:C:496:HOH:O	2.15	0.46
1:B:113[B]:THR:HG21	1:B:118:ASP:HB2	1.98	0.46
1:C:80:ALA:HA	1:C:107:VAL:O	2.16	0.46
1:A:88:THR:HA	1:A:116:THR:HB	1.97	0.46
1:C:223:SER:HB2	4:C:471:HOH:O	2.17	0.45
1:D:119:GLN:HB3	1:D:120:PRO:HD2	1.99	0.45
1:A:161:MET:HG3	3:A:302:CL:CL	2.54	0.45
1:A:258:MET:O	1:A:262:MET:HB2	2.18	0.44
1:A:197:PRO:HA	1:A:226:ALA:O	2.17	0.44
1:C:95:LYS:HE2	4:C:474:HOH:O	2.18	0.43
1:B:219:TYR:CE1	1:B:227:LYS:HG3	2.54	0.43
1:A:145:ILE:O	1:A:146:TYR:C	2.61	0.43
1:B:100:ARG:NH1	4:B:402:HOH:O	2.46	0.43
1:A:216:LEU:N	1:A:217:PRO:HD2	2.34	0.42
1:A:210:PRO:HB2	1:A:212:ASN:OD1	2.19	0.42
1:B:160:SER:HA	1:B:183:ALA:O	2.20	0.42
1:B:180:ALA:HA	1:B:199:LEU:O	2.20	0.41
1:C:141:SER:HA	1:C:146:TYR:CG	2.55	0.41
1:C:182:GLN:HA	1:C:201:PHE:HB2	2.02	0.41
1:A:230:LEU:HD21	1:A:281:VAL:HG22	2.01	0.41
1:D:65:ALA:HB2	4:D:423:HOH:O	2.20	0.41
1:A:38:PRO:HB3	1:A:268:TYR:CZ	2.55	0.41
1:D:270:THR:HB	4:D:478:HOH:O	2.20	0.41
1:A:51:THR:O	1:A:71:PRO:HA	2.20	0.41
1:D:121:SER:HB2	4:D:494:HOH:O	2.21	0.41
1:C:206:ASP:CG	1:C:237:HIS:HD1	2.29	0.40
1:C:203:CYS:O	1:C:204:GLU:C	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/290 (90%)	254 (98%)	6 (2%)	0	100	100
1	B	260/290 (90%)	255 (98%)	5 (2%)	0	100	100
1	C	261/290 (90%)	252 (97%)	9 (3%)	0	100	100
1	D	260/290 (90%)	253 (97%)	7 (3%)	0	100	100
All	All	1041/1160 (90%)	1014 (97%)	27 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	208/226 (92%)	206 (99%)	2 (1%)	68	75
1	B	208/226 (92%)	204 (98%)	4 (2%)	50	56
1	C	208/226 (92%)	206 (99%)	2 (1%)	68	75
1	D	208/226 (92%)	206 (99%)	2 (1%)	68	75
All	All	832/904 (92%)	822 (99%)	10 (1%)	63	70

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	94	ILE
1	A	125	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	61	SER
1	B	95	LYS
1	B	115	SER
1	B	214	SER
1	C	125	SER
1	C	185	TRP
1	D	94	ILE
1	D	160	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	119	GLN
1	A	288	ASN
1	B	91	GLN
1	B	233	ASN
1	B	264	ASN
1	B	288	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	301	-	4,4,4	0.30	0	6,6,6	0.66	0
2	SO4	C	301	-	4,4,4	0.43	0	6,6,6	0.61	0
2	SO4	C	302	-	4,4,4	0.50	0	6,6,6	0.20	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	262/290 (90%)	0.28	2 (0%) 82 82	16, 24, 31, 40	0
1	B	261/290 (90%)	0.39	4 (1%) 72 71	13, 24, 34, 45	1 (0%)
1	C	262/290 (90%)	0.45	4 (1%) 72 71	15, 24, 35, 51	1 (0%)
1	D	261/290 (90%)	0.28	2 (0%) 82 82	12, 23, 30, 41	1 (0%)
All	All	1046/1160 (90%)	0.35	12 (1%) 78 77	12, 24, 33, 51	3 (0%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	280	ARG	2.9
1	D	29	THR	2.7
1	B	284	PHE	2.7
1	C	273	CYS	2.6
1	B	29	THR	2.6
1	B	289	CYS	2.5
1	C	289	CYS	2.3
1	A	29	THR	2.2
1	C	248	ALA	2.2
1	D	280	ARG	2.1
1	A	37	ASN	2.0
1	C	29	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	C	302	5/5	0.84	0.12	56,61,65,66	0
2	SO4	A	301	5/5	0.95	0.09	27,31,37,37	0
3	CL	B	301	1/1	0.95	0.07	25,25,25,25	0
3	CL	A	302	1/1	0.96	0.05	20,20,20,20	0
2	SO4	C	301	5/5	0.96	0.09	31,34,38,41	0
3	CL	C	303	1/1	0.97	0.09	21,21,21,21	0
3	CL	D	301	1/1	0.98	0.05	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.