



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 04:07 AM UTC

PDB ID : 6QGT / pdb_00006qgt
Title : The carbon monoxide inhibition of F420-reducing [NiFe] hydrogenase complex from *Methanosarcina barkeri*
Authors : Ilina, Y.; Lorent, C.; Katz, S.; Jeoung, J.H.; Shima, S.; Horch, M.; Zebger, I.; Dobbek, H.
Deposited on : 2019-01-12
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	253	 91% 9%
2	A	438	 92% 8%
3	B	291	 93% 6%

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 8237 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Coenzyme F420 hydrogenase subunit gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	252	Total	C	N	O	S	0	0	0
			1911	1200	322	366	23			

- Molecule 2 is a protein called F420-reducing hydrogenase, subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	437	Total	C	N	O	S	0	3	0
			3421	2154	609	631	27			

- Molecule 3 is a protein called Coenzyme F420 hydrogenase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	B	291	Total	C	N	O	S	0	1	0
			2271	1442	390	425	14			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	265	GLN	GLU	conflict	UNP A0A0E3QWH3

- Molecule 4 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	G	1	Total	Fe	S	0	0
			8	4	4		
4	G	1	Total	Fe	S	0	0
			8	4	4		
4	G	1	Total	Fe	S	0	0
			8	4	4		
4	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 5 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe_2S_2).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	G	1	Total	Fe	S	0	0
			3	1	2		

- Molecule 6 is TRIS-HYDROXYMETHYL-METHYL-AMMONIUM (CCD ID: 144) (formula: $C_4H_{12}NO_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	G	1	Total	C	N	O	0	0
			8	4	1	3		
6	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 7 is (R,R)-2,3-BUTANEDIOL (CCD ID: BU3) (formula: $C_4H_{10}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	G	1	Total	C	O	0	0
			6	4	2		
7	G	1	Total	C	O	0	0
			6	4	2		
7	B	1	Total	C	O	0	0
			6	4	2		

- Molecule 8 is dicyano-(oxidaniumylidynemethylnickelio)-(oxidanylidene)iron (CCD ID: J52) (formula: $C_4FeN_2NiO_2$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
8	A	1	Total	C	Fe	N	Ni	O	0	0
			10	4	1	2	1	2		


- Molecule 9 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total	Fe	0	0
			1	1		

- Molecule 10 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	1	Total	Mg	0	0
			1	1		

- Molecule 11 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 12 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 13 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
13	G	102	Total O 104 104	0	2
13	A	257	Total O 260 260	0	3
13	B	127	Total O 128 128	0	1

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Coenzyme F420 hydrogenase subunit gamma

Chain G:  91% 9%



- Molecule 2: F420-reducing hydrogenase, subunit alpha

Chain A:  92% 8%



- Molecule 3: Coenzyme F420 hydrogenase subunit beta

Chain B:  93% 6%



4 Data and refinement statistics

Property	Value	Source
Space group	F 2 3	Depositor
Cell constants a, b, c, α , β , γ	235.43Å 235.43Å 235.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.06 – 1.99 48.06 – 1.99	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.06-1.99) 99.9 (48.06-1.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.96 (at 1.98Å)	Xtriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, R_{free}	0.169 , 0.209 0.173 , 0.213	Depositor DCC
R_{free} test set	2100 reflections (2.84%)	wwPDB-VP
Wilson B-factor (Å ²)	39.7	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 40.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.034 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8237	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 144, J52, FAD, MPD, SF4, MG, FES, BU3, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	G	0.49	0/1943	0.66	0/2626
2	A	0.49	0/3502	0.64	0/4757
3	B	0.40	0/2315	0.58	0/3125
All	All	0.47	0/7760	0.63	0/10508

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	1911	0	1893	19	0
2	A	3421	0	3394	20	0
3	B	2271	0	2305	10	0
4	B	8	0	0	0	0
4	G	24	0	0	1	0
5	G	3	0	0	0	0
6	A	8	0	12	0	0
6	G	8	0	11	3	0
7	B	6	0	10	0	0
7	G	12	0	20	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	A	10	0	0	2	0
9	A	1	0	0	0	0
10	A	1	0	0	0	0
11	A	8	0	14	0	0
12	B	53	0	30	1	0
13	A	260	0	0	1	0
13	B	128	0	0	1	0
13	G	104	0	0	2	0
All	All	8237	0	7689	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:153:ARG:HH12	7:G:306:BU3:H43	1.50	0.77
2:A:348[B]:ASN:ND2	13:A:601:HOH:O	2.23	0.71
1:G:89:ASP:OD1	6:G:305:144:H13	1.93	0.69
1:G:153:ARG:HH12	7:G:306:BU3:C4	2.07	0.68
3:B:263:ALA:O	3:B:267:ILE:HG12	1.96	0.64
6:G:305:144:H42	2:A:9:PRO:HD3	1.81	0.62
2:A:148:ILE:HG23	2:A:186:LEU:HB3	1.82	0.60
2:A:148:ILE:HG21	2:A:190:CYS:HB2	1.84	0.59
2:A:301:VAL:HG13	2:A:408:PRO:HG3	1.85	0.58
2:A:62:VAL:HG21	2:A:399:MET:HE1	1.86	0.57
3:B:137:PHE:CD2	3:B:213:GLY:HA3	2.40	0.56
3:B:199:VAL:HG22	3:B:274:VAL:HG22	1.88	0.54
3:B:146:SER:HB3	3:B:192:TYR:HB3	1.90	0.51
3:B:250:GLU:H	3:B:250:GLU:CD	2.16	0.51
2:A:14:GLU:HB3	2:A:432:CYS:HA	1.93	0.50
3:B:39:TYR:OH	3:B:237:LYS:HG2	2.12	0.50
1:G:22:LYS:NZ	13:G:406:HOH:O	2.43	0.49
1:G:142:ILE:HG23	1:G:151:LEU:HD23	1.94	0.49
1:G:65:ARG:O	6:G:305:144:H41	2.12	0.49
2:A:119:ILE:HD12	2:A:124:LYS:HD3	1.94	0.49
3:B:138:CYS:O	3:B:266:LYS:NZ	2.45	0.47
2:A:280:PRO:HA	2:A:281:ILE:HA	1.66	0.46
1:G:206:MET:HE1	1:G:245:PRO:HG2	1.97	0.46
1:G:266:LYS:HB3	1:G:266:LYS:HE2	1.56	0.45
1:G:24:GLY:HA3	1:G:70:MET:SD	2.56	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:153:ARG:NH1	7:G:306:BU3:H43	2.26	0.45
4:G:301:SF4:S4	2:A:61:ARG:HG2	2.57	0.45
2:A:402:PRO:HG2	8:A:501:J52:C2	2.46	0.44
2:A:432:CYS:HB3	2:A:435:CYS:HB2	1.99	0.44
3:B:9:LYS:HB2	3:B:9:LYS:HE2	1.64	0.44
2:A:97:LEU:HD13	2:A:157:GLY:HA3	1.99	0.44
2:A:328:THR:HG23	2:A:428:ALA:HB2	2.00	0.43
1:G:27:HIS:CD2	13:G:429:HOH:O	2.71	0.43
1:G:257:PHE:HA	3:B:118:LEU:HD11	2.01	0.43
3:B:250:GLU:HA	13:B:489:HOH:O	2.19	0.43
1:G:80:CYS:HB2	1:G:129:TYR:CE2	2.54	0.42
1:G:206:MET:CE	1:G:245:PRO:HG2	2.49	0.42
1:G:252:ASP:OD1	1:G:252:ASP:N	2.53	0.42
1:G:34:CYS:SG	1:G:104:GLY:HA3	2.59	0.42
2:A:317:LEU:HB3	2:A:323:PHE:HB2	2.02	0.42
2:A:430:ASP:N	2:A:431:PRO:HD3	2.35	0.42
12:B:302:FAD:H9	12:B:302:FAD:H1'1	1.85	0.42
2:A:308:PRO:HB3	2:A:408:PRO:HG2	2.02	0.41
1:G:236:LYS:HB3	1:G:236:LYS:HE3	1.88	0.41
2:A:380:ARG:CZ	8:A:501:J52:O	2.70	0.40
2:A:34:TRP:CH2	2:A:431:PRO:HB3	2.56	0.40
1:G:60:THR:O	2:A:12:ARG:HA	2.21	0.40
1:G:235:ILE:H	1:G:235:ILE:HG13	1.73	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	250/253 (99%)	238 (95%)	11 (4%)	1 (0%)	30	20
2	A	437/438 (100%)	430 (98%)	7 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	B	290/291 (100%)	280 (97%)	10 (3%)	0	100	100
All	All	977/982 (100%)	948 (97%)	28 (3%)	1 (0%)	48	41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	58	CYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	214/215 (100%)	212 (99%)	2 (1%)	70	67
2	A	370/368 (100%)	366 (99%)	4 (1%)	65	61
3	B	247/246 (100%)	240 (97%)	7 (3%)	38	29
All	All	831/829 (100%)	818 (98%)	13 (2%)	55	50

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	69	GLU
1	G	206	MET
2	A	13	LEU
2	A	104	HIS
2	A	225	LYS
2	A	309	VAL
3	B	42	GLU
3	B	140	GLU
3	B	199	VAL
3	B	250	GLU
3	B	272	LYS
3	B	275	GLU
3	B	287	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	122	ASN
2	A	232	GLN
2	A	376	ASN
3	B	232	ASN
3	B	271	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 2 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	144	G	305	-	1,7,7	1.26	0	3,9,9	0.38	0
11	MPD	A	505	-	7,7,7	0.31	0	9,10,10	0.73	0
12	FAD	B	302	-	58,58,58	3.22	26 (44%)	85,89,89	2.58	25 (29%)
5	FES	G	304	1	0,2,4	-	-	-	-	-
7	BU3	G	306	-	4,5,5	0.81	0	6,6,6	0.89	0
6	144	A	504	-	1,7,7	5.02	1 (100%)	3,9,9	0.22	0
8	J52	A	501	2	1,9,9	0.16	0	-	-	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SF4	G	302	1	0,12,12	-	-	-		
4	SF4	G	303	1	0,12,12	-	-	-		
4	SF4	B	303	3	0,12,12	-	-	-		
7	BU3	B	301	-	4,5,5	1.01	0	6,6,6	0.51	0
4	SF4	G	301	1	0,12,12	-	-	-		
7	BU3	G	307	-	4,5,5	0.78	0	6,6,6	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	144	G	305	-	-	0/0/9/9	-
12	FAD	B	302	-	-	4/34/50/50	0/6/6/6
7	BU3	G	306	-	-	0/4/4/4	-
6	144	A	504	-	-	0/0/9/9	-
4	SF4	G	302	1	-	-	0/6/5/5
4	SF4	G	303	1	-	-	0/6/5/5
4	SF4	G	301	1	-	-	0/6/5/5
4	SF4	B	303	3	-	-	0/6/5/5
7	BU3	B	301	-	-	4/4/4/4	-
11	MPD	A	505	-	-	0/5/5/5	-
7	BU3	G	307	-	-	4/4/4/4	-

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	B	302	FAD	P-O3P	8.57	1.68	1.59
12	B	302	FAD	C8A-N7A	7.85	1.46	1.31
12	B	302	FAD	O4B-C1B	7.63	1.59	1.42
12	B	302	FAD	C4X-N5	7.02	1.45	1.30
12	B	302	FAD	C10-N1	5.32	1.44	1.33
12	B	302	FAD	C6A-N6A	5.12	1.47	1.34
12	B	302	FAD	C4A-N3A	5.04	1.43	1.34
6	A	504	144	C1-N	-5.02	1.30	1.50
12	B	302	FAD	C2B-C1B	-4.69	1.38	1.53
12	B	302	FAD	PA-O3P	-4.49	1.54	1.59
12	B	302	FAD	C7M-C7	4.46	1.59	1.51
12	B	302	FAD	O4B-C4B	-4.42	1.35	1.45
12	B	302	FAD	O4-C4	-4.40	1.15	1.23
12	B	302	FAD	C5A-N7A	4.12	1.46	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	B	302	FAD	C9-C9A	4.06	1.46	1.39
12	B	302	FAD	C9-C8	-3.66	1.34	1.39
12	B	302	FAD	O2B-C2B	3.52	1.51	1.43
12	B	302	FAD	C2-N1	3.47	1.44	1.36
12	B	302	FAD	C5X-N5	3.26	1.45	1.39
12	B	302	FAD	C2A-N3A	3.00	1.39	1.33
12	B	302	FAD	C8M-C8	2.95	1.56	1.51
12	B	302	FAD	C5'-C4'	2.78	1.55	1.51
12	B	302	FAD	C4-N3	2.58	1.43	1.38
12	B	302	FAD	O3B-C3B	-2.56	1.36	1.43
12	B	302	FAD	O2'-C2'	-2.54	1.38	1.43
12	B	302	FAD	C2A-N1A	2.38	1.38	1.33
12	B	302	FAD	C10-N10	2.30	1.42	1.37

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	B	302	FAD	C7M-C7-C6	-10.13	101.72	119.57
12	B	302	FAD	C7M-C7-C8	7.93	136.95	120.76
12	B	302	FAD	N3A-C2A-N1A	-6.69	118.46	128.58
12	B	302	FAD	N6A-C6A-N1A	-6.06	104.88	118.38
12	B	302	FAD	C5A-C4A-N3A	-5.88	118.62	126.72
12	B	302	FAD	N9A-C8A-N7A	-5.87	105.61	113.94
12	B	302	FAD	C4A-N9A-C8A	5.11	111.11	105.74
12	B	302	FAD	C9-C9A-N10	-3.96	116.53	121.85
12	B	302	FAD	C5A-C6A-N6A	3.85	132.82	123.29
12	B	302	FAD	C2A-N3A-C4A	3.84	121.20	111.83
12	B	302	FAD	C4X-C10-N10	3.74	121.83	116.48
12	B	302	FAD	O4B-C1B-N9A	-3.56	101.25	108.09
12	B	302	FAD	N3A-C4A-N9A	3.30	132.78	127.17
12	B	302	FAD	C5X-N5-C4X	-2.96	113.30	118.09
12	B	302	FAD	O4-C4-C4X	-2.92	118.82	126.53
12	B	302	FAD	C5X-C9A-N10	2.88	120.58	117.97
12	B	302	FAD	C4X-C4-N3	2.62	119.92	113.25
12	B	302	FAD	C5A-C4A-N9A	2.56	108.60	105.81
12	B	302	FAD	C4A-C5A-N7A	-2.44	107.79	110.58
12	B	302	FAD	C8M-C8-C7	-2.26	116.14	120.76
12	B	302	FAD	C9A-N10-C10	-2.20	117.40	120.75
12	B	302	FAD	C5A-N7A-C8A	2.19	106.89	103.45
12	B	302	FAD	C4-N3-C2	-2.16	121.81	125.64
12	B	302	FAD	C4A-N9A-C1B	-2.11	121.69	126.63
12	B	302	FAD	O3B-C3B-C2B	-2.03	105.30	111.82

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	G	307	BU3	O5-C2-C3-O6
7	G	307	BU3	C1-C2-C3-O6
7	G	307	BU3	O5-C2-C3-C4
7	G	307	BU3	C1-C2-C3-C4
7	B	301	BU3	O5-C2-C3-O6
7	B	301	BU3	C1-C2-C3-O6
7	B	301	BU3	O5-C2-C3-C4
12	B	302	FAD	PA-O3P-P-O2P
7	B	301	BU3	C1-C2-C3-C4
12	B	302	FAD	O4B-C4B-C5B-O5B
12	B	302	FAD	PA-O3P-P-O1P
12	B	302	FAD	C3B-C4B-C5B-O5B

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	G	305	144	3	0
12	B	302	FAD	1	0
7	G	306	BU3	3	0
8	A	501	J52	2	0
4	G	301	SF4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	G	252/253 (99%)	-0.10	0 100 100	31, 41, 60, 94	0
2	A	437/438 (99%)	-0.38	0 100 100	15, 36, 49, 61	3 (0%)
3	B	291/291 (100%)	-0.03	0 100 100	21, 44, 62, 74	1 (0%)
All	All	980/982 (99%)	-0.21	0 100 100	15, 39, 58, 94	4 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	144	A	504	8/8	0.81	0.16	46,46,47,50	8
7	BU3	G	306	6/6	0.81	0.17	42,51,54,55	6
7	BU3	B	301	6/6	0.81	0.19	43,50,53,54	6
7	BU3	G	307	6/6	0.83	0.15	52,56,64,70	0
11	MPD	A	505	8/8	0.90	0.17	44,48,52,54	8

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	144	G	305	8/8	0.93	0.08	39,46,50,56	0
12	FAD	B	302	53/53	0.97	0.06	31,38,43,45	0
10	MG	A	503	1/1	0.98	0.04	37,37,37,37	1
4	SF4	G	302	8/8	0.98	0.04	32,35,36,38	0
4	SF4	G	301	8/8	0.98	0.04	35,35,36,37	0
8	J52	A	501	10/10	0.99	0.05	31,33,37,37	2
5	FES	G	304	3/4	0.99	0.04	35,35,38,43	2
4	SF4	G	303	8/8	0.99	0.04	36,37,39,39	0
4	SF4	B	303	8/8	0.99	0.04	38,40,41,42	0
9	FE	A	502	1/1	1.00	0.01	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.