



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 08:26 PM UTC

PDB ID : 1QHH / pdb_00001qhh
Title : STRUCTURE OF DNA HELICASE WITH ADPNP
Authors : Soultanas, P.; Dillingham, M.S.; Velankar, S.S.; Wigley, D.B.
Deposited on : 1999-05-14
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	167	2% 62% 32% ...
2	B	273	3% 58% 29% 7% ..
3	C	115	2% 63% 20% 5% 11%
4	D	169	2% 38% 15% .. 43%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ATP	A	700	X	-	-	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5291 atoms, of which 114 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (PCRA (SUBUNIT)).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	164	Total	C	H	N	O	S	0	0	0
			1330	823	28	238	234	7			

- Molecule 2 is a protein called PROTEIN (PCRA (SUBUNIT)).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	261	Total	C	H	N	O	S	0	0	0
			2250	1390	62	388	405	5			


- Molecule 3 is a protein called PROTEIN (PCRA (SUBUNIT)).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	102	Total	C	H	N	O	S	0	0	0
			817	510	14	129	162	2			

- Molecule 4 is a protein called PROTEIN (PCRA (SUBUNIT)).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	D	96	Total	C	H	N	O	S	0	0	0
			769	477	10	129	147	6			

- Molecule 5 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	32	Total	O	0	0
			32	32		
6	B	46	Total	O	0	0
			46	46		
6	C	3	Total	O	0	0
			3	3		
6	D	13	Total	O	0	0
			13	13		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PROTEIN (PCRA (SUBUNIT))



- Molecule 2: PROTEIN (PCRA (SUBUNIT))




- Molecule 3: PROTEIN (PCRA (SUBUNIT))



- Molecule 4: PROTEIN (PCRA (SUBUNIT))





4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	138.94Å 138.94Å 111.64Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 2.50 10.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	92.5 (10.00-2.50) 89.2 (10.00-2.50)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.00 (at 2.50Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.228 , 0.285 0.209 , 0.256	Depositor DCC
R_{free} test set	1862 reflections (4.42%)	wwPDB-VP
Wilson B-factor (Å ²)	45.7	Xtriage
Anisotropy	0.321	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 74.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.038 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5291	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	5/1323 (0.4%)	1.81	26/1783 (1.5%)
2	B	0.97	6/2231 (0.3%)	1.78	51/3016 (1.7%)
3	C	0.80	1/812 (0.1%)	1.65	7/1094 (0.6%)
4	D	1.03	4/772 (0.5%)	1.77	15/1042 (1.4%)
All	All	0.96	16/5138 (0.3%)	1.77	99/6935 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	302	HIS	CD2-NE2	-6.86	1.30	1.37
1	A	134	ILE	CA-CB	6.73	1.63	1.54
3	C	521	HIS	CD2-NE2	-6.62	1.30	1.37
4	D	644	HIS	CD2-NE2	-6.52	1.30	1.37
2	B	211	HIS	CD2-NE2	-6.05	1.31	1.37

The worst 5 of 99 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	425	SER	N-CA-C	15.52	131.49	109.15
4	D	639	ASN	CA-CB-CG	11.61	124.21	112.60
4	D	591	LEU	O-C-N	11.09	135.62	122.20
1	A	3	PHE	CA-CB-CG	9.83	123.63	113.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	388	ASP	CA-CB-CG	9.38	121.98	112.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	135	ASP	Peptide
1	A	73	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1302	28	1356	21	0
2	B	2188	62	2168	46	0
3	C	803	14	800	9	0
4	D	759	10	744	17	0
5	A	31	0	11	2	0
6	A	32	0	0	1	0
6	B	46	0	0	1	0
6	C	3	0	0	0	0
6	D	13	0	0	1	0
All	All	5177	114	5079	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 72 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:289:THR:HG21	6:B:731:HOH:O	1.87	0.75
2:B:173:VAL:HA	2:B:176:VAL:HG12	1.69	0.73
4:D:639:ASN:HB3	6:D:779:HOH:O	1.96	0.66
2:B:328:ASN:HA	4:D:622:MET:O	1.96	0.65
2:B:291:ARG:HG3	4:D:645:LEU:HD22	1.80	0.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	162/167 (97%)	149 (92%)	10 (6%)	3 (2%)	6	11
2	B	259/273 (95%)	248 (96%)	7 (3%)	4 (2%)	8	16
3	C	100/115 (87%)	89 (89%)	8 (8%)	3 (3%)	3	5
4	D	94/169 (56%)	86 (92%)	6 (6%)	2 (2%)	5	9
All	All	615/724 (85%)	572 (93%)	31 (5%)	12 (2%)	6	10

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	135	ASP
2	B	424	ALA
3	C	453	LEU
3	C	455	ALA
1	A	138	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/143 (99%)	128 (91%)	13 (9%)	8	19
2	B	233/243 (96%)	207 (89%)	26 (11%)	6	12
3	C	86/96 (90%)	70 (81%)	16 (19%)	1	3

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	D	82/136 (60%)	73 (89%)	9 (11%)	6	13
All	All	542/618 (88%)	478 (88%)	64 (12%)	5	11

5 of 64 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	588	ASN
4	D	606	VAL
2	B	265	GLN
2	B	242	ARG
4	D	610	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	416	ASN
3	C	511	ASN
4	D	644	HIS
4	D	587	HIS
2	B	232	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ATP	A	700	-	32,33,33	2.70	11 (34%)	48,52,52	3.17	17 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ATP	A	700	-	1/1/7/7	6/22/38/38	0/3/3/3

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	700	ATP	PA-O3A	-8.17	1.50	1.59
5	A	700	ATP	PB-O3A	-8.17	1.50	1.59
5	A	700	ATP	C6-N6	4.90	1.46	1.34
5	A	700	ATP	C8-N7	3.91	1.39	1.31
5	A	700	ATP	C5-C4	-3.43	1.33	1.39

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	700	ATP	C6-C5-C4	10.52	131.55	117.18
5	A	700	ATP	C5-C4-N3	-7.72	116.09	126.72
5	A	700	ATP	C6-C5-N7	-7.71	117.22	132.09
5	A	700	ATP	O4'-C1'-N9	5.96	119.53	108.09
5	A	700	ATP	O3G-PG-O3B	5.54	123.23	104.64

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	A	700	ATP	C3'

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	700	ATP	C5'-O5'-PA-O1A
5	A	700	ATP	C5'-O5'-PA-O2A
5	A	700	ATP	C5'-O5'-PA-O3A
5	A	700	ATP	C3'-C4'-C5'-O5'
5	A	700	ATP	PB-O3A-PA-O1A

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	700	ATP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	164/167 (98%)	-0.26	3 (1%) 67 64	14, 34, 77, 91	0
2	B	261/273 (95%)	-0.35	9 (3%) 48 43	14, 32, 72, 100	0
3	C	102/115 (88%)	0.09	2 (1%) 65 61	20, 51, 81, 90	0
4	D	96/169 (56%)	-0.51	3 (3%) 51 47	15, 27, 64, 85	0
All	All	623/724 (86%)	-0.28	17 (2%) 56 51	14, 34, 78, 100	0

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	650	SER	5.1
1	A	134	ILE	4.2
4	D	649	ALA	3.9
2	B	421	GLY	3.7
2	B	420	ARG	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	ATP	A	700	31/31	0.83	0.12	2,11,40,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.